



## Types, Frames, and Applicative Structures

# Def.: Types



Let  $\mathcal{T}$  be the least set s.t:

$$o \in \mathcal{T}$$

$$\iota \in \mathcal{T}$$

$$\forall \alpha, \beta \in \mathcal{T} : (\alpha\beta) \in \mathcal{T}$$

# Def.: Types



Let  $\mathcal{T}$  be the least set s.t:

$$o \in \mathcal{T}$$

$$\iota \in \mathcal{T}$$

$$\forall \alpha, \beta \in \mathcal{T} : (\alpha\beta) \in \mathcal{T}$$

We say that  $\alpha \in \mathcal{T}$  is a **simple type** (or type).  
 $(\alpha\beta)$  is called a **function type**.

# Def.: Types

Let  $\mathcal{T}$  be the least set s.t:

$$o \in \mathcal{T}$$

$$\iota \in \mathcal{T}$$

$$\forall \alpha, \beta \in \mathcal{T} : (\alpha\beta) \in \mathcal{T}$$

We say that  $\alpha \in \mathcal{T}$  is a **simple type** (or type).  
 $(\alpha\beta)$  is called a **function type**.

- The set  $\mathcal{T}$  is defined inductively.
- The set  $\mathcal{T}$  is "freely generated".



# Ex.: Freely Generated



Consider the set  $\mathbb{N} = \{0, 1, 2, \dots\}$ .

- $0 \in \mathbb{N}$

# Ex.: Freely Generated



Consider the set  $\mathbb{N} = \{0, 1, 2, \dots\}$ .

- $0 \in \mathbb{N}$
- $\forall n \in \mathbb{N} : s(n) \in \mathbb{N}$ .

# Ex.: Freely Generated



Consider the set  $\mathbb{N} = \{0, 1, 2, \dots\}$ .

- $0 \in \mathbb{N}$
- $\forall n \in \mathbb{N} : s(n) \in \mathbb{N}.$
- $\forall n : 0 \neq s(n).$

# Ex.: Freely Generated



Consider the set  $\mathbb{N} = \{0, 1, 2, \dots\}$ .

- $0 \in \mathbb{N}$
- $\forall n \in \mathbb{N} : s(n) \in \mathbb{N}.$
- $\forall n : 0 \neq s(n).$
- $\forall m, n : s(m) = s(n) \Rightarrow m = n.$

# Ex.: Freely Generated



Consider the set  $\mathbb{N} = \{0, 1, 2, \dots\}$ .

- $0 \in \mathbb{N}$
- $\forall n \in \mathbb{N} : s(n) \in \mathbb{N}$ .
- $\forall n : 0 \neq s(n)$ .
- $\forall m, n : s(m) = s(n) \Rightarrow m = n$ .

The set  $\mathbb{N}$  is "freely generated".

# Ex.: Freely Generated

Consider the set  $\mathbb{N} = \{0, 1, 2, \dots\}$ .

- $0 \in \mathbb{N}$
- $\forall n \in \mathbb{N} : s(n) \in \mathbb{N}$ .
- $\forall n : 0 \neq s(n)$ .
- $\forall m, n : s(m) = s(n) \Rightarrow m = n$ .

The set  $\mathbb{N}$  is "freely generated".

Contrast  $\mathbb{N}$  to  $\mathbb{Z} = \{\dots, -1, 0, 1, \dots\}$ .

Note that  $\mathbb{Z}$  contains 0 and is closed under successor, but is not the least such set.

# Ex.: Freely Generated



The set  $\mathcal{T}$  is "freely generated":

■  $o \neq \iota$

# Ex.: Freely Generated



The set  $\mathcal{T}$  is "freely generated":

- $\circ \neq \iota$
- $\circ \neq (\alpha\beta)$



# Ex.: Freely Generated



The set  $\mathcal{T}$  is "freely generated":

- $\circ \neq \iota$
- $\circ \neq (\alpha\beta)$
- $\iota \neq (\alpha\beta)$

# Ex.: Freely Generated



The set  $\mathcal{T}$  is "freely generated":

- $\circ \neq \iota$
- $\circ \neq (\alpha\beta)$
- $\iota \neq (\alpha\beta)$
- $(\alpha\beta) = (\gamma\delta) \Rightarrow \alpha = \gamma \wedge \beta = \delta$

# Ex.: Types

---



- $(o\iota) \in \mathcal{T}$

# Ex.: Types



- $(o\iota) \in \mathcal{T}$
- $(o(o\iota)) \in \mathcal{T}$

# Ex.: Types



- $(o\iota) \in \mathcal{T}$
- $(o(o\iota)) \in \mathcal{T}$
- $(\iota\iota) \in \mathcal{T}$

# Ex.: Types



- $(o\iota) \in \mathcal{T}$
- $(o(o\iota)) \in \mathcal{T}$
- $(\iota\iota) \in \mathcal{T}$
- $((o\iota)\iota) \in \mathcal{T}$

# Ex.: Types



- $(o\iota) \in \mathcal{T}$
- $(o(o\iota)) \in \mathcal{T}$
- $(\iota\iota) \in \mathcal{T}$
- $((o\iota)\iota) \in \mathcal{T}$

Is  $(o\iota\iota)$  also a type?

# Ex.: Types



- $(o\iota) \in \mathcal{T}$
- $(o(o\iota)) \in \mathcal{T}$
- $(\iota\iota) \in \mathcal{T}$
- $((o\iota)\iota) \in \mathcal{T}$

Is  $(o\iota\iota)$  also a type? – no



# Ex.: Types

- $(o\iota) \in \mathcal{T}$
- $(o(o\iota)) \in \mathcal{T}$
- $(\iota\iota) \in \mathcal{T}$
- $((o\iota)\iota) \in \mathcal{T}$

Is  $(o\iota\iota)$  also a type? – no

But we can and will consider it shorthand by replacing missing parenthesis, associating to the left:  $(o\iota\iota) = ((o\iota)\iota) \neq (o(\iota\iota))$ .

# Def.: Functions

---



Let  $A, B$  be sets.

# Def.: Functions

---



Let  $A, B$  be sets.

$f : B \rightarrow A$  : a function from  $B$  to  $A$ .

# Def.: Functions

---



Let  $A, B$  be sets.

$f : B \rightarrow A$  : a function from  $B$  to  $A$ .

$A^B$  : set of functions from  $B$  to  $A$ .

# Def.: Functions

Let  $A, B$  be sets.

$f : B \rightarrow A$  : a function from  $B$  to  $A$ .

$A^B$  : set of functions from  $B$  to  $A$ .

Assume (only for the moment) that  $A, B$  are finite.

# Def.: Functions

Let  $A, B$  be sets.

$f : B \rightarrow A$  : a function from  $B$  to  $A$ .

$A^B$  : set of functions from  $B$  to  $A$ .

Assume (only for the moment) that  $A, B$  are finite.

Let  $|A| = m, |B| = n$ . Then  $|A^B| = m^n = |A|^{|B|}$ .

# Def.: Functions

Let  $A, B$  be sets.

$f : B \rightarrow A$  : a function from  $B$  to  $A$ .

$A^B$  : set of functions from  $B$  to  $A$ .

Assume (only for the moment) that  $A, B$  are finite.

Let  $|A| = m, |B| = n$ . Then  $|A^B| = m^n = |A|^{|B|}$ .

Example:

# Def.: Functions

Let  $A, B$  be sets.

$f : B \rightarrow A$  : a function from  $B$  to  $A$ .

$A^B$  : set of functions from  $B$  to  $A$ .

Assume (only for the moment) that  $A, B$  are finite.

Let  $|A| = m, |B| = n$ . Then  $|A^B| = m^n = |A|^{|B|}$ .

Example:

- $f : \{0, 1, 2\} \rightarrow \{0, 1\}$



# Def.: Functions

Let  $A, B$  be sets.

$f : B \rightarrow A$  : a function from  $B$  to  $A$ .

$A^B$  : set of functions from  $B$  to  $A$ .

Assume (only for the moment) that  $A, B$  are finite.

Let  $|A| = m, |B| = n$ . Then  $|A^B| = m^n = |A|^{|B|}$ .

Example:

- $f : \{0, 1, 2\} \rightarrow \{0, 1\}$
- $f(0), f(1), f(2) \in \{0, 1\}$

# Def.: Functions

Let  $A, B$  be sets.

$f : B \rightarrow A$  : a function from  $B$  to  $A$ .

$A^B$  : set of functions from  $B$  to  $A$ .

Assume (only for the moment) that  $A, B$  are finite.

Let  $|A| = m, |B| = n$ . Then  $|A^B| = m^n = |A|^{|B|}$ .

Example:

- $f : \{0, 1, 2\} \rightarrow \{0, 1\}$
- $f(0), f(1), f(2) \in \{0, 1\}$
- $A = \{0, 1\}, B = \{0, 1, 2\}$

# Def.: Functions

Let  $A, B$  be sets.

$f : B \rightarrow A$  : a function from  $B$  to  $A$ .

$A^B$  : set of functions from  $B$  to  $A$ .

Assume (only for the moment) that  $A, B$  are finite.

Let  $|A| = m, |B| = n$ . Then  $|A^B| = m^n = |A|^{|B|}$ .

Example:

- $f : \{0, 1, 2\} \rightarrow \{0, 1\}$
- $f(0), f(1), f(2) \in \{0, 1\}$
- $A = \{0, 1\}, B = \{0, 1, 2\}$
- $|A^B| = 2 \cdot 2 \cdot 2 = 2^3 = 8$

# Ex.: Sets of Functions



Let  $F = \{f : B \rightarrow A \mid \forall x, y \in B : x \leq y \Rightarrow f(x) \leq f(y)\} \subseteq A^B$ .

# Ex.: Sets of Functions



Let  $F = \{f : B \rightarrow A \mid \forall x, y \in B : x \leq y \Rightarrow f(x) \leq f(y)\} \subseteq A^B$ .

$|F| = ?$

# Ex.: Sets of Functions

Let  $F = \{f : B \rightarrow A \mid \forall x, y \in B : x \leq y \Rightarrow f(x) \leq f(y)\} \subseteq A^B$ .

$|F| = ?$

$A^B$	$f(0)$	$f(1)$	$f(2)$
$K_0 \in F$	0	0	0
$\in F$	0	0	1
$\notin F$	0	1	0
$\in F$	0	1	1
$g \notin F$	1	0	0
$\notin F$	1	0	1
$\notin F$	1	1	0
$K_1 \in F$	1	1	1

Consider:

$g : x = 0, y = 1, x \leq y$ , but  
 $f(x) \geq f(y) \Rightarrow g \notin F$ .

# Ex.: Sets of Functions

Let  $F = \{f : B \rightarrow A \mid \forall x, y \in B : x \leq y \Rightarrow f(x) \leq f(y)\} \subseteq A^B$ .

$|F| = ?$

$A^B$	$f(0)$	$f(1)$	$f(2)$
$K_0 \in F$	0	0	0
$\in F$	0	0	1
$\notin F$	0	1	0
$\in F$	0	1	1
$g \notin F$	1	0	0
$\notin F$	1	0	1
$\notin F$	1	1	0
$K_1 \in F$	1	1	1

Consider:

$g : x = 0, y = 1, x \leq y$ , but  
 $f(x) \geq f(y) \Rightarrow g \notin F$ .

$|F| = 4$

# Ex.: Sets of Labelled Functions



$$C = \{\text{red, blue, green}\}$$



# Ex.: Sets of Labelled Functions



$$C = \{\text{red, blue, green}\}$$

$$F_C = \{\langle c, f \rangle \mid c \in C, f \in F\}$$

# Ex.: Sets of Labelled Functions



$$C = \{\text{red, blue, green}\}$$

$$F_C = \{\langle c, f \rangle \mid c \in C, f \in F\}$$

$$|F_C| = 3 \cdot 4 = 12$$

# Def.: Frames

---



A **frame** is a family  $(D_\alpha)_{\alpha \in \mathcal{I}}$  of nonempty sets s.t:

# Def.: Frames

---



A **frame** is a family  $(D_\alpha)_{\alpha \in \mathcal{T}}$  of nonempty sets s.t:

$$\forall \alpha, \beta \in \mathcal{T} : D_{\alpha\beta} \subseteq D_\alpha^{D_\beta}$$

# Def.: Frames



A **frame** is a family  $(D_\alpha)_{\alpha \in \mathcal{T}}$  of nonempty sets s.t:

$$\forall \alpha, \beta \in \mathcal{T} : D_{\alpha\beta} \subseteq D_\alpha^{D_\beta}$$

A Frame is called **standard** if

# Def.: Frames

A **frame** is a family  $(D_\alpha)_{\alpha \in \mathcal{T}}$  of nonempty sets s.t:

$$\forall \alpha, \beta \in \mathcal{T} : D_{\alpha\beta} \subseteq D_\alpha^{D_\beta}$$

A Frame is called **standard** if

$$D_{\alpha\beta} = D_\alpha^{D_\beta} \quad \forall \alpha, \beta \in \mathcal{T}$$

# Ex.: Frames

---



$$D_o = \{\perp, \top\}$$

# Ex.: Frames

---



$$D_o = \{\perp, \top\}$$

$$D_\iota = \{1\}$$



# Ex.: Frames



$$D_o = \{\perp, \top\}$$

$$D_\iota = \{1\}$$

$$D_{\alpha\beta} = D_\alpha^{D_\beta}$$

# Ex.: Frames

$$D_o = \{\perp, \top\}$$

$$D_i = \{1\}$$

$$D_{\alpha\beta} = D_{\alpha}^{D_{\beta}}$$

$D$ : the standard frame with  $D_o = \{\perp, \top\}$ ,  $D_i = \{1\}$

# Ex.: Frames (Contd.)



Consider the set  $D_{o(\iota)((o(\iota o)))}$ . Is the set empty?

# Ex.: Frames (Contd.)



Consider the set  $D_{o(\iota)((o(\iota o)))}$ . Is the set empty? — no!

# Ex.: Frames (Contd.)

Consider the set  $D_{o(u)((o(uo)))}$ . Is the set empty? — no!

**Claim:**  $\forall \alpha \in \mathcal{T} : D_\alpha \neq \emptyset$ .

# Ex.: Frames (Contd.)

Consider the set  $D_{o(\iota)((o(\iota o)))}$ . Is the set empty? — no!

**Claim:**  $\forall \alpha \in \mathcal{T} : D_\alpha \neq \emptyset$ .

**Proof:** induction on type.

# Ex.: Frames (Contd.)

Consider the set  $D_{o(\iota)((o(\iota o)))}$ . Is the set empty? — no!

**Claim:**  $\forall \alpha \in \mathcal{T} : D_\alpha \neq \emptyset$ .

**Proof:** induction on type.

- Base:  $D_o = \{\perp, \top\} \neq \emptyset, D_i = \{1\} \neq \emptyset$ .

# Ex.: Frames (Contd.)

Consider the set  $D_{o(\iota)((o(\iota o)))}$ . Is the set empty? — no!

**Claim:**  $\forall \alpha \in \mathcal{T} : D_\alpha \neq \emptyset$ .

**Proof:** induction on type.

- Base:  $D_o = \{\perp, \top\} \neq \emptyset, D_i = \{1\} \neq \emptyset$ .
- Step: Assume  $D_\alpha \neq \emptyset \wedge D_\beta \neq \emptyset$ . Want to show:  $D_{\alpha\beta} \neq \emptyset$ .



# Ex.: Frames (Contd.)

Consider the set  $D_{o(\iota)((o(\iota o)))}$ . Is the set empty? — no!

**Claim:**  $\forall \alpha \in \mathcal{T} : D_\alpha \neq \emptyset$ .

**Proof:** induction on type.

- Base:  $D_o = \{\perp, \top\} \neq \emptyset, D_i = \{1\} \neq \emptyset$ .
- Step: Assume  $D_\alpha \neq \emptyset \wedge D_\beta \neq \emptyset$ . Want to show:  $D_{\alpha\beta} \neq \emptyset$ .  
Since  $D_\alpha \neq \emptyset \Rightarrow \exists a \in D_\alpha$ ,

# Ex.: Frames (Contd.)

Consider the set  $D_{o(\iota)((o(\iota o)))}$ . Is the set empty? — no!

**Claim:**  $\forall \alpha \in \mathcal{T} : D_\alpha \neq \emptyset$ .

**Proof:** induction on type.

- Base:  $D_o = \{\perp, \top\} \neq \emptyset, D_i = \{1\} \neq \emptyset$ .
- Step: Assume  $D_\alpha \neq \emptyset \wedge D_\beta \neq \emptyset$ . Want to show:  $D_{\alpha\beta} \neq \emptyset$ .  
Since  $D_\alpha \neq \emptyset \Rightarrow \exists a \in D_\alpha$ , hence  $K_a \in D_{\alpha\beta}$ .

# Ex.: Frames (Contd.)

Consider the set  $D_{o(\iota)((o(\iota o)))}$ . Is the set empty? — no!

**Claim:**  $\forall \alpha \in \mathcal{T} : D_\alpha \neq \emptyset$ .

**Proof:** induction on type.

- Base:  $D_o = \{\perp, \top\} \neq \emptyset, D_i = \{1\} \neq \emptyset$ .
- Step: Assume  $D_\alpha \neq \emptyset \wedge D_\beta \neq \emptyset$ . Want to show:  $D_{\alpha\beta} \neq \emptyset$ .  
Since  $D_\alpha \neq \emptyset \Rightarrow \exists a \in D_\alpha$ , hence  $K_a \in D_{\alpha\beta}$ .

(Here  $K_a$  is the constant function which always returns  $a$ . We will often use this notation for constant functions.)

# Def.: Typed Applicative Structure



A (typed) applicative structure is a tuple

# Def.: Typed Applicative Structure



A (typed) applicative structure is a tuple

$$\langle D, @ \rangle$$

where

# Def.: Typed Applicative Structure



A (typed) applicative structure is a tuple

$$\langle D, @ \rangle$$

where

- $D := (D_\alpha)_{\alpha \in \mathcal{T}}$  is a family of nonempty sets

# Def.: Typed Applicative Structure

A (typed) applicative structure is a tuple

$$\langle D, @ \rangle$$

where

- $D := (D_\alpha)_{\alpha \in \mathcal{T}}$  is a family of nonempty sets
- $@ := (@^{\alpha\beta} : D_{\alpha\beta} \times D_\beta \rightarrow D_\alpha)_{\alpha, \beta \in \mathcal{T}}$

# Def.: Typed Applicative Structure

A (typed) applicative structure is a tuple

$$\langle D, @ \rangle$$

where

- $D := (D_\alpha)_{\alpha \in \mathcal{T}}$  is a family of nonempty sets
- $@ := (@^{\alpha\beta} : D_{\alpha\beta} \times D_\beta \rightarrow D_\alpha)_{\alpha, \beta \in \mathcal{T}}$

Usually we write  $f@b$  for  $@^{\alpha\beta}(f, b)$  when  $f \in D_{\alpha\beta} \wedge b \in D_\beta$



# Rem.: Curryng

---



The application operator  $@$  in an applicative structure is an abstract version of function application.

The application operator  $@$  in an applicative structure is an abstract version of function application. It is no restriction to exclusively use a binary application operator, which corresponds to unary function application,

The application operator  $@$  in an applicative structure is an abstract version of function application. It is no restriction to exclusively use a binary application operator, which corresponds to unary function application, since we can define higher-arity application operators from the binary one by setting  $f@(a^1, \dots, a^n) := (\dots (f@a^1) \dots @a^n)$  (“Currying”).

# Interesting Properties

---



Let  $\mathcal{D}$  be a frame.

# Interesting Properties



Let  $D$  be a frame.

$$\forall f, g \in D_{\alpha\beta} \quad (\forall b \in D_{\beta} : f(b) = g(b)) \Rightarrow f = g.$$

# Interesting Properties



Let  $D$  be a frame.

$$\forall f, g \in D_{\alpha\beta} \quad (\forall b \in D_{\beta} : f(b) = g(b)) \Rightarrow f = g.$$

Let  $\langle D, @ \rangle$  be an applicative structure. Consider the property:

# Interesting Properties



Let  $D$  be a frame.

$$\forall f, g \in D_{\alpha\beta} \quad (\forall b \in D_{\beta} : f(b) = g(b)) \Rightarrow f = g.$$

Let  $\langle D, @ \rangle$  be an applicative structure. Consider the property:

$$\forall f, g \in D_{\alpha\beta} \quad (\forall b \in D_{\beta} : f@b = g@b) \Rightarrow f = g.$$

# Def.: Functional Applicative Structures



Given an applicative structure  $\langle D, @ \rangle$ .



# Def.: Functional Applicative Structures



Given an applicative structure  $\langle D, @ \rangle$ . We say that  $\langle D, @ \rangle$  is **functional** if

# Def.: Functional Applicative Structures



Given an applicative structure  $\langle D, @ \rangle$ . We say that  $\langle D, @ \rangle$  is **functional** if

$$\forall \alpha, \beta \in \mathcal{T} : \forall f, g \in D_{\alpha\beta} (\forall b \in D_{\beta} : f@b = g@b) \Rightarrow f = g$$

# Def.: Full Applicative Structures



Given an applicative structure  $\langle D, @ \rangle$ .

# Def.: Full Applicative Structures



Given an applicative structure  $\langle D, @ \rangle$ . We say that  $\langle D, @ \rangle$  is **full** if

# Def.: Full Applicative Structures



Given an applicative structure  $\langle D, @ \rangle$ . We say that  $\langle D, @ \rangle$  is **full** if

$$\forall \alpha, \beta \quad \forall h : D_\beta \rightarrow D_\alpha \quad \exists f \in D_{\alpha\beta} \forall b \in D_\beta : f@b = h(b)$$

# Def.: Standard Applicative Structures



An applicative structure  $\mathcal{A} := \langle D, @ \rangle$  is called **standard** if

# Def.: Standard Applicative Structures



An applicative structure  $\mathcal{A} := \langle D, @ \rangle$  is called **standard** if it is a frame structure (i.e.  $@$  is function application) where  $\mathcal{D}$  is standard.

# Def.: Standard Applicative Structures



An applicative structure  $\mathcal{A} := \langle D, @ \rangle$  is called **standard** if it is a frame structure (i.e.  $@$  is function application) where  $\mathcal{D}$  is standard.

Note that the definitions of functional, full, and standard impose restrictions on the domains for function types only.



# Rem.: Frames and Applicative Structures



It is easy to show that every frame is functional.

# Rem.: Frames and Applicative Structures



It is easy to show that every frame is functional.

Furthermore, an applicative structure is standard iff it is a full frame.

# Example: Full Functional Appl. Structure



Let  $D_\alpha = \{1\} \quad \forall \alpha$

# Example: Full Functional Appl. Structure



Let  $D_\alpha = \{1\} \quad \forall \alpha$

Let  $f@b = 1 \quad \forall f \in D_{\alpha\beta} \quad \forall b \in D_\beta$

# Example: Full Functional Appl. Structure



Let  $D_\alpha = \{1\} \quad \forall \alpha$

Let  $f@b = 1 \quad \forall f \in D_{\alpha\beta} \quad \forall b \in D_\beta$

$\langle D, @ \rangle$  is a full functional applicative structure, but it is not a frame.

# Example: Full Functional Appl. Structure



Let  $D_\alpha = \{1\} \quad \forall \alpha$

Let  $f@b = 1 \quad \forall f \in D_{\alpha\beta} \quad \forall b \in D_\beta$

$\langle D, @ \rangle$  is a full functional applicative structure, but it is not a frame.

$1 \in D_{\circ\circ}$  but  $1 \notin D_{\circ}^{D_{\circ}} \Rightarrow D_{\circ\circ} \not\subseteq D_{\circ}^{D_{\circ}}$

# Def.: Homomorphic Appl. Structures



Let  $\langle D^1, @^1 \rangle$  and  $\langle D^2, @^2 \rangle$  are applicative structures.

# Def.: Homomorphic Appl. Structures



Let  $\langle D^1, @^1 \rangle$  and  $\langle D^2, @^2 \rangle$  are applicative structures. We say that  $\kappa$  is a **homomorphism** from  $\langle D^1, @^1 \rangle$  to  $\langle D^2, @^2 \rangle$  if



# Def.: Homomorphic Appl. Structures



Let  $\langle D^1, @^1 \rangle$  and  $\langle D^2, @^2 \rangle$  are applicative structures. We say that  $\kappa$  is a **homomorphism** from  $\langle D^1, @^1 \rangle$  to  $\langle D^2, @^2 \rangle$  if

- $\kappa_\alpha : D_\alpha^1 \rightarrow D_\alpha^2 \quad \forall \alpha \in \mathcal{T}$

# Def.: Homomorphic Appl. Structures



Let  $\langle D^1, @^1 \rangle$  and  $\langle D^2, @^2 \rangle$  are applicative structures. We say that  $\kappa$  is a **homomorphism** from  $\langle D^1, @^1 \rangle$  to  $\langle D^2, @^2 \rangle$  if

- $\kappa_\alpha : D_\alpha^1 \rightarrow D_\alpha^2 \quad \forall \alpha \in \mathcal{T}$
- $\forall \alpha, \beta \in \mathcal{T}, \quad \forall f \in D_{\alpha\beta}^1, \quad \forall b \in D_\beta^1:$

$$\kappa(f)@^2\kappa(b) = \kappa(f@^1b)$$

# Def.: Isomorphic Appl. Structures



We say that  $\langle D^1, @^1 \rangle$  and  $\langle D^2, @^2 \rangle$  are **isomorphic** if  $\exists i, j$  s.t:

# Def.: Isomorphic Appl. Structures

We say that  $\langle D^1, @^1 \rangle$  and  $\langle D^2, @^2 \rangle$  are **isomorphic** if  $\exists i, j$  s.t:

- $i$  is a homomorphism from  $\langle D^1, @^1 \rangle$  to  $\langle D^2, @^2 \rangle$

# Def.: Isomorphic Appl. Structures

We say that  $\langle D^1, @^1 \rangle$  and  $\langle D^2, @^2 \rangle$  are **isomorphic** if  $\exists i, j$  s.t:

- $i$  is a homomorphism from  $\langle D^1, @^1 \rangle$  to  $\langle D^2, @^2 \rangle$
- $j$  is a homomorphism from  $\langle D^2, @^2 \rangle$  to  $\langle D^1, @^1 \rangle$

# Def.: Isomorphic Appl. Structures

We say that  $\langle D^1, @^1 \rangle$  and  $\langle D^2, @^2 \rangle$  are **isomorphic** if  $\exists i, j$  s.t:

- $i$  is a homomorphism from  $\langle D^1, @^1 \rangle$  to  $\langle D^2, @^2 \rangle$
- $j$  is a homomorphism from  $\langle D^2, @^2 \rangle$  to  $\langle D^1, @^1 \rangle$
- $i$  and  $j$  are inverses (i.e  $i(j(a^2)) = a^2$  and  $j(i(a^1)) = a^1$ ).



# Simply Typed $\lambda$ -Calculus

# Def.: Untyped $\lambda$ -Calculus



Let  $\Sigma = (\mathcal{V}, \mathcal{C})$  be a signature where



# Def.: Untyped $\lambda$ -Calculus



Let  $\Sigma = (\mathcal{V}, \mathcal{C})$  be a signature where

- $\mathcal{V}$  — countably infinite set of variables

# Def.: Untyped $\lambda$ -Calculus



Let  $\Sigma = (\mathcal{V}, \mathcal{C})$  be a signature where

- $\mathcal{V}$  — countably infinite set of variables
- $\mathcal{C}$  — possibly empty set of constants

# Def.: Untyped $\lambda$ -Calculus

Let  $\Sigma = (\mathcal{V}, \mathcal{C})$  be a signature where

- $\mathcal{V}$  — countably infinite set of variables
- $\mathcal{C}$  — possibly empty set of constants

We define the set  $\Lambda = wff_{\Sigma}(\Sigma)$  to be the smallest set s.t:

# Def.: Untyped $\lambda$ -Calculus

Let  $\Sigma = (\mathcal{V}, \mathcal{C})$  be a signature where

- $\mathcal{V}$  — countably infinite set of variables
- $\mathcal{C}$  — possibly empty set of constants

We define the set  $\Lambda = wff_{\Sigma}(\Sigma)$  to be the smallest set s.t:

- $x \in \mathcal{V}$  then  $x \in \Lambda$

# Def.: Untyped $\lambda$ -Calculus

Let  $\Sigma = (\mathcal{V}, \mathcal{C})$  be a signature where

- $\mathcal{V}$  — countably infinite set of variables
- $\mathcal{C}$  — possibly empty set of constants

We define the set  $\Lambda = wff_{\Sigma}(\Sigma)$  to be the smallest set s.t:

- $x \in \mathcal{V}$  then  $x \in \Lambda$
- $c \in \mathcal{C}$  then  $c \in \Lambda$

# Def.: Untyped $\lambda$ -Calculus

Let  $\Sigma = (\mathcal{V}, \mathcal{C})$  be a signature where

- $\mathcal{V}$  — countably infinite set of variables
- $\mathcal{C}$  — possibly empty set of constants

We define the set  $\Lambda = wff_{\Sigma}(\Sigma)$  to be the smallest set s.t:

- $x \in \mathcal{V}$  then  $x \in \Lambda$
- $c \in \mathcal{C}$  then  $c \in \Lambda$
- $A \in \Lambda, B \in \Lambda$  then  $(A B) \in \Lambda$

# Def.: Untyped $\lambda$ -Calculus

Let  $\Sigma = (\mathcal{V}, \mathcal{C})$  be a signature where

- $\mathcal{V}$  — countably infinite set of variables
- $\mathcal{C}$  — possibly empty set of constants

We define the set  $\Lambda = wff_{\Sigma}(\Sigma)$  to be the smallest set s.t:

- $x \in \mathcal{V}$  then  $x \in \Lambda$
- $c \in \mathcal{C}$  then  $c \in \Lambda$
- $A \in \Lambda, B \in \Lambda$  then  $(A B) \in \Lambda$
- $x \in \mathcal{V}, A \in \Lambda$  then  $(\lambda x.A) \in \Lambda$

# Simply Typed $\lambda$ -Calculus



Let  $\Sigma = (\mathcal{V}^\alpha, \mathcal{C}^\alpha)$  be a signature where



# Simply Typed $\lambda$ -Calculus



Let  $\Sigma = (\mathcal{V}^\alpha, \mathcal{C}^\alpha)$  be a signature where

- $\mathcal{V}^\alpha = \bigcup_{\alpha \in \mathcal{I}} \mathcal{V}_\alpha$  — countably infinite sets of variables

# Simply Typed $\lambda$ -Calculus



Let  $\Sigma = (\mathcal{V}^\alpha, \mathcal{C}^\alpha)$  be a signature where

- $\mathcal{V}^\alpha = \bigcup_{\alpha \in \mathcal{T}} \mathcal{V}_\alpha$  — countably infinite sets of variables
- $\mathcal{C}^\alpha = \bigcup_{\alpha \in \mathcal{T}} \mathcal{C}_\alpha$  — possibly empty sets of constants

# Simply Typed $\lambda$ -Calculus

Let  $\Sigma = (\mathcal{V}^\alpha, \mathcal{C}^\alpha)$  be a signature where

- $\mathcal{V}^\alpha = \bigcup_{\alpha \in \mathcal{T}} \mathcal{V}_\alpha$  — countably infinite sets of variables
- $\mathcal{C}^\alpha = \bigcup_{\alpha \in \mathcal{T}} \mathcal{C}_\alpha$  — possibly empty sets of constants

We define the set  $\Lambda^\alpha = wff_\Sigma(\Sigma)_\alpha = \bigcup_{\alpha \in \mathcal{T}} \Lambda_\alpha$  to be the smallest set s.t:

# Simply Typed $\lambda$ -Calculus

Let  $\Sigma = (\mathcal{V}^\alpha, \mathcal{C}^\alpha)$  be a signature where

- $\mathcal{V}^\alpha = \bigcup_{\alpha \in \mathcal{T}} \mathcal{V}_\alpha$  — countably infinite sets of variables
- $\mathcal{C}^\alpha = \bigcup_{\alpha \in \mathcal{T}} \mathcal{C}_\alpha$  — possibly empty sets of constants

We define the set  $\Lambda^\alpha = wff_\Sigma(\Sigma)_\alpha = \bigcup_{\alpha \in \mathcal{T}} \Lambda_\alpha$  to be the smallest set s.t:

- $x_\alpha \in \mathcal{V}_\alpha$  then  $x_\alpha \in \Lambda_\alpha$

# Simply Typed $\lambda$ -Calculus

Let  $\Sigma = (\mathcal{V}^\alpha, \mathcal{C}^\alpha)$  be a signature where

- $\mathcal{V}^\alpha = \bigcup_{\alpha \in \mathcal{T}} \mathcal{V}_\alpha$  — countably infinite sets of variables
- $\mathcal{C}^\alpha = \bigcup_{\alpha \in \mathcal{T}} \mathcal{C}_\alpha$  — possibly empty sets of constants

We define the set  $\Lambda^\alpha = wff_\Sigma(\Sigma)_\alpha = \bigcup_{\alpha \in \mathcal{T}} \Lambda_\alpha$  to be the smallest set s.t:

- $x_\alpha \in \mathcal{V}_\alpha$  then  $x_\alpha \in \Lambda_\alpha$
- $c_\alpha \in \mathcal{C}_\alpha$  then  $c_\alpha \in \Lambda_\alpha$

# Simply Typed $\lambda$ -Calculus

Let  $\Sigma = (\mathcal{V}^\alpha, \mathcal{C}^\alpha)$  be a signature where

- $\mathcal{V}^\alpha = \bigcup_{\alpha \in \mathcal{T}} \mathcal{V}_\alpha$  — countably infinite sets of variables
- $\mathcal{C}^\alpha = \bigcup_{\alpha \in \mathcal{T}} \mathcal{C}_\alpha$  — possibly empty sets of constants

We define the set  $\Lambda^\alpha = wff_\Sigma(\Sigma)_\alpha = \bigcup_{\alpha \in \mathcal{T}} \Lambda_\alpha$  to be the smallest set s.t:

- $x_\alpha \in \mathcal{V}_\alpha$  then  $x_\alpha \in \Lambda_\alpha$
- $c_\alpha \in \mathcal{C}_\alpha$  then  $c_\alpha \in \Lambda_\alpha$
- $A_{\alpha\beta} \in \Lambda_{\alpha\beta}, B_\beta \in \Lambda_\beta$  then  $(A B) \in \Lambda_\alpha$

# Simply Typed $\lambda$ -Calculus

Let  $\Sigma = (\mathcal{V}^\alpha, \mathcal{C}^\alpha)$  be a signature where

- $\mathcal{V}^\alpha = \bigcup_{\alpha \in \mathcal{T}} \mathcal{V}_\alpha$  — countably infinite sets of variables
- $\mathcal{C}^\alpha = \bigcup_{\alpha \in \mathcal{T}} \mathcal{C}_\alpha$  — possibly empty sets of constants

We define the set  $\Lambda^\alpha = wff_\Sigma(\Sigma)_\alpha = \bigcup_{\alpha \in \mathcal{T}} \Lambda_\alpha$  to be the smallest set s.t:

- $x_\alpha \in \mathcal{V}_\alpha$  then  $x_\alpha \in \Lambda_\alpha$
- $c_\alpha \in \mathcal{C}_\alpha$  then  $c_\alpha \in \Lambda_\alpha$
- $A_{\alpha\beta} \in \Lambda_{\alpha\beta}, B_\beta \in \Lambda_\beta$  then  $(A B) \in \Lambda_\alpha$
- $x_\alpha \in \mathcal{V}_\alpha, A_\beta \in \Lambda_\beta$  then  $(\lambda x_\alpha. A_\beta)_{\beta\alpha} \in \Lambda_{\beta\alpha}$

# Notational Conventions



- brackets may be avoided:  $A B C \rightsquigarrow ((A B) C)$



# Notational Conventions

- brackets may be avoided:  $A B C \rightsquigarrow ((A B) C)$
- $\lambda x_\iota . A_{o\iota} B_\iota C_\iota$  — dots as far to the right as is consistent:  
 $((\lambda x_\iota . A_{o\iota} B_\iota) C_\iota)$

# Notational Conventions

- brackets may be avoided:  $A B C \rightsquigarrow ((A B) C)$
- $\lambda x_\iota . A_{o_\iota} B_\iota C_\iota$  — dots as far to the right as is consistent:  
 $((\lambda x_\iota . A_{o_\iota} B_\iota) C_\iota)$
- $\lambda x, y . A \rightsquigarrow (\lambda x . (\lambda y . A))$

# Notational Conventions

- brackets may be avoided:  $A B C \rightsquigarrow ((A B) C)$
- $\lambda x_\iota . A_{o_\iota} B_\iota C_\iota$  — dots as far to the right as is consistent:  
 $((\lambda x_\iota . A_{o_\iota} B_\iota) C_\iota)$
- $\lambda x, y . A \rightsquigarrow (\lambda x . (\lambda y . A))$
- $\lambda \bar{x}^n . A \rightsquigarrow (\lambda x_1 . (\dots (\lambda x_n . A) \dots))$

# Notational Conventions

- brackets may be avoided:  $A B C \rightsquigarrow ((A B) C)$
- $\lambda x_\iota . A_{o_\iota} B_\iota C_\iota$  — dots as far to the right as is consistent:  
 $((\lambda x_\iota . A_{o_\iota} B_\iota) C_\iota)$
- $\lambda x, y . A \rightsquigarrow (\lambda x . (\lambda y . A))$
- $\lambda \bar{x}^n . A \rightsquigarrow (\lambda x_1 . (\dots (\lambda x_n . A) \dots))$
- $\lambda \bar{x} . A$  —  $n$  is not important

# Notational Conventions

- brackets may be avoided:  $A B C \rightsquigarrow ((A B) C)$
- $\lambda x_\iota . A_{o_\iota} B_\iota C_\iota$  — dots as far to the right as is consistent:  
 $((\lambda x_\iota . A_{o_\iota} B_\iota) C_\iota)$
- $\lambda x, y . A \rightsquigarrow (\lambda x . (\lambda y . A))$
- $\lambda \bar{x}^n . A \rightsquigarrow (\lambda x_1 . (\dots (\lambda x_n . A) \dots))$
- $\lambda \bar{x} . A$  —  $n$  is not important
- $(f \bar{A}^n) \rightsquigarrow (\dots ((f A^1) A^2) \dots A^n)$

# Def.: Positions in $\lambda$ -Terms



Consider the following term:

$$((\lambda x.x)((\lambda y.y)(\lambda z.z)))$$

# Def.: Positions in $\lambda$ -Terms

Consider the following term:

$$((\lambda x.x)((\lambda y.y)(\lambda z.z)))$$

The **position** [212] points to the red **y** in

$$((\lambda x.x)((\lambda y.y)(\lambda z.z)))$$

# Def.: Positions in $\lambda$ -Terms

Consider the following term:

$$((\lambda x.x)((\lambda y.y)(\lambda z.z)))$$

The **position** [212] points to the red **y** in

$$((\lambda x.x)((\lambda y.\textcolor{red}{y})(\lambda z.z)))$$

... Graphics on Blackboard ...



# Def.: Position (Contd.)



The expression

$A_p$

refers to the **subterm of A at position p**.

# Def.: Position (Contd.)



The expression

$$A_p$$

refers to the **subterm of A at position p**.

Example: Consider  $T := ((\lambda x.x)((\lambda y.y)(\lambda z.z)))$

# Def.: Position (Contd.)



The expression

$$A_p$$

refers to the **subterm of A at position p**.

Example: Consider  $T := ((\lambda x.x)((\lambda y.y)(\lambda z.z)))$

$$T_{[212]} = y$$

# Def.: Replacement at Position



Replacement of  $A_p$  in  $A$  by a term  $B$  is denoted as

$$A[B]_p$$

# Def.: Replacement at Position

Replacement of  $A_p$  in  $A$  by a term  $B$  is denoted as

$$A[B]_p$$

Example:

$$T[(f\ x)]_{[212]} = ((\lambda x.x)((\lambda y.(f\ x))(\lambda z.z)))$$

# Def.: Scope of $\lambda$ -Term



$(\lambda x.A)$  : We say that  $A$  is in the **scope** of  $\lambda$ -binder that binds  $x$ .

# Def.: Free and Bound Variables



An occurrence of a variable  $x$  in a term  $A$  is called **bound** if it is in the scope of a  $\lambda$ -binder that binds  $x$ .

# Def.: Free and Bound Variables

An occurrence of a variable  $x$  in a term  $A$  is called **bound** if it is in the scope of a  $\lambda$ -binder that binds  $x$ .

Otherwise it is called **free**.



# Def.: Free and Bound Variables

An occurrence of a variable  $x$  in a term  $A$  is called **bound** if it is in the scope of a  $\lambda$ -binder that binds  $x$ .

Otherwise it is called **free**.

We denote the **set of all free variables** in a  $\lambda$ -term as  $FV(A)$ .



# Syntax: Simply Typed $\lambda$ -Calculus (Contd.)

# Def.: Substitution

---



Substitution is a map

# Def.: Substitution



Substitution is a map

$$[A/x] : \Lambda \rightarrow \Lambda \quad (\text{untyped})$$

# Def.: Substitution



Substitution is a map

$$\begin{aligned} [A/x] &: \Lambda \rightarrow \Lambda \quad (\text{untyped}) \\ [A_\alpha/x_\alpha] &: \Lambda_\alpha \rightarrow \Lambda_\alpha \quad (\text{typed}) \end{aligned}$$

# Def.: Substitution

Substitution is a map

$$\begin{aligned} [A/x] &: \Lambda \rightarrow \Lambda \quad (\text{untyped}) \\ [A_\alpha/x_\alpha] &: \Lambda_\alpha \rightarrow \Lambda_\alpha \quad (\text{typed}) \end{aligned}$$

and is defined as follows:

# Def.: Substitution

Substitution is a map

$$[A/x] : \Lambda \rightarrow \Lambda \quad (\text{untyped})$$

$$[A_\alpha/x_\alpha] : \Lambda_\alpha \rightarrow \Lambda_\alpha \quad (\text{typed})$$

and is defined as follows:

$$1. [N_\alpha/x_\alpha]x_\alpha = N_\alpha$$

# Def.: Substitution

Substitution is a map

$$[A/x] : \Lambda \rightarrow \Lambda \quad (\text{untyped})$$

$$[A_\alpha/x_\alpha] : \Lambda_\alpha \rightarrow \Lambda_\alpha \quad (\text{typed})$$

and is defined as follows:

1.  $[N_\alpha/x_\alpha]x_\alpha = N_\alpha$
2.  $[N_\alpha/x_\alpha]a_\beta = a_\beta$  if  $a_\beta \neq x_\alpha \wedge a_\beta \in \mathcal{V}_\beta \cup \mathcal{C}_\beta$



# Def.: Substitution

Substitution is a map

$$[A/x] : \Lambda \rightarrow \Lambda \quad (\text{untyped})$$

$$[A_\alpha/x_\alpha] : \Lambda_\alpha \rightarrow \Lambda_\alpha \quad (\text{typed})$$

and is defined as follows:

1.  $[N_\alpha/x_\alpha]x_\alpha = N_\alpha$
2.  $[N_\alpha/x_\alpha]a_\beta = a_\beta$  if  $a_\beta \neq x_\alpha \wedge a_\beta \in \mathcal{V}_\beta \cup \mathcal{C}_\beta$
3.  $[N_\alpha/x_\alpha](A_{\alpha\alpha} B_\beta) = ([N_\alpha/x_\alpha]A)([N_\alpha/x_\alpha]B)$

# Def.: Substitution

Substitution is a map

$$[A/x] : \Lambda \rightarrow \Lambda \quad (\text{untyped})$$

$$[A_\alpha/x_\alpha] : \Lambda_\alpha \rightarrow \Lambda_\alpha \quad (\text{typed})$$

and is defined as follows:

1.  $[N_\alpha/x_\alpha]x_\alpha = N_\alpha$
2.  $[N_\alpha/x_\alpha]a_\beta = a_\beta$  if  $a_\beta \neq x_\alpha \wedge a_\beta \in \mathcal{V}_\beta \cup \mathcal{C}_\beta$
3.  $[N_\alpha/x_\alpha](A_{\alpha\alpha}B_\beta) = ([N_\alpha/x_\alpha]A)([N_\alpha/x_\alpha]B)$
4.  $[N_\alpha/x_\alpha](\lambda x_\alpha.A_\gamma) = (\lambda x_\alpha.A_\gamma)$

# Def.: Substitution

Substitution is a map

$$[A/x] : \Lambda \rightarrow \Lambda \quad (\text{untyped})$$

$$[A_\alpha/x_\alpha] : \Lambda_\alpha \rightarrow \Lambda_\alpha \quad (\text{typed})$$

and is defined as follows:

1.  $[N_\alpha/x_\alpha]x_\alpha = N_\alpha$
2.  $[N_\alpha/x_\alpha]a_\beta = a_\beta$  if  $a_\beta \neq x_\alpha \wedge a_\beta \in \mathcal{V}_\beta \cup \mathcal{C}_\beta$
3.  $[N_\alpha/x_\alpha](A_\alpha B_\beta) = ([N_\alpha/x_\alpha]A)([N_\alpha/x_\alpha]B)$
4.  $[N_\alpha/x_\alpha](\lambda x_\alpha. A_\gamma) = (\lambda x_\alpha. A_\gamma)$
5.  $[N_\alpha/x_\alpha](\lambda y_\beta. A_\gamma) = (\lambda y_\beta. [N_\alpha/x_\alpha]A_\gamma)$  if  
 $x_\alpha \neq y_\beta \wedge (y_\beta \notin \text{FV}(N_\alpha) \vee x_\alpha \notin \text{FV}(A_\gamma))$

# Def.: Substitution

Substitution is a map

$$[A/x] : \Lambda \rightarrow \Lambda \quad (\text{untyped})$$

$$[A_\alpha/x_\alpha] : \Lambda_\alpha \rightarrow \Lambda_\alpha \quad (\text{typed})$$

and is defined as follows:

1.  $[N_\alpha/x_\alpha]x_\alpha = N_\alpha$
2.  $[N_\alpha/x_\alpha]a_\beta = a_\beta$  if  $a_\beta \neq x_\alpha \wedge a_\beta \in \mathcal{V}_\beta \cup \mathcal{C}_\beta$
3.  $[N_\alpha/x_\alpha](A_\alpha B_\beta) = ([N_\alpha/x_\alpha]A)([N_\alpha/x_\alpha]B)$
4.  $[N_\alpha/x_\alpha](\lambda x_\alpha. A_\gamma) = (\lambda x_\alpha. A_\gamma)$
5.  $[N_\alpha/x_\alpha](\lambda y_\beta. A_\gamma) = (\lambda y_\beta. [N_\alpha/x_\alpha]A_\gamma)$  if  
 $x_\alpha \neq y_\beta \wedge (y_\beta \notin \text{FV}(N_\alpha) \vee x_\alpha \notin \text{FV}(A_\gamma))$
6.  $[N_\alpha/x_\alpha](\lambda y_\alpha. A_\gamma) = (\lambda z_\beta. [N_\alpha/x_\alpha][z_\beta/y_\beta]A_\gamma)$  if  $x_\alpha \neq y_\beta \wedge$   
 $(y_\beta \in \text{FV}(N_\alpha) \wedge x_\alpha \in \text{FV}(A_\gamma))$  and  $z$  is a 'fresh' variable.

# Ex.: Substitution

- $[y/x](\lambda y.x)$  — the occurrence of  $x$  is free  
 $\neq (\lambda y.y)$  — if we replace  $x$  with  $y$ , the variable  $y$  becomes bound.

# Ex.: Substitution

- $[y/x](\lambda y.x)$  — the occurrence of  $x$  is free  
 $\neq (\lambda y.y)$  — if we replace  $x$  with  $y$ , the variable  $y$  becomes bound.
- $[y/x](\lambda y.x)$  — the occurrence of  $x$  is free  
 $= (\lambda z[y/x][z/y]x)$  — we need a fresh variable  
 $= (\lambda z.y)$  — the occurrence of  $y$  is free

# Ex.: Substitution

- $[y/x](\lambda y.x)$  — the occurrence of  $x$  is free  
 $\neq (\lambda y.y)$  — if we replace  $x$  with  $y$ , the variable  $y$  becomes bound.
- $[y/x](\lambda y.x)$  — the occurrence of  $x$  is free  
 $= (\lambda z[y/x][z/y]x)$  — we need a fresh variable  
 $= (\lambda z.y)$  — the occurrence of  $y$  is free
- Further Examples on Blackboard

# Ex.: Substitution

- $[y/x](\lambda y.x)$  — the occurrence of  $x$  is free  
 $\neq (\lambda y.y)$  — if we replace  $x$  with  $y$ , the variable  $y$  becomes bound.
- $[y/x](\lambda y.x)$  — the occurrence of  $x$  is free  
 $= (\lambda z[y/x][z/y]x)$  — we need a fresh variable  
 $= (\lambda z.y)$  — the occurrence of  $y$  is free
- Further Examples on Blackboard
- Claim:  $[N/x]A = A$  if  $x \notin FV(A)$   
Proof: Induction on  $A$



# Def.: $\alpha$ -Conversion



$$[\lambda x. M] \rightarrow_{\alpha} [\lambda y. [y/x]M]$$

where  $y \notin FV(M)$

# Def.: $\alpha$ -Conversion

$$[\lambda x. M] \rightarrow_{\alpha} [\lambda y. [y/x]M]$$

where  $y \notin FV(M)$

$$A =^{\alpha} B$$

if  $A$  can be converted to  $B$  by renaming the bound variables. We read  $A =_{\alpha} B$  as  $A$  is  $\alpha$ -equal to  $B$ .

# Def.: $\alpha$ -Conversion

$$[\lambda x. M] \rightarrow_{\alpha} [\lambda y. [y/x]M]$$

where  $y \notin FV(M)$

$$A =^{\alpha} B$$

if  $A$  can be converted to  $B$  by renaming the bound variables. We read  $A =_{\alpha} B$  as  $A$  is  $\alpha$ -equal to  $B$ .

From now on  $(\lambda y. y) = (\lambda z. z)$ , that is, we will say that two terms are simply equal, if they are  $\alpha$ -equal. Two terms are equal means that two terms are  $\alpha$ -convertible.

# Def.: $\beta$ -Conversion



A  $\beta$ -redex is a term  $((\lambda x. A)B)$ . The  $\beta$ -reduct of this redex is  $[B/x]A$ .

# Def.: $\beta$ -Conversion



A  $\beta$ -redex is a term  $((\lambda x. A)B)$ . The  $\beta$ -reduct of this redex is  $[B/x]A$ .

We say  $M \rightarrow_{\beta} N$ , ie.  $\beta$ -reduces in 1 step, if

$$M = P[(\lambda x. A)B]_p$$

$$N = P[[B/x]A]_p$$

# Def.: $\beta$ -Conversion

A  $\beta$ -redex is a term  $((\lambda x. A)B)$ . The  $\beta$ -reduct of this redex is  $[B/x]A$ .

We say  $M \rightarrow_{\beta} N$ , ie.  $\beta$ -reduces in 1 step, if

$$M = P[(\lambda x. A)B]_p$$

$$N = P[[B/x]A]_p$$

We say  $M \rightarrow_{\beta}^* N$ , ie.  $\beta$ -reduces in several steps, if  $\exists M^1, \dots, M^n$  for  $n \geq 1$  such that  $M = M^1$  and  $N = M^n$  and  $M^i \rightarrow_{\beta} M^{i+1}$ .

# Def.: $\beta$ -Normal Form



A term is called  $\beta$ -normal if it contains no  $\beta$ -redexes.

# Def.: $\beta$ -Normal Form



A term is called  $\beta$ -normal if it contains no  $\beta$ -redexes.

Any term that does not contain  $\lambda$ -abstractions is  $\beta$ -normal.



# Def.: $\beta$ -Normal Form



A term is called  $\beta$ -normal if it contains no  $\beta$ -redexes.

Any term that does not contain  $\lambda$ -abstractions is  $\beta$ -normal.

A term is called  $\beta$ -head normal if the head term of its outermost application can not be further reduced.

# Def.: $\beta$ -Normal Form




A term is called  $\beta$ -normal if it contains no  $\beta$ -redexes.

Any term that does not contain  $\lambda$ -abstractions is  $\beta$ -normal.


A term is called  $\beta$ -head normal if the head term of its outermost application can not be further reduced.

Any term that does not contain  $\lambda$ -abstractions is  $\beta$ -head normal.

# Thm.: Church-Rosser Property for $\rightarrow_{\beta}$




# Thm.: Church-Rosser Property for $\rightarrow_{\beta}$



If  $T_{\alpha}$   $\beta$ -reduces in multiple steps with one strategy to  $L_{\alpha}$  and with another strategy to  $R_{\alpha}$  then there exists a term  $B_{\alpha}$  such that  $L_{\alpha}$  and  $R_{\alpha}$   $\beta$ -reduce in multiple steps to  $B_{\alpha}$ .


# Thm.: Church-Rosser Property for $\rightarrow_{\beta}$



If  $T_{\alpha}$   $\beta$ -reduces in multiple steps with one strategy to  $L_{\alpha}$  and with another strategy to  $R_{\alpha}$  then there exists a term  $B_{\alpha}$  such that  $L_{\alpha}$  and  $R_{\alpha}$   $\beta$ -reduce in multiple steps to  $B_{\alpha}$ .

Note that  $B_{\alpha}$  is not necessarily in normal form.

# Thm.: Church-Rosser Property for $\rightarrow_{\beta}$




If  $T_{\alpha}$   $\beta$ -reduces in multiple steps with one strategy to  $L_{\alpha}$  and with another strategy to  $R_{\alpha}$  then there exists a term  $B_{\alpha}$  such that  $L_{\alpha}$  and  $R_{\alpha}$   $\beta$ -reduce in multiple steps to  $B_{\alpha}$ .

Note that  $B_{\alpha}$  is not necessarily in normal form.

The Church-Rosser Property for  $\rightarrow_{\beta}$  holds for  $\Lambda$  and  $\Lambda^{\alpha}$ .

# Ex.: Church-Rosser Property for $\rightarrow_{\beta}$



Do we always get a  $\beta$ -normal form as we apply  $\beta$ -reduction?



Do we always get a  $\beta$ -normal form as we apply  $\beta$ -reduction?

**Typed Case:** For all  $A_\alpha$  there exists a unique (up to  $\alpha$ -conversion)  $\beta$ -normal term  $B$  such that  $A \rightarrow_\beta B$

Do we always get a  $\beta$ -normal form as we apply  $\beta$ -reduction?

**Typed Case:** For all  $A_\alpha$  there exists a unique (up to  $\alpha$ -conversion)  $\beta$ -normal term  $B$  such that  $A \rightarrow_\beta B$

**Untyped Case:** Consider the term  $\omega = (\lambda x. xx)$

$$(\lambda x. xx)(\lambda x. xx) \rightarrow_\beta^1 \omega\omega$$

# Def.: $\eta$ -Conversion



A  $\eta$ -redex is a term of the form  $(\lambda x_{\beta}. F_{\alpha\beta} x)$  where  $x \notin FV(F)$ . The  $\eta$ -reduct of this term is  $F$ .

# Def.: $\eta$ -Conversion

A  $\eta$ -redex is a term of the form  $(\lambda x_{\beta}. F_{\alpha\beta} x)$  where  $x \notin FV(F)$ . The  $\eta$ -reduct of this term is  $F$ .

We say  $M \rightarrow_{\eta} N$ , ie.  $\eta$ -reduces in 1 step, if

$$\begin{aligned} M &= P[(\lambda x_{\beta}. F_{\alpha\beta} x)]_p \\ N &= P[F]_p \end{aligned}$$

# Def.: $\eta$ -Conversion

A  $\eta$ -redex is a term of the form  $(\lambda x_{\beta}. F_{\alpha\beta} x)$  where  $x \notin FV(F)$ . The  $\eta$ -reduct of this term is  $F$ .

We say  $M \rightarrow_{\eta} N$ , ie.  $\eta$ -reduces in 1 step, if

$$\begin{aligned} M &= P[(\lambda x_{\beta}. F_{\alpha\beta} x)]_p \\ N &= P[F]_p \end{aligned}$$


We say  $M \twoheadrightarrow_{\eta} N$ , ie.  $\eta$ -reduces in several steps, if  $\exists M^1, \dots, M^n$  for  $n \geq 1$  such that  $M = M^1$  and  $N = M^n$  and  $M^i \rightarrow_{\beta} M^{i+1}$ .

# Def.: $\eta$ -Normal Form




A term is called  $\eta$ -normal if it contains no  $\eta$ -redexes.

# Thm.: Church-Rosser Property for $\rightarrow_{\eta}$




# Thm.: Church-Rosser Property for $\rightarrow_{\eta}$



If  $T_{\alpha}$   $\eta$ -reduces in multiple steps with one strategy to  $L_{\alpha}$  and with another strategy to  $R_{\alpha}$  then there exists a term  $B_{\alpha}$  such that  $L_{\alpha}$  and  $R_{\alpha}$   $\eta$ -reduce in multiple steps to  $B_{\alpha}$ .



# Thm.: Church-Rosser Property for $\rightarrow_{\eta}$



If  $T_{\alpha}$   $\eta$ -reduces in multiple steps with one strategy to  $L_{\alpha}$  and with another strategy to  $R_{\alpha}$  then there exists a term  $B_{\alpha}$  such that  $L_{\alpha}$  and  $R_{\alpha}$   $\eta$ -reduce in multiple steps to  $B_{\alpha}$ .

The Church-Rosser Property for  $\rightarrow_{\eta}$  holds for  $\Lambda$  and  $\Lambda^{\alpha}$ .

# Def.: $\beta\eta$ -Conversion



$$\rightarrow_{\beta\eta} := \rightarrow_{\beta} \cup \rightarrow_{\eta}$$

# Def.: $\beta\eta$ -Conversion



$$\rightarrow_{\beta\eta} := \rightarrow_{\beta} \cup \rightarrow_{\eta}$$

If  $M \rightarrow_{\beta\eta} N$  we say  $M$   $\beta\eta$ -reduces in 1 step to  $N$ .

# Def.: $\beta\eta$ -Conversion



$$\rightarrow_{\beta\eta} := \rightarrow_{\beta} \cup \rightarrow_{\eta}$$

If  $M \rightarrow_{\beta\eta} N$  we say  $M$   $\beta\eta$ -reduces in 1 step to  $N$ .


We say  $M \twoheadrightarrow_{\beta\eta} N$ , ie.  $\eta$ -reduces in several steps, if  $\exists M^1, \dots, M^n$  for  $n \geq 1$  such that  $M = M^1$  and  $N = M^n$  and  $M^i \rightarrow_{\beta\eta} M^{i+1}$ .

# Def.: $\beta\eta$ -Normal Form




A term is  $\beta\eta$ -normal if it contains no  $\beta$ -redexes and no  $\eta$ -redexes.

# Thm.: Church-Rosser Property for $\rightarrow_{\beta\eta}$




# Thm.: Church-Rosser Property for $\rightarrow_{\beta\eta}$



If  $T_\alpha$   $\beta\eta$ -reduces in multiple steps with one strategy to  $L_\alpha$  and with another strategy to  $R_\alpha$  then there exists a term  $B_\alpha$  such that  $L_\alpha$  and  $R_\alpha$   $\beta\eta$ -reduce in multiple steps to  $B_\alpha$ .

# Thm.: Church-Rosser Property for $\rightarrow_{\beta\eta}$



If  $T_\alpha$   $\beta\eta$ -reduces in multiple steps with one strategy to  $L_\alpha$  and with another strategy to  $R_\alpha$  then there exists a term  $B_\alpha$  such that  $L_\alpha$  and  $R_\alpha$   $\beta\eta$ -reduce in multiple steps to  $B_\alpha$ .

The Church-Rosser Property for  $\rightarrow_{\beta\eta}$  holds for  $\Lambda$  and  $\Lambda^\alpha$ .



# Thm.: Strong Church-Rosser Property



In  $\Lambda^\alpha$  (simply typed  $\lambda$ -calculus) the relations  $\rightarrow_\beta$  and  $\rightarrow_{\beta\eta}$  have the **strong Church Rosser property**:

# Thm.: Strong Church-Rosser Property



In  $\Lambda^\alpha$  (simply typed  $\lambda$ -calculus) the relations  $\rightarrow_\beta$  and  $\rightarrow_{\beta\eta}$  have the **strong Church Rosser property**: for every term  $A_\tau$  there exists a unique (up to  $\alpha$ -renaming)  $\beta$ -normal resp.  $\beta\eta$ -normal term  $B_\tau$  such that  $A_\tau \rightarrow_\beta B_\tau$  resp.  $A_\tau \rightarrow_{\beta\eta} B_\tau$ .

# Def.: Long $\beta\eta$ -Normal Form

Let  $n \geq 0$ ,  $\alpha^1, \dots, \alpha^n \in \mathcal{T}$ , and  $\beta \in \{\mathbf{o}, \iota\}$ . A term  $A$  of type  $(\beta, \alpha^n, \dots, \alpha^1)$  is in **long  $\beta\eta$ -normal form** if it is of form

$$\lambda x_{\alpha^1}^1 \dots x_{\alpha^n}^n. (h_{\beta\gamma^m \dots \gamma^1} A_{\gamma^1}^1 \dots A_{\gamma^m}^m)$$

for a variable or constant  $h_{\beta\gamma^m \dots \gamma^1}$ ,  $m \geq 0$  and long  $\beta\eta$ -normal forms  $A_{\gamma^1}^1, \dots, A_{\gamma^m}^m$ .

# Def.: Long $\beta\eta$ -Normal Form

Let  $n \geq 0$ ,  $\alpha^1, \dots, \alpha^n \in \mathcal{T}$ , and  $\beta \in \{o, \iota\}$ . A term  $A$  of type  $(\beta, \alpha^n, \dots, \alpha^1)$  is in **long  $\beta\eta$ -normal form** if it is of form

$$\lambda x_{\alpha^1}^1 \dots x_{\alpha^n}^n. (h_{\beta\gamma^m \dots \gamma^1} A_{\gamma^1}^1 \dots A_{\gamma^m}^m)$$

for a variable or constant  $h_{\beta\gamma^m \dots \gamma^1}$ ,  $m \geq 0$  and long  $\beta\eta$ -normal forms  $A_{\gamma^1}^1, \dots, A_{\gamma^m}^m$ . Note that this is an inductive definition; the base case is when  $m = 0$ .

# Def.: Long $\beta\eta$ -Normal Form

Let  $n \geq 0$ ,  $\alpha^1, \dots, \alpha^n \in \mathcal{T}$ , and  $\beta \in \{\mathbf{o}, \iota\}$ . A term  $A$  of type  $(\beta, \alpha^n, \dots, \alpha^1)$  is in **long  $\beta\eta$ -normal form** if it is of form

$$\lambda x_{\alpha^1}^1 \dots x_{\alpha^n}^n. (h_{\beta\gamma^m \dots \gamma^1} A_{\gamma^1}^1 \dots A_{\gamma^m}^m)$$

for a variable or constant  $h_{\beta\gamma^m \dots \gamma^1}$ ,  $m \geq 0$  and long  $\beta\eta$ -normal forms  $A_{\gamma^1}^1, \dots, A_{\gamma^m}^m$ . Note that this is an inductive definition; the base case is when  $m = 0$ . Note that if  $\lambda \bar{x}^n. (h \bar{A}^m)$  is in long  $\beta\eta$ -normal form then  $(h \bar{A}^m)$  is of base type.

# Ex.: Long $\beta\eta$ -Normal Form



Consider the  $\beta\eta$ -normal term  $f_{\iota(\iota)}$ .

$$\begin{array}{c} f_{\iota(\iota)} \\ \uparrow \eta \\ \lambda w_{\iota\iota}. (f_{\iota(\iota)} w_{\iota\iota}) \\ \uparrow \eta \\ \lambda w_{\iota\iota}. (f(\lambda x_{\iota}. w_{\iota\iota} x)) \end{array}$$

# Thm.: Long $\beta\eta$ -Normal Form



For every term  $A$  there is unique long  $\beta\eta$ -normal form  $B$  such that  $A =^{\beta\eta} B$ .

# Rem.: $\beta\eta$ -Head Normal Form



Instead of terms in long  $\beta\eta$ -normal form we often use in practice terms in  $\beta\eta$ -head normal form.



# Rem.: $\beta\eta$ -Head Normal Form



Instead of terms in long  $\beta\eta$ -normal form we often use in practice terms in  **$\beta\eta$ -head normal form**. Definition is similar to long  $\beta\eta$ -normal, but we do not require the embedded terms  $A_{\gamma^i}^i$  to be in normal form.

- $A \downarrow_{\beta}$  is the  $\beta$ -normal form of  $A$ .

- $A \downarrow_{\beta}$  is the  $\beta$ -normal form of  $A$ .
- $A \downarrow_{\eta}$  is the  $\eta$ -normal form of  $A$ .

- $A \downarrow_{\beta}$  is the  $\beta$ -normal form of  $A$ .
- $A \downarrow_{\eta}$  is the  $\eta$ -normal form of  $A$ .
- $A \downarrow$  is the  $\beta\eta$ -normal form of  $A$ .

- $A \downarrow_{\beta}$  is the  $\beta$ -normal form of  $A$ .
- $A \downarrow_{\eta}$  is the  $\eta$ -normal form of  $A$ .
- $A \downarrow$  is the  $\beta\eta$ -normal form of  $A$ .
- $A \updownarrow$  is the long  $\beta\eta$ -normal form of  $A$ .



Semantics:  $\Sigma$ -Evaluations

# Ex.: An Interesting Applicative Structure



$$D_\alpha := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed}\}.$$

- Is  $D_\alpha$  non-empty for all  $\alpha$ ?

# Ex.: An Interesting Applicative Structure



$$D_\alpha := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed}\}.$$

- Is  $D_\alpha$  non-empty for all  $\alpha$ ?
- If  $\mathcal{C}_i \neq \emptyset$  and  $\mathcal{C}_o \neq \emptyset$ , then  $\forall \alpha \in \mathcal{T}. \Lambda_\alpha \neq \emptyset$ .



# Ex.: An Interesting Applicative Structure



$$D_\alpha := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed}\}.$$

- Is  $D_\alpha$  non-empty for all  $\alpha$ ?
- If  $\mathcal{C}_i \neq \emptyset$  and  $\mathcal{C}_o \neq \emptyset$ , then  $\forall \alpha \in \mathcal{T}. \Lambda_\alpha \neq \emptyset$ .
- Is  $D_{\alpha\beta}$  a set of functions? (ie.  $D_{\alpha\beta} \subseteq (D_\alpha)^{D_\beta}$ ?) — No!

# Ex.: An Interesting Applicative Structure



$D_\alpha := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed}\}.$

- Is  $D_\alpha$  non-empty for all  $\alpha$ ?
- If  $\mathcal{C}_i \neq \emptyset$  and  $\mathcal{C}_o \neq \emptyset$ , then  $\forall \alpha \in \mathcal{T}. \Lambda_\alpha \neq \emptyset.$
- Is  $D_{\alpha\beta}$  a set of functions? (ie.  $D_{\alpha\beta} \subseteq (D_\alpha)^{D_\beta}$ ?) — No!
- Is  $(\lambda x_i x) \in D_{ii}$ ? — Yes!

# Ex.: An Interesting Applicative Structure



$$D_\alpha := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed}\}.$$

- Is  $D_\alpha$  non-empty for all  $\alpha$ ?
- If  $\mathcal{C}_i \neq \emptyset$  and  $\mathcal{C}_o \neq \emptyset$ , then  $\forall \alpha \in \mathcal{T}. \Lambda_\alpha \neq \emptyset$ .
- Is  $D_{\alpha\beta}$  a set of functions? (ie.  $D_{\alpha\beta} \subseteq (D_\alpha)^{D_\beta}$ ?) — No!
- Is  $(\lambda x_i. x) \in D_{ii}$ ? — Yes!
- $D = (D_\alpha)_{\alpha \in \mathcal{T}}$  is not a frame!

# Ex.: An Interesting Applicative Structure



$$D_\alpha := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed}\}.$$

- Is  $D_\alpha$  non-empty for all  $\alpha$ ?
- If  $\mathcal{C}_i \neq \emptyset$  and  $\mathcal{C}_o \neq \emptyset$ , then  $\forall \alpha \in \mathcal{T}. \Lambda_\alpha \neq \emptyset$ .
- Is  $D_{\alpha\beta}$  a set of functions? (ie.  $D_{\alpha\beta} \subseteq (D_\alpha)^{D_\beta}$ ?) — No!
- Is  $(\lambda x_i x) \in D_{ii}$ ? — Yes!
- $D = (D_\alpha)_{\alpha \in \mathcal{T}}$  is not a frame!
- It requires a specific application operator  $@ : D_{\alpha\beta} \times D_\beta \rightarrow D_\alpha$

# Ex.: An Interesting Applicative Structure



$$D_\alpha := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed}\}.$$

- Is  $D_\alpha$  non-empty for all  $\alpha$ ?
- If  $\mathcal{C}_i \neq \emptyset$  and  $\mathcal{C}_o \neq \emptyset$ , then  $\forall \alpha \in \mathcal{T}. \Lambda_\alpha \neq \emptyset$ .
- Is  $D_{\alpha\beta}$  a set of functions? (ie.  $D_{\alpha\beta} \subseteq (D_\alpha)^{D_\beta}$ ?) — No!
- Is  $(\lambda x_i x) \in D_{ii}$ ? — Yes!
- $D = (D_\alpha)_{\alpha \in \mathcal{T}}$  is not a frame!
- It requires a specific application operator  $@ : D_{\alpha\beta} \times D_\beta \rightarrow D_\alpha$
- If  $\Lambda_\alpha$  is non-empty for all  $\alpha \in \mathcal{T}$ , then  $\langle D, @ \rangle$  is an applicative structure.

# Ex.: Interpretation of Terms



Syntax      Semantics       $\langle D, @ \rangle$   
 $(\lambda x_\iota. x)$

# Ex.: Interpretation of Terms



Syntax	Semantics	$\langle D, @ \rangle$
$(\lambda x_\iota. x)$	$(\lambda x_\iota. x)$	

# Ex.: Interpretation of Terms



Syntax	Semantics	$\langle D, @ \rangle$
$(\lambda x_{\iota}. x)$	$(\lambda x_{\iota}. x)$	$\in D_{\iota\iota}$



# Ex.: Interpretation of Terms



Syntax	Semantics	$\langle D, @ \rangle$
$(\lambda x_\iota. x)$	$(\lambda x_\iota. x)$	$\in D_{\iota\iota}$
$y_\iota$		

# Ex.: Interpretation of Terms



Syntax	Semantics	$\langle D, @ \rangle$
$(\lambda x_\iota. x)$	$(\lambda x_\iota. x)$	$\in D_{\iota\iota}$
$y_\iota$	$\varphi(y)$	

# Ex.: Interpretation of Terms



Syntax	Semantics	$\langle D, @ \rangle$
$(\lambda x_\iota. x)$	$(\lambda x_\iota. x)$	$\in D_{\iota\iota}$
$y_\iota$	$\varphi(y)$	$\in D_\iota$

# Ex.: Interpretation of Terms



Syntax	Semantics	$\langle D, @ \rangle$
$(\lambda x_\iota. x)$	$(\lambda x_\iota. x)$	$\in D_{\iota\iota}$
$y_\iota$	$\varphi(y)$	$\in D_\iota$
$a_\iota \in C$		

# Ex.: Interpretation of Terms



Syntax	Semantics	$\langle D, @ \rangle$
$(\lambda x_\iota. x)$	$(\lambda x_\iota. x)$	$\in D_{\iota\iota}$
$y_\iota$	$\varphi(y)$	$\in D_\iota$
$a_\iota \in C$	$a$	

# Ex.: Interpretation of Terms



Syntax	Semantics	$\langle D, @ \rangle$
$(\lambda x_\iota. x)$	$(\lambda x_\iota. x)$	$\in D_{\iota\iota}$
$y_\iota$	$\varphi(y)$	$\in D_\iota$
$a_\iota \in C$	$a$	$\in D_\iota$

# Ex.: Interpretation of Terms



Syntax	Semantics	$\langle D, @ \rangle$
$(\lambda x_\iota. x)$	$(\lambda x_\iota. x)$	$\in D_{\iota\iota}$
$y_\iota$	$\varphi(y)$	$\in D_\iota$
$a_\iota \in C$	$a$	$\in D_\iota$
$(\lambda x_\iota. x)a_\iota$		

# Ex.: Interpretation of Terms



Syntax	Semantics	$\langle D, @ \rangle$
$(\lambda x_\iota. x)$	$(\lambda x_\iota. x)$	$\in D_{\iota\iota}$
$y_\iota$	$\varphi(y)$	$\in D_\iota$
$a_\iota \in C$	$a$	$\in D_\iota$
$(\lambda x_\iota. x)a_\iota$	$(\lambda x_\iota. x)@a_\iota$	



# Ex.: Interpretation of Terms



Syntax	Semantics	$\langle D, @ \rangle$
$(\lambda x_\iota. x)$	$(\lambda x_\iota. x)$	$\in D_\iota$
$y_\iota$	$\varphi(y)$	$\in D_\iota$
$a_\iota \in C$	$a$	$\in D_\iota$
$(\lambda x_\iota. x)a_\iota$	$(\lambda x_\iota. x)@a_\iota$	$\in D_\iota$

# Ex.: Interpretation of Terms

Syntax	Semantics	$\langle D, @ \rangle$
$(\lambda x_\iota. x)$	$(\lambda x_\iota. x)$	$\in D_{\iota\iota}$
$y_\iota$	$\varphi(y)$	$\in D_\iota$
$a_\iota \in C$	$a$	$\in D_\iota$
$(\lambda x_\iota. x)a_\iota$	$(\lambda x_\iota. x)@a_\iota$	$\in D_\iota$

Remark: The variable  $y_\iota$  is a non-closed well-formed formula of type  $\iota$ . We need an assignment  $\varphi_\alpha : V_\alpha \rightarrow D_\alpha$  to give it a meaning.

# Ex.: Interesting Applicative Structures



- Let  $D_\alpha \downarrow_\beta := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed and } A \text{ is in } \beta\text{-normal form}\}$

# Ex.: Interesting Applicative Structures



- Let  $D_\alpha \downarrow_\beta := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed and } A \text{ is in } \beta\text{-normal form}\}$
- Let  $D := (D_\alpha \downarrow_\beta)_{\alpha \in \mathcal{T}}$

# Ex.: Interesting Applicative Structures



- Let  $D_\alpha \downarrow_\beta := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed and } A \text{ is in } \beta\text{-normal form}\}$
- Let  $D := (D_\alpha \downarrow_\beta)_{\alpha \in \mathcal{T}}$
- Let  $@_{\gamma\delta}^\beta : D_{\gamma\delta} \times D_\delta \rightarrow D_\gamma$  be defined by

$$F_{\gamma\delta} @_{\gamma\delta}^\beta G_\delta = (F G) \downarrow_\beta$$

for all  $F_{\gamma\delta} \in D_{\gamma\delta}$  and  $G_\delta \in D_\delta$ .

# Ex.: Interesting Applicative Structures



- Let  $D_\alpha \downarrow_\beta := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed and } A \text{ is in } \beta\text{-normal form}\}$
- Let  $D := (D_\alpha \downarrow_\beta)_{\alpha \in \mathcal{T}}$
- Let  $@_{\gamma\delta}^\beta : D_{\gamma\delta} \times D_\delta \rightarrow D_\gamma$  be defined by

$$F_{\gamma\delta} @_{\gamma\delta}^\beta G_\delta = (F G) \downarrow_\beta$$

for all  $F_{\gamma\delta} \in D_{\gamma\delta}$  and  $G_\delta \in D_\delta$ .

- $@^\beta = (@_{\gamma\delta}^\beta)_{\gamma\delta \in \mathcal{T}}$

# Ex.: Interesting Applicative Structures

- Let  $D_\alpha \downarrow_\beta := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed and } A \text{ is in } \beta\text{-normal form}\}$
- Let  $D := (D_\alpha \downarrow_\beta)_{\alpha \in \mathcal{T}}$
- Let  $@_{\gamma\delta}^\beta : D_{\gamma\delta} \times D_\delta \rightarrow D_\gamma$  be defined by

$$F_{\gamma\delta} @_{\gamma\delta}^\beta G_\delta = (F G) \downarrow_\beta$$

for all  $F_{\gamma\delta} \in D_{\gamma\delta}$  and  $G_\delta \in D_\delta$ .

- $@^\beta = (@_{\gamma\delta}^\beta)_{\gamma\delta \in \mathcal{T}}$

Claim: If  $\mathcal{C}_i \neq \emptyset$  and  $\mathcal{C}_o \neq \emptyset$  (i.e., at least one constant for each base type is given), then  $(D, @^\beta)$  is an applicative structure.

# Ex.: Interesting Applicative Structures



Proof:

- Is  $D_\alpha \downarrow_\beta$  nonempty for all  $\alpha \in \mathcal{T}$ ?



# Ex.: Interesting Applicative Structures



Proof:

- Is  $D_\alpha \downarrow_\beta$  nonempty for all  $\alpha \in \mathcal{T}$ ?
- Yes! This follows since  $\mathcal{C}_\iota \neq \emptyset$  and  $\mathcal{C}_\lambda \neq \emptyset$ .

# Ex.: Interesting Applicative Structures



Proof:

- Is  $D_\alpha \downarrow_\beta$  nonempty for all  $\alpha \in \mathcal{T}$ ?
- Yes! This follows since  $\mathcal{C}_\iota \neq \emptyset$  and  $\mathcal{C}_\iota \neq \emptyset$ .
- Is  $F_{\gamma\delta} @_{\gamma\delta}^\beta G_\delta \in D_\gamma \downarrow_\beta$ ?

# Ex.: Interesting Applicative Structures



Proof:

- Is  $D_\alpha \downarrow_\beta$  nonempty for all  $\alpha \in \mathcal{T}$ ?
- Yes! This follows since  $\mathcal{C}_\iota \neq \emptyset$  and  $\mathcal{C}_\lambda \neq \emptyset$ .
- Is  $F_{\gamma\delta} @_{\gamma\delta}^\beta G_\delta \in D_\gamma \downarrow_\beta$ ?
- Let's check:  $F_{\gamma\delta} @_{\gamma\delta}^\beta G_\delta = (F G) \downarrow_\beta \in D_\gamma \downarrow_\beta$

# Ex.: Interesting Applicative Structures



- Let  $D_\alpha \downarrow_{\beta\eta} := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed and } A \text{ is in } \beta\eta\text{-normal form}\}$

# Ex.: Interesting Applicative Structures



- Let  $D_\alpha \downarrow_{\beta\eta} := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed and } A \text{ is in } \beta\eta\text{-normal form}\}$
- Let  $D := (D_\alpha \downarrow_{\beta\eta})_{\alpha \in \mathcal{T}}$

# Ex.: Interesting Applicative Structures



- Let  $D_\alpha \downarrow_{\beta\eta} := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed and } A \text{ is in } \beta\eta\text{-normal form}\}$
- Let  $D := (D_\alpha \downarrow_{\beta\eta})_{\alpha \in \mathcal{T}}$
- Let  $@_{\gamma\delta}^{\beta\eta} : D_{\gamma\delta} \times D_\delta \rightarrow D_\gamma$  be defined by

$$F_{\gamma\delta} @_{\gamma\delta}^{\beta\eta} G_\delta = (F G) \downarrow$$

for all  $F_{\gamma\delta} \in D_{\gamma\delta}$  and  $G_\delta \in D_\delta$ .

# Ex.: Interesting Applicative Structures

- Let  $D_\alpha \downarrow_{\beta\eta} := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed and } A \text{ is in } \beta\eta\text{-normal form}\}$
- Let  $D := (D_\alpha \downarrow_{\beta\eta})_{\alpha \in \mathcal{T}}$
- Let  $@_{\gamma\delta}^{\beta\eta} : D_{\gamma\delta} \times D_\delta \rightarrow D_\gamma$  be defined by

$$F_{\gamma\delta} @_{\gamma\delta}^{\beta\eta} G_\delta = (F G) \downarrow$$

for all  $F_{\gamma\delta} \in D_{\gamma\delta}$  and  $G_\delta \in D_\delta$ .

- $@^{\beta\eta} = (@_{\gamma\delta}^{\beta\eta})_{\gamma\delta \in \mathcal{T}}$

# Ex.: Interesting Applicative Structures

- Let  $D_\alpha \downarrow_{\beta\eta} := \{A_\alpha \in \Lambda_\alpha \mid A \text{ is closed and } A \text{ is in } \beta\eta\text{-normal form}\}$
- Let  $D := (D_\alpha \downarrow_{\beta\eta})_{\alpha \in \mathcal{T}}$
- Let  $@_{\gamma\delta}^{\beta\eta} : D_{\gamma\delta} \times D_\delta \rightarrow D_\gamma$  be defined by

$$F_{\gamma\delta} @_{\gamma\delta}^{\beta\eta} G_\delta = (F G) \downarrow$$

for all  $F_{\gamma\delta} \in D_{\gamma\delta}$  and  $G_\delta \in D_\delta$ .

- $@^{\beta\eta} = (@_{\gamma\delta}^{\beta\eta})_{\gamma\delta \in \mathcal{T}}$

Claim: If  $\mathcal{C}_\iota \neq \emptyset$  and  $\mathcal{C}_o \neq \emptyset$  (i.e., at least one constant for each base type is given), then  $(D, @^{\beta\eta})$  is an applicative structure.



# Ex.: Interesting Applicative Structures



Proof:

- ...analogous ...

# Def.: Variable Assignment



Let  $\mathcal{A} := (\mathcal{D}, @)$  be an applicative structure.

# Def.: Variable Assignment

Let  $\mathcal{A} := (\mathcal{D}, @)$  be an applicative structure.

A typed function  $\varphi: \mathcal{V} \longrightarrow \mathcal{D} := (\varphi_\alpha: \mathcal{V}_\alpha \longrightarrow \mathcal{D}_\alpha)_{\alpha \in \mathcal{T}}$  is called a **variable assignment** into  $\mathcal{A}$ .

# Def.: Variable Assignment

Let  $\mathcal{A} := (\mathcal{D}, @)$  be an applicative structure.

A typed function  $\varphi: \mathcal{V} \longrightarrow \mathcal{D} := (\varphi_\alpha: \mathcal{V}_\alpha \longrightarrow \mathcal{D}_\alpha)_{\alpha \in \mathcal{T}}$  is called a **variable assignment** into  $\mathcal{A}$ .

Given a variable assignment  $\varphi$ , variable  $X_\alpha$ , and value  $a \in \mathcal{D}_\alpha$ ,

# Def.: Variable Assignment

Let  $\mathcal{A} := (\mathcal{D}, @)$  be an applicative structure.

A typed function  $\varphi: \mathcal{V} \longrightarrow \mathcal{D} := (\varphi_\alpha: \mathcal{V}_\alpha \longrightarrow \mathcal{D}_\alpha)_{\alpha \in \mathcal{T}}$  is called a **variable assignment** into  $\mathcal{A}$ .

Given a variable assignment  $\varphi$ , variable  $X_\alpha$ , and value  $a \in \mathcal{D}_\alpha$ , we use  $\varphi, [a/X]$  to denote the variable assignment with

# Def.: Variable Assignment

Let  $\mathcal{A} := (\mathcal{D}, @)$  be an applicative structure.

A typed function  $\varphi: \mathcal{V} \longrightarrow \mathcal{D} := (\varphi_\alpha: \mathcal{V}_\alpha \longrightarrow \mathcal{D}_\alpha)_{\alpha \in \mathcal{T}}$  is called a **variable assignment** into  $\mathcal{A}$ .

Given a variable assignment  $\varphi$ , variable  $X_\alpha$ , and value  $a \in \mathcal{D}_\alpha$ , we use  $\varphi, [a/X]$  to denote the variable assignment with

$$(\varphi, [a/X])(X) = a$$

# Def.: Variable Assignment

Let  $\mathcal{A} := (\mathcal{D}, @)$  be an applicative structure.

A typed function  $\varphi: \mathcal{V} \longrightarrow \mathcal{D} := (\varphi_\alpha: \mathcal{V}_\alpha \longrightarrow \mathcal{D}_\alpha)_{\alpha \in \mathcal{T}}$  is called a **variable assignment** into  $\mathcal{A}$ .

Given a variable assignment  $\varphi$ , variable  $X_\alpha$ , and value  $a \in \mathcal{D}_\alpha$ , we use  $\varphi, [a/X]$  to denote the variable assignment with

$$(\varphi, [a/X])(X) = a$$

and

$$(\varphi, [a/X])(Y) = \varphi(Y)$$

for variables  $Y$  other than  $X$ .

# Some Assumptions

---



From now on, we assume the signature  $\Sigma_\alpha = (\mathcal{V}, \mathcal{C})$  to be infinite for each type  $\alpha$ .



# Some Assumptions



From now on, we assume the signature  $\Sigma_\alpha = (\mathcal{V}, \mathcal{C})$  to be infinite for each type  $\alpha$ . Furthermore, we assume there is a particular cardinal  $\aleph_s$  such that  $\Sigma_\alpha$  has cardinality  $\aleph_s$  for every type  $\alpha$ .

# Some Assumptions

From now on, we assume the signature  $\Sigma_\alpha = (\mathcal{V}, \mathcal{C})$  to be infinite for each type  $\alpha$ . Furthermore, we assume there is a particular cardinal  $\aleph_s$  such that  $\Sigma_\alpha$  has cardinality  $\aleph_s$  for every type  $\alpha$ . Since  $\mathcal{V}$  is countable, this implies  $wff_\alpha(\Sigma) := \Lambda^\alpha$  and  $cwff_\alpha(\Sigma) := \{A \in \Lambda^\alpha \mid A \text{ closed}\}$  have cardinality  $\aleph_s$  for each type  $\alpha$ .

# Some Assumptions

From now on, we assume the signature  $\Sigma_\alpha = (\mathcal{V}, \mathcal{C})$  to be infinite for each type  $\alpha$ . Furthermore, we assume there is a particular cardinal  $\aleph_s$  such that  $\Sigma_\alpha$  has cardinality  $\aleph_s$  for every type  $\alpha$ . Since  $\mathcal{V}$  is countable, this implies  $wff_\alpha(\Sigma) := \Lambda^\alpha$  and  $cwff_\alpha(\Sigma) := \{A \in \Lambda^\alpha \mid A \text{ closed}\}$  have cardinality  $\aleph_s$  for each type  $\alpha$ . Also, whether or not primitive equality is included in the signature, there can only be finitely many logical constants in  $\Sigma_\alpha$  for each particular type  $\alpha$ .

# Some Assumptions

From now on, we assume the signature  $\Sigma_\alpha = (\mathcal{V}, \mathcal{C})$  to be infinite for each type  $\alpha$ . Furthermore, we assume there is a particular cardinal  $\aleph_s$  such that  $\Sigma_\alpha$  has cardinality  $\aleph_s$  for every type  $\alpha$ . Since  $\mathcal{V}$  is countable, this implies  $wff_\alpha(\Sigma) := \Lambda^\alpha$  and  $cwff_\alpha(\Sigma) := \{A \in \Lambda^\alpha \mid A \text{ closed}\}$  have cardinality  $\aleph_s$  for each type  $\alpha$ . Also, whether or not primitive equality is included in the signature, there can only be finitely many logical constants in  $\Sigma_\alpha$  for each particular type  $\alpha$ . Thus, the cardinality of the set of parameters in  $\Sigma_\alpha$  is also  $\aleph_s$ . In the countable case,  $\aleph_s$  is  $\aleph_0$ .

# $\Sigma$ -Evaluations

---



Let  $\Sigma$  be a signature.

Let  $\Sigma$  be a signature. We build on the notion of applicative structures to **define  $\Sigma$ -evaluations**, where the evaluation function is assumed to **respect application and  $\beta$ -conversion**.

Let  $\Sigma$  be a signature. We build on the notion of applicative structures to **define  $\Sigma$ -evaluations**, where the evaluation function is assumed to **respect application and  $\beta$ -conversion**.

**In such models, a function is not uniquely determined by its behavior on all possible arguments.**

Let  $\Sigma$  be a signature. We build on the notion of applicative structures to **define  $\Sigma$ -evaluations**, where the evaluation function is assumed to **respect application and  $\beta$ -conversion**.

**In such models, a function is not uniquely determined by its behavior on all possible arguments.**

Such models can be constructed, for example, by labeling for functions (e.g., a green and a red version of a function  $f$ ) in order to differentiate between them, even though they are functionally equivalent.



Let  $\mathcal{E}: \mathcal{F}_{\mathcal{T}}(\mathcal{V}; \mathcal{D}) \longrightarrow \mathcal{F}_{\mathcal{T}}(wff(\Sigma), \mathcal{D})$  be a total function, where  $\mathcal{F}_{\mathcal{T}}(\mathcal{V}; \mathcal{D})$  is the set of variable assignments and  $\mathcal{F}_{\mathcal{T}}(wff(\Sigma), \mathcal{D})$  is the set of typed functions mapping terms into objects in  $\mathcal{D}$ .

Let  $\mathcal{E}: \mathcal{F}_{\mathcal{T}}(\mathcal{V}; \mathcal{D}) \longrightarrow \mathcal{F}_{\mathcal{T}}(wff(\Sigma), \mathcal{D})$  be a total function, where  $\mathcal{F}_{\mathcal{T}}(\mathcal{V}; \mathcal{D})$  is the set of variable assignments and  $\mathcal{F}_{\mathcal{T}}(wff(\Sigma), \mathcal{D})$  is the set of typed functions mapping terms into objects in  $\mathcal{D}$ . We will write the argument of  $\mathcal{E}$  as a subscript. So, for each assignment  $\varphi$ , we have a typed function

$$\mathcal{E}_{\varphi}: wff(\Sigma) \longrightarrow \mathcal{D}$$

Let  $\mathcal{E}: \mathcal{F}_{\mathcal{T}}(\mathcal{V}; \mathcal{D}) \longrightarrow \mathcal{F}_{\mathcal{T}}(wff(\Sigma), \mathcal{D})$  be a total function, where  $\mathcal{F}_{\mathcal{T}}(\mathcal{V}; \mathcal{D})$  is the set of variable assignments and  $\mathcal{F}_{\mathcal{T}}(wff(\Sigma), \mathcal{D})$  is the set of typed functions mapping terms into objects in  $\mathcal{D}$ . We will write the argument of  $\mathcal{E}$  as a subscript. So, for each assignment  $\varphi$ , we have a typed function

$$\mathcal{E}_{\varphi}: wff(\Sigma) \longrightarrow \mathcal{D}$$

What properties shall  $\mathcal{E}$  fulfill?

# Def.: Evaluation Function

$\mathcal{E}$  is called an **evaluation function** for an applicative structure

$$\mathcal{A} = (\mathcal{D}, @)$$

# Def.: Evaluation Function

$\mathcal{E}$  is called an **evaluation function** for an applicative structure  $\mathcal{A} = (\mathcal{D}, @)$  if for any assignments  $\varphi$  and  $\psi$  into  $\mathcal{A}$ , we have

# Def.: Evaluation Function

$\mathcal{E}$  is called an **evaluation function** for an applicative structure  $\mathcal{A} = (\mathcal{D}, @)$  if for any assignments  $\varphi$  and  $\psi$  into  $\mathcal{A}$ , we have

1.  $\mathcal{E}_\varphi|_\nu = \varphi$

# Def.: Evaluation Function

$\mathcal{E}$  is called an **evaluation function** for an applicative structure  $\mathcal{A} = (\mathcal{D}, @)$  if for any assignments  $\varphi$  and  $\psi$  into  $\mathcal{A}$ , we have

1.  $\mathcal{E}_\varphi|_{\mathcal{V}} = \varphi$
2.  $\mathcal{E}_\varphi(\mathbf{FA}) = \mathcal{E}_\varphi(\mathbf{F})@ \mathcal{E}_\varphi(\mathbf{A})$  for any  $\mathbf{F} \in wff_{\alpha \rightarrow \beta}(\Sigma)$  and  $\mathbf{A} \in wff_\alpha(\Sigma)$  and types  $\alpha$  and  $\beta$ .

# Def.: Evaluation Function

$\mathcal{E}$  is called an **evaluation function** for an applicative structure  $\mathcal{A} = (\mathcal{D}, @)$  if for any assignments  $\varphi$  and  $\psi$  into  $\mathcal{A}$ , we have

1.  $\mathcal{E}_\varphi|_{\mathcal{V}} = \varphi$
2.  $\mathcal{E}_\varphi(\mathbf{F}\mathbf{A}) = \mathcal{E}_\varphi(\mathbf{F})@\mathcal{E}_\varphi(\mathbf{A})$  for any  $\mathbf{F} \in wff_{\alpha \rightarrow \beta}(\Sigma)$  and  $\mathbf{A} \in wff_\alpha(\Sigma)$  and types  $\alpha$  and  $\beta$ .
3.  $\mathcal{E}_\varphi(\mathbf{A}) = \mathcal{E}_\psi(\mathbf{A})$  for any type  $\alpha$  and  $\mathbf{A} \in wff_\alpha(\Sigma)$ , whenever  $\varphi$  and  $\psi$  coincide on  $FV(\mathbf{A})$ .



# Def.: Evaluation Function

$\mathcal{E}$  is called an **evaluation function** for an applicative structure  $\mathcal{A} = (\mathcal{D}, @)$  if for any assignments  $\varphi$  and  $\psi$  into  $\mathcal{A}$ , we have

1.  $\mathcal{E}_\varphi|_{\mathcal{V}} = \varphi$
2.  $\mathcal{E}_\varphi(\mathbf{F}\mathbf{A}) = \mathcal{E}_\varphi(\mathbf{F})@\mathcal{E}_\varphi(\mathbf{A})$  for any  $\mathbf{F} \in wff_{\alpha \rightarrow \beta}(\Sigma)$  and  $\mathbf{A} \in wff_\alpha(\Sigma)$  and types  $\alpha$  and  $\beta$ .
3.  $\mathcal{E}_\varphi(\mathbf{A}) = \mathcal{E}_\psi(\mathbf{A})$  for any type  $\alpha$  and  $\mathbf{A} \in wff_\alpha(\Sigma)$ , whenever  $\varphi$  and  $\psi$  coincide on  $FV(\mathbf{A})$ .
4.  $\mathcal{E}_\varphi(\mathbf{A}) = \mathcal{E}_\varphi(\mathbf{A}|_\beta)$  for all  $\mathbf{A} \in wff_\alpha(\Sigma)$ .

# Def.: $\Sigma$ -Evaluation

We call  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  a  $\Sigma$ -evaluation if  $(\mathcal{D}, @)$  is an applicative structure and  $\mathcal{E}$  is an evaluation function for  $(\mathcal{D}, @)$ . We call  $\mathcal{E}_\varphi(\mathbf{A}_\alpha) \in \mathcal{D}_\alpha$  the denotation of  $\mathbf{A}_\alpha$  in  $\mathcal{J}$  for  $\varphi$ .

# Def.: $\Sigma$ -Evaluation

We call  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  a  $\Sigma$ -evaluation if  $(\mathcal{D}, @)$  is an applicative structure and  $\mathcal{E}$  is an evaluation function for  $(\mathcal{D}, @)$ . We call  $\mathcal{E}_\varphi(\mathbf{A}_\alpha) \in \mathcal{D}_\alpha$  the denotation of  $\mathbf{A}_\alpha$  in  $\mathcal{J}$  for  $\varphi$ .

Remark: since  $\mathcal{E}$  is a function, the denotation in  $\mathcal{J}$  is unique. However, for a given applicative structure  $\mathcal{A}$ , there may be many possible evaluation functions.

# Def.: $\Sigma$ -Evaluation

We call  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  a  $\Sigma$ -evaluation if  $(\mathcal{D}, @)$  is an applicative structure and  $\mathcal{E}$  is an evaluation function for  $(\mathcal{D}, @)$ . We call  $\mathcal{E}_\varphi(\mathbf{A}_\alpha) \in \mathcal{D}_\alpha$  the denotation of  $\mathbf{A}_\alpha$  in  $\mathcal{J}$  for  $\varphi$ .

Remark: since  $\mathcal{E}$  is a function, the denotation in  $\mathcal{J}$  is unique. However, for a given applicative structure  $\mathcal{A}$ , there may be many possible evaluation functions.

If  $\mathbf{A}$  is a closed formula, then  $\mathcal{E}_\varphi(\mathbf{A})$  is independent of  $\varphi$ , since  $\text{Free}(\mathbf{A}) = \emptyset$ . In these cases we sometimes drop the reference to  $\varphi$  from  $\mathcal{E}_\varphi(\mathbf{A})$  and simply write  $\mathcal{E}(\mathbf{A})$ .

# Def.: Functional/Full/Standard $\Sigma$ -Eval.



We call a  $\Sigma$ -evaluation  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  **functional** [**full**, **standard**] if the applicative structure  $(\mathcal{D}, @)$  is **functional** [**full**, **standard**].

# Def.: Functional/Full/Standard $\Sigma$ -Eval.



We call a  $\Sigma$ -evaluation  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  **functional** [**full**, **standard**] if the applicative structure  $(\mathcal{D}, @)$  is **functional** [**full**, **standard**].

We say  $\mathcal{J}$  is a  **$\Sigma$ -evaluation over a frame** if  $(\mathcal{D}, @)$  is a frame.

# What is the Idea?

---



$\Sigma$ -evaluations **generalize  $\Sigma$ -evaluations over frames**, which are the basis for Henkin models, **to the non-functional case**.

# What is the Idea?



$\Sigma$ -evaluations **generalize  $\Sigma$ -evaluations over frames**, which are the basis for Henkin models, **to the non-functional case**.

The existence of an evaluation function that meets the conditions as presented seems to be the weakest situation where one would like to speak of a model.



# What is the Idea?



$\Sigma$ -evaluations **generalize  $\Sigma$ -evaluations over frames**, which are the basis for Henkin models, **to the non-functional case**.

The existence of an evaluation function that meets the conditions as presented seems to be the weakest situation where one would like to speak of a model.

We cannot in general assume the evaluation function is uniquely determined by its values on constants as this requires functionality.

# What is the Idea?

$\Sigma$ -evaluations **generalize  $\Sigma$ -evaluations over frames**, which are the basis for Henkin models, **to the non-functional case**.

The existence of an evaluation function that meets the conditions as presented seems to be the weakest situation where one would like to speak of a model.

We cannot in general assume the evaluation function is uniquely determined by its values on constants as this requires functionality. Example: two evaluation functions  $\mathcal{E}$  and  $\mathcal{E}'$  on the same applicative structure may agree on all constants, but give a different value to the term  $(\lambda X.X)$ .

# Lemma: $\Sigma$ -Evaluations respect $\beta$ -Equality



Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and  $A =_{\beta} B$ . For all assignments  $\varphi$  into  $(\mathcal{D}, @)$ , we have

▪

# Lemma: $\Sigma$ -Evaluations respect $\beta$ -Equality



Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and  $A =_{\beta} B$ . For all assignments  $\varphi$  into  $(\mathcal{D}, @)$ , we have

$$\mathcal{E}_{\varphi}(A) = \mathcal{E}_{\varphi}(B)$$

▪

# Lemma: $\Sigma$ -Evaluations respect $\beta$ -Equality



Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and  $\mathbf{A} =_{\beta} \mathbf{B}$ . For all assignments  $\varphi$  into  $(\mathcal{D}, @)$ , we have

$$\mathcal{E}_{\varphi}(\mathbf{A}) = \mathcal{E}_{\varphi}(\mathbf{A} \downarrow_{\beta}) \quad \mathcal{E}_{\varphi}(\mathbf{B} \downarrow_{\beta}) = \mathcal{E}_{\varphi}(\mathbf{B})$$

▪

# Lemma: $\Sigma$ -Evaluations respect $\beta$ -Equality



Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and  $\mathbf{A} =_{\beta} \mathbf{B}$ . For all assignments  $\varphi$  into  $(\mathcal{D}, @)$ , we have

$$\mathcal{E}_{\varphi}(\mathbf{A}) = \mathcal{E}_{\varphi}(\mathbf{A} \downarrow_{\beta}) = \mathcal{E}_{\varphi}(\mathbf{B} \downarrow_{\beta}) = \mathcal{E}_{\varphi}(\mathbf{B})$$

▪

# Thm.: Substitution-Value Lemma



Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and  $\varphi$  be an assignment into  $\mathcal{J}$ .

# Thm.: Substitution-Value Lemma

Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and  $\varphi$  be an assignment into  $\mathcal{J}$ . For any types  $\alpha$  and  $\beta$ , variables  $X_\beta$ , and formulae  $A \in wff_\alpha(\Sigma)$  and  $B \in wff_\beta(\Sigma)$ , we have



# Thm.: Substitution-Value Lemma

Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and  $\varphi$  be an assignment into  $\mathcal{J}$ . For any types  $\alpha$  and  $\beta$ , variables  $X_\beta$ , and formulae  $\mathbf{A} \in wff_\alpha(\Sigma)$  and  $\mathbf{B} \in wff_\beta(\Sigma)$ , we have

$$\mathcal{E}_{\varphi, [\mathcal{E}_\varphi(\mathbf{B})/X]}(\mathbf{A}) = \mathcal{E}_\varphi([\mathbf{B}/X]\mathbf{A})$$

▪

# Prf.: Substitution-Value Lemma



Proof:

.

# Prf.: Substitution-Value Lemma



Proof: Using the fact that  $\mathcal{E}$  respects  $\beta$ -equality and the other properties of  $\mathcal{E}$ , we can compute

.

# Prf.: Substitution-Value Lemma



Proof: Using the fact that  $\mathcal{E}$  respects  $\beta$ -equality and the other properties of  $\mathcal{E}$ , we can compute

$$\mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}(\mathbf{A}) =$$

.

# Prf.: Substitution-Value Lemma



Proof: Using the fact that  $\mathcal{E}$  respects  $\beta$ -equality and the other properties of  $\mathcal{E}$ , we can compute

$$\mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}(\mathbf{A}) = \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}((\lambda \mathbf{X}. \mathbf{A})\mathbf{X})$$

.

# Prf.: Substitution-Value Lemma

Proof: Using the fact that  $\mathcal{E}$  respects  $\beta$ -equality and the other properties of  $\mathcal{E}$ , we can compute

$$\begin{aligned}\mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}(\mathbf{A}) &= \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}((\lambda \mathbf{X}. \mathbf{A})\mathbf{X}) \\ &= \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}(\lambda \mathbf{X}. \mathbf{A}) @ \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}(\mathbf{X})\end{aligned}$$

.

# Prf.: Substitution-Value Lemma

Proof: Using the fact that  $\mathcal{E}$  respects  $\beta$ -equality and the other properties of  $\mathcal{E}$ , we can compute

$$\begin{aligned}\mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}(\mathbf{A}) &= \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}((\lambda \mathbf{X}. \mathbf{A})\mathbf{X}) \\ &= \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}(\lambda \mathbf{X}. \mathbf{A}) @ \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}(\mathbf{X}) \\ &= \mathcal{E}_{\varphi}(\lambda \mathbf{X}. \mathbf{A}) @ \mathcal{E}_{\varphi}(\mathbf{B})\end{aligned}$$

.

# Prf.: Substitution-Value Lemma

Proof: Using the fact that  $\mathcal{E}$  respects  $\beta$ -equality and the other properties of  $\mathcal{E}$ , we can compute

$$\begin{aligned}
 \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}(\mathbf{A}) &= \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}((\lambda \mathbf{X}. \mathbf{A})\mathbf{X}) \\
 &= \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}(\lambda \mathbf{X}. \mathbf{A}) @ \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/\mathbf{X}]}(\mathbf{X}) \\
 &= \mathcal{E}_{\varphi}(\lambda \mathbf{X}. \mathbf{A}) @ \mathcal{E}_{\varphi}(\mathbf{B}) \\
 &= \mathcal{E}_{\varphi}((\lambda \mathbf{X}. \mathbf{A})\mathbf{B})
 \end{aligned}$$

.



# Prf.: Substitution-Value Lemma

Proof: Using the fact that  $\mathcal{E}$  respects  $\beta$ -equality and the other properties of  $\mathcal{E}$ , we can compute

$$\begin{aligned}
 \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/X]}(\mathbf{A}) &= \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/X]}((\lambda X. \mathbf{A})X) \\
 &= \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/X]}(\lambda X. \mathbf{A}) @ \mathcal{E}_{\varphi, [\mathcal{E}_{\varphi}(\mathbf{B})/X]}(X) \\
 &= \mathcal{E}_{\varphi}(\lambda X. \mathbf{A}) @ \mathcal{E}_{\varphi}(\mathbf{B}) \\
 &= \mathcal{E}_{\varphi}((\lambda X. \mathbf{A})\mathbf{B}) \\
 &= \mathcal{E}_{\varphi}([\mathbf{B}/X]\mathbf{A}).
 \end{aligned}$$

# Weaker Notions of Functionality



We will consider two weaker notions of functionality. These forms are often discussed in the literature (cf. [HindleySeldin86]).

# Weaker Notions of Functionality



We will consider two weaker notions of functionality. These forms are often discussed in the literature (cf. [HindleySeldin86]).

- $\eta$ -functionality simply means the evaluation respects  $\eta$ -conversion.

# Weaker Notions of Functionality



We will consider two weaker notions of functionality. These forms are often discussed in the literature (cf. [HindleySeldin86]).

- $\eta$ -functionality simply means the evaluation respects  $\eta$ -conversion.
- $\xi$ -functionality means we have functionality (only) with respect to  $\lambda$ -abstractions.

# Def.: $\eta$ -Functional

---



Let  $\mathcal{J} = (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation.

# Def.: $\eta$ -Functional

---



Let  $\mathcal{J} = (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation.

We say  $\mathcal{J}$  is  $\eta$ -functional if

# Def.: $\eta$ -Functional

Let  $\mathcal{J} = (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation.

We say  $\mathcal{J}$  is  $\eta$ -functional if

$$\mathcal{E}_\varphi(\mathbf{A}) = \mathcal{E}_\varphi(\mathbf{A} \downarrow_{\beta\eta})$$

for any type  $\alpha$ , formula  $\mathbf{A} \in wff_\alpha(\Sigma)$ , and assignment  $\varphi$ .

# Def.: $\xi$ -Functional

---



Let  $\mathcal{J} = (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation.



# Def.: $\xi$ -Functional

---



Let  $\mathcal{J} = (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation. We say  $\mathcal{J}$  is  $\xi$ -functional if

# Def.: $\xi$ -Functional



Let  $\mathcal{J} = (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation. We say  $\mathcal{J}$  is  $\xi$ -functional if for all  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{M}, \mathbf{N} \in wff_{\beta}(\Sigma)$ , assignments  $\varphi$ , and variables  $X_{\alpha}$ ,

# Def.: $\xi$ -Functional

Let  $\mathcal{J} = (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation. We say  $\mathcal{J}$  is  $\xi$ -functional if for all  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{M}, \mathbf{N} \in wff_\beta(\Sigma)$ , assignments  $\varphi$ , and variables  $X_\alpha$ ,

$$\mathcal{E}_\varphi(\lambda X_\alpha. \mathbf{M}_\beta) = \mathcal{E}_\varphi(\lambda X_\alpha. \mathbf{N}_\beta)$$

# Def.: $\xi$ -Functional

Let  $\mathcal{J} = (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation. We say  $\mathcal{J}$  is  $\xi$ -functional if for all  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{M}, \mathbf{N} \in wff_\beta(\Sigma)$ , assignments  $\varphi$ , and variables  $X_\alpha$ ,

$$\mathcal{E}_\varphi(\lambda X_\alpha. \mathbf{M}_\beta) = \mathcal{E}_\varphi(\lambda X_\alpha. \mathbf{N}_\beta)$$

whenever

$$\mathcal{E}_{\varphi, [a/X]}(\mathbf{M}) = \mathcal{E}_{\varphi, [a/X]}(\mathbf{N})$$

for every  $a \in \mathcal{D}_\alpha$ .

# Lemma: Functionality and $\eta$



Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a functional  $\Sigma$ -evaluation.

# Lemma: Functionality and $\eta$

Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a functional  $\Sigma$ -evaluation.

1. For any assignment  $\varphi$  into  $\mathcal{J}$  and  $\mathbf{F} \in wff_{\alpha \rightarrow \beta}(\Sigma)$  where  $X_\alpha \notin \mathbf{Free}(\mathbf{F})$ , we have

# Lemma: Functionality and $\eta$

Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a functional  $\Sigma$ -evaluation.

1. For any assignment  $\varphi$  into  $\mathcal{J}$  and  $\mathbf{F} \in wff_{\alpha \rightarrow \beta}(\Sigma)$  where  $X_\alpha \notin \mathbf{Free}(\mathbf{F})$ , we have

$$\mathcal{E}_\varphi(\lambda X_\alpha. \mathbf{F} X) = \mathcal{E}_\varphi(\mathbf{F})$$

# Lemma: Functionality and $\eta$

Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a functional  $\Sigma$ -evaluation.

1. For any assignment  $\varphi$  into  $\mathcal{J}$  and  $\mathbf{F} \in wff_{\alpha \rightarrow \beta}(\Sigma)$  where  $X_\alpha \notin \mathbf{Free}(\mathbf{F})$ , we have

$$\mathcal{E}_\varphi(\lambda X_\alpha. \mathbf{F} X) = \mathcal{E}_\varphi(\mathbf{F})$$

2. If a formula  $\mathbf{A}$   $\eta$ -reduces to  $\mathbf{B}$  in one step, then for any assignment  $\varphi$  into  $\mathcal{J}$ , we have



# Lemma: Functionality and $\eta$

Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a functional  $\Sigma$ -evaluation.

1. For any assignment  $\varphi$  into  $\mathcal{J}$  and  $\mathbf{F} \in wff_{\alpha \rightarrow \beta}(\Sigma)$  where  $X_\alpha \notin \mathbf{Free}(\mathbf{F})$ , we have

$$\mathcal{E}_\varphi(\lambda X_\alpha. \mathbf{F} X) = \mathcal{E}_\varphi(\mathbf{F})$$

2. If a formula  $\mathbf{A}$   $\eta$ -reduces to  $\mathbf{B}$  in one step, then for any assignment  $\varphi$  into  $\mathcal{J}$ , we have

$$\mathcal{E}_\varphi(\mathbf{A}) = \mathcal{E}_\varphi(\mathbf{B})$$

# Lemma: Functionality and $\eta$

Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a functional  $\Sigma$ -evaluation.

1. For any assignment  $\varphi$  into  $\mathcal{J}$  and  $\mathbf{F} \in wff_{\alpha \rightarrow \beta}(\Sigma)$  where  $X_\alpha \notin \mathbf{Free}(\mathbf{F})$ , we have

$$\mathcal{E}_\varphi(\lambda X_\alpha. \mathbf{F} X) = \mathcal{E}_\varphi(\mathbf{F})$$

2. If a formula  $\mathbf{A}$   $\eta$ -reduces to  $\mathbf{B}$  in one step, then for any assignment  $\varphi$  into  $\mathcal{J}$ , we have

$$\mathcal{E}_\varphi(\mathbf{A}) = \mathcal{E}_\varphi(\mathbf{B})$$

Proof: Exercise

# Lemma: Functionality and $\eta+\xi$



Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation.

# Lemma: Functionality and $\eta+\xi$



Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation. Then  $\mathcal{J}$  is functional iff it is both  $\eta$ -functional and  $\xi$ -functional.

# Lemma: Functionality and $\eta+\xi$



Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation. Then  $\mathcal{J}$  is functional iff it is both  $\eta$ -functional and  $\xi$ -functional.

Proof: Exercise

# Logical Constants in Signature



Let  $\Sigma := (\mathcal{V}, \mathcal{C})$  be a signature.

# Logical Constants in Signature



Let  $\Sigma := (\mathcal{V}, \mathcal{C})$  be a signature.

The following **logical constants may** or **may not** be in the set  $\mathcal{C}$  of constants:

# Logical Constants in Signature

Let  $\Sigma := (\mathcal{V}, \mathcal{C})$  be a signature.

The following **logical constants may or may not** be in the set  $\mathcal{C}$  of constants:

$$\top_0, \perp_0, \neg_{00}, \vee_{000}, \wedge_{000}, \supset_{000}, \Leftrightarrow_{000}$$



# Logical Constants in Signature

Let  $\Sigma := (\mathcal{V}, \mathcal{C})$  be a signature.

The following **logical constants may or may not** be in the set  $\mathcal{C}$  of constants:

$$\top_o, \perp_o, \neg_{oo}, \vee_{ooo}, \wedge_{ooo}, \supset_{ooo}, \Leftrightarrow_{ooo}$$

$$\prod_{o(o\alpha)}^\alpha (\prod_{o\alpha}^\alpha F_{o\alpha} \sim \forall x_\alpha Fx), \quad \sum_{o(o\alpha)}^\alpha (\sum_{o\alpha}^\alpha F_{o\alpha} \sim \exists x_\alpha Fx)$$

# Logical Constants in Signature

Let  $\Sigma := (\mathcal{V}, \mathcal{C})$  be a signature.

The following **logical constants may** or **may not** be in the set  $\mathcal{C}$  of constants:

$$\top_o, \perp_o, \neg_{oo}, \vee_{ooo}, \wedge_{ooo}, \supset_{ooo}, \Leftrightarrow_{ooo}$$

$$\prod_{o(o\alpha)}^\alpha (\prod_{o\alpha}^\alpha F_{o\alpha} \sim \forall x_\alpha Fx), \quad \sum_{o(o\alpha)}^\alpha (\sum_{o\alpha}^\alpha F_{o\alpha} \sim \exists x_\alpha Fx)$$

$$=_{o\alpha\alpha}^\alpha$$

# Logical Constants in Signature

Let  $\Sigma := (\mathcal{V}, \mathcal{C})$  be a signature.

The following **logical constants may or may not** be in the set  $\mathcal{C}$  of constants:

$$\top_o, \perp_o, \neg_{oo}, \vee_{ooo}, \wedge_{ooo}, \supset_{ooo}, \Leftrightarrow_{ooo}$$

$$\prod_{o(o\alpha)}^\alpha (\prod_{o\alpha}^\alpha F_{o\alpha} \sim \forall x_\alpha Fx), \quad \sum_{o(o\alpha)}^\alpha (\sum_{o\alpha}^\alpha F_{o\alpha} \sim \exists x_\alpha Fx)$$

$$=_{o\alpha\alpha}^\alpha$$

for all  $\alpha \in \mathcal{T}$

# Once More: Cantor's Theorem



For any set  $A$ ,

$$|A| < |\mathcal{P}(A)|$$

# Once More: Cantor's Theorem



For any set  $A$ ,

$$|A| < |\mathcal{P}(A)|$$

i.e.,  $\neg \exists g : A \rightarrow \mathcal{P}(A)$  with  $g$  surjective.

# Once More: Cantor's Theorem



Assume the set  $A$  is associated with  $\iota$ .

# Once More: Cantor's Theorem



Assume the set  $A$  is associated with  $\iota$ . Then  $\mathcal{P}(A)$  has type  $o_\iota$ , i.e. the type of "sets" (or **characteristic functions**) over  $\iota$ .

# Once More: Cantor's Theorem



Assume the set  $A$  is associated with  $\iota$ . Then  $\mathcal{P}(A)$  has type  $o_\iota$ , i.e. the type of "sets" (or **characteristic functions**) over  $\iota$ .

$D_{o_\iota}$



# Once More: Cantor's Theorem



Assume the set  $A$  is associated with  $\iota$ . Then  $\mathcal{P}(A)$  has type  $o\iota$ , i.e. the type of "sets" (or **characteristic functions**) over  $\iota$ .

$$D_{o\iota} = D_o^{D_\iota}$$

# Once More: Cantor's Theorem



Assume the set  $A$  is associated with  $\iota$ . Then  $\mathcal{P}(A)$  has type  $o_\iota$ , i.e. the type of "sets" (or **characteristic functions**) over  $\iota$ .

$$\begin{aligned} D_{o_\iota} &= D_o^{D_\iota} \\ &= \{\perp, \top\}^{D_\iota} \end{aligned}$$

# Once More: Cantor's Theorem



Assume the set  $A$  is associated with  $\iota$ . Then  $\mathcal{P}(A)$  has type  $o_\iota$ , i.e. the type of "sets" (or **characteristic functions**) over  $\iota$ .

$$\begin{aligned} D_{o_\iota} &= D_o^{D_\iota} \\ &= \{\perp, \top\}^{D_\iota} \\ &= \{f \mid f : D_\iota \rightarrow \{\perp, \top\}\} \end{aligned}$$

# Once More: Cantor's Theorem



Assume the set  $A$  is associated with  $\iota$ . Then  $\mathcal{P}(A)$  has type  $o_\iota$ , i.e. the type of "sets" (or **characteristic functions**) over  $\iota$ .

$$\begin{aligned} D_{o_\iota} &= D_o^{D_\iota} \\ &= \{\perp, \top\}^{D_\iota} \\ &= \{f \mid f : D_\iota \rightarrow \{\perp, \top\}\} \\ &\cong \{X \mid X \subseteq D_\iota\} \end{aligned}$$

# Once More: Cantor's Theorem



Assume the set  $A$  is associated with  $\iota$ . Then  $\mathcal{P}(A)$  has type  $o_\iota$ , i.e. the type of "sets" (or **characteristic functions**) over  $\iota$ .

$$\begin{aligned} D_{o_\iota} &= D_o^{D_\iota} \\ &= \{\perp, \top\}^{D_\iota} \\ &= \{f \mid f : D_\iota \rightarrow \{\perp, \top\}\} \\ &\cong \{X \mid X \subseteq D_\iota\} \\ &= \mathcal{P}(D_\iota) \end{aligned}$$

# Once More: Cantor's Theorem



We can now formulate Cantor's Theorem with typed terms (as seen before):

# Once More: Cantor's Theorem



We can now formulate Cantor's Theorem with typed terms (as seen before):

$$\neg \exists g_{o \iota \iota} \forall f_{o \iota} \exists x_{\iota} : gx = f$$

# Once More: Cantor's Theorem



We can now formulate Cantor's Theorem with typed terms (as seen before):

$$\neg \exists g_{o \iota \iota} \forall f_{o \iota} \exists x_{\iota} : gx = f$$

which is shorthand for:



# Once More: Cantor's Theorem



We can now formulate Cantor's Theorem with typed terms (as seen before):

$$\neg \exists g_{o\iota\iota} \forall f_{o\iota} \exists x_{\iota} : gx = f$$

which is shorthand for:

$$\neg_{oo} \Sigma_{o(o(o\iota\iota))}^{o\iota\iota} \left( \lambda g_{o\iota\iota} . \Pi_{o(o(o\iota))}^{o\iota} \left( \lambda f_{o\iota} . \Sigma_{o(o\iota)}^{\iota} \left( \lambda x_{\iota} . =_{o(o\iota)(o\iota)}^{o\iota} (gx) f \right) \right) \right)$$

# Once More: Cantor's Theorem

We can now formulate Cantor's Theorem with typed terms (as seen before):

$$\neg \exists g_{o\iota\iota} \forall f_{o\iota} \exists x_{\iota} : gx = f$$

which is shorthand for:

$$\neg_{oo} \Sigma_{o(o(o\iota\iota))}^{o\iota\iota} \left( \lambda g_{o\iota\iota} . \Pi_{o(o(o\iota))}^{o\iota} \left( \lambda f_{o\iota} . \Sigma_{o(o\iota)}^{\iota} \left( \lambda x_{\iota} . =_{o(o\iota)(o\iota)}^{o\iota} (gx) f \right) \right) \right)$$

**Note:** for this term to be in the set  $cwff_{\alpha}(\Sigma)$ , the constants  $\neg_{oo}$ ,  $\Sigma_{o(o(o\iota\iota))}^{o\iota\iota}$ ,  $\Pi_{o(o(o\iota))}^{o\iota}$ ,  $\Sigma^{\iota}$  and  $=^{o\iota}$  have to be in the set  $\mathcal{C}$ .

# Once More: Cantor's Theorem



Proof:

# Once More: Cantor's Theorem



Proof: Assume such a function  $g$  exists.

# Once More: Cantor's Theorem



Proof: Assume such a function  $g$  exists.

Let  $f = \{x \mid x \notin gx\}$  that is  $f = (\lambda x. \neg gxx)$ .

# Once More: Cantor's Theorem



Proof: Assume such a function  $g$  exists.

Let  $f = \{x \mid x \notin gx\}$  that is  $f = (\lambda x. \neg gxx)$ .

$g$  is surjective,

# Once More: Cantor's Theorem

Proof: Assume such a function  $g$  exists.

Let  $f = \{x \mid x \notin gx\}$  that is  $f = (\lambda x. \neg gx)$ .

$g$  is surjective, hence

$$(\exists y. gy = (\lambda x. \neg gx))$$

# Once More: Cantor's Theorem



Proof: Assume such a function  $g$  exists.

Let  $f = \{x \mid x \notin gx\}$  that is  $f = (\lambda x. \neg gxx)$ .

$g$  is surjective, hence

$$(\exists y. : gy = [\lambda x. \neg gxx])$$

hence

$$(gyy \Leftrightarrow \neg gyy)$$



# Once More: Cantor's Theorem



Proof: Assume such a function  $g$  exists.

Let  $f = \{x \mid x \notin gx\}$  that is  $f = (\lambda x. \neg gxx)$ .

$g$  is surjective, hence

$$(\exists y. : gy = [\lambda x. \neg gxx])$$

hence

$$(gyy \Leftrightarrow \neg gyy)$$

Contradiction!

# Once More: Cantor's Theorem

Proof: Assume such a function  $g$  exists.

Let  $f = \{x \mid x \notin gx\}$  that is  $f = (\lambda x. \neg gx)$ .

$g$  is surjective, hence

$$(\exists y. : gy = [ \lambda x. \neg gx ])$$

hence

$$(gyy \Leftrightarrow \neg gyy)$$

Contradiction!

Note that the proof uses  $\neg$ .



Semantics:  $\Sigma$ -Models

# Def.: Properties of Logical Constants



Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ).

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_0 \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c		

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c	$\beta$	

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c	$\beta$	$\mathcal{L}_c(( )a)$ holds when



# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c	$\beta$	$\mathcal{L}_c(( )a)$ holds when
T		

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c	$\beta$	$\mathcal{L}_c(( )a)$ holds when
T	o	

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c	$\beta$	$\mathcal{L}_c(( )a)$ holds when
T	o	$v(a) = T$

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$		

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$		

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_0 \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$0$	$v(a) = T$
$\perp$	$0$	$v(a) = F$
$\neg$	$00$	



# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_0 \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_0$
$\vee$	$ooo$	

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$		

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

c	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_0 \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_0 \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$		



# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_0 \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$	$ooo$	

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$	$ooo$	$v(a@b@c) = T$ iff $v(b) = F$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$	$ooo$	$v(a@b@c) = T$ iff $v(b) = F$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\Leftrightarrow$		

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$	$ooo$	$v(a@b@c) = T$ iff $v(b) = F$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\Leftrightarrow$	$ooo$	

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$	$ooo$	$v(a@b@c) = T$ iff $v(b) = F$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\Leftrightarrow$	$ooo$	$v(a@b@c) = T$ iff $v(b) = v(c) \ \forall b, c \in \mathcal{D}_o$

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$	$ooo$	$v(a@b@c) = T$ iff $v(b) = F$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\Leftrightarrow$	$ooo$	$v(a@b@c) = T$ iff $v(b) = v(c) \ \forall b, c \in \mathcal{D}_o$
$=^\alpha$		

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$	$ooo$	$v(a@b@c) = T$ iff $v(b) = F$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\Leftrightarrow$	$ooo$	$v(a@b@c) = T$ iff $v(b) = v(c) \ \forall b, c \in \mathcal{D}_o$
$=^\alpha$	$o\alpha\alpha$	

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$	$ooo$	$v(a@b@c) = T$ iff $v(b) = F$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\Leftrightarrow$	$ooo$	$v(a@b@c) = T$ iff $v(b) = v(c) \ \forall b, c \in \mathcal{D}_o$
$=^\alpha$	$o\alpha\alpha$	$v(a@b@c) = T$ iff $b = c \ \forall b, c \in \mathcal{D}_o$



# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$	$ooo$	$v(a@b@c) = T$ iff $v(b) = F$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\Leftrightarrow$	$ooo$	$v(a@b@c) = T$ iff $v(b) = v(c) \ \forall b, c \in \mathcal{D}_o$
$=^\alpha$	$o\alpha\alpha$	$v(a@b@c) = T$ iff $b = c \ \forall b, c \in \mathcal{D}_o$
$\prod^\alpha$		

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$	$ooo$	$v(a@b@c) = T$ iff $v(b) = F$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\Leftrightarrow$	$ooo$	$v(a@b@c) = T$ iff $v(b) = v(c) \ \forall b, c \in \mathcal{D}_o$
$=^\alpha$	$o\alpha\alpha$	$v(a@b@c) = T$ iff $b = c \ \forall b, c \in \mathcal{D}_o$
$\Pi^\alpha$	$o(o\alpha)$	

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$	$ooo$	$v(a@b@c) = T$ iff $v(b) = F$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\Leftrightarrow$	$ooo$	$v(a@b@c) = T$ iff $v(b) = v(c) \ \forall b, c \in \mathcal{D}_o$
$=^\alpha$	$o\alpha\alpha$	$v(a@b@c) = T$ iff $b = c \ \forall b, c \in \mathcal{D}_o$
$\prod^\alpha$	$o(o\alpha)$	$v(a@f) = T$ iff $\forall b \in \mathcal{D}_\alpha : v(f@b) = T \ \forall f \in \mathcal{D}_{o\alpha}$

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$	$ooo$	$v(a@b@c) = T$ iff $v(b) = F$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\Leftrightarrow$	$ooo$	$v(a@b@c) = T$ iff $v(b) = v(c) \ \forall b, c \in \mathcal{D}_o$
$=^\alpha$	$o\alpha\alpha$	$v(a@b@c) = T$ iff $b = c \ \forall b, c \in \mathcal{D}_o$
$\prod^\alpha$	$o(o\alpha)$	$v(a@f) = T$ iff $\forall b \in \mathcal{D}_\alpha : v(f@b) = T \ \forall f \in \mathcal{D}_{o\alpha}$
$\sum^\alpha$		

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$	$ooo$	$v(a@b@c) = T$ iff $v(b) = F$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\Leftrightarrow$	$ooo$	$v(a@b@c) = T$ iff $v(b) = v(c) \ \forall b, c \in \mathcal{D}_o$
$=^\alpha$	$o\alpha\alpha$	$v(a@b@c) = T$ iff $b = c \ \forall b, c \in \mathcal{D}_o$
$\Pi^\alpha$	$o(o\alpha)$	$v(a@f) = T$ iff $\forall b \in \mathcal{D}_\alpha : v(f@b) = T \ \forall f \in \mathcal{D}_{o\alpha}$
$\Sigma^\alpha$	$o(o\alpha)$	

# Def.: Properties of Logical Constants

Let  $(D, @)$  be an applicative structure and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a function (for given  $T \neq F$ ). For each logical constant  $c_\beta$  and for  $a \in \mathcal{D}_\beta$ , we define the proposition  $\mathcal{L}_c(( )a)$  with respect to  $v$ :

$c$	$\beta$	$\mathcal{L}_c(( )a)$ holds when
$\top$	$o$	$v(a) = T$
$\perp$	$o$	$v(a) = F$
$\neg$	$oo$	$v(a@b) = T$ iff $v(b) = F \ \forall b \in \mathcal{D}_o$
$\vee$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\wedge$	$ooo$	$v(a@b@c) = T$ iff $v(b) = T$ and $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\supset$	$ooo$	$v(a@b@c) = T$ iff $v(b) = F$ or $v(c) = T \ \forall b, c \in \mathcal{D}_o$
$\Leftrightarrow$	$ooo$	$v(a@b@c) = T$ iff $v(b) = v(c) \ \forall b, c \in \mathcal{D}_o$
$=^\alpha$	$o\alpha\alpha$	$v(a@b@c) = T$ iff $b = c \ \forall b, c \in \mathcal{D}_o$
$\Pi^\alpha$	$o(o\alpha)$	$v(a@f) = T$ iff $\forall b \in \mathcal{D}_\alpha : v(f@b) = T \ \forall f \in \mathcal{D}_{o\alpha}$
$\Sigma^\alpha$	$o(o\alpha)$	$v(a@f) = T$ iff $\exists b \in \mathcal{D}_\alpha : v(f@b) = T \ \forall f \in \mathcal{D}_{o\alpha}$

# Def.: $\Sigma$ -Valuation



Let  $\mathcal{J} := (D, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and  $v : \mathcal{D}_o \rightarrow \{T, F\}$ .

# Def.: $\Sigma$ -Valuation

Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and  $v : \mathcal{D}_o \rightarrow \{\mathbf{T}, \mathbf{F}\}$ . We say  $v$  is a  $\Sigma$ -valuation w.r.t  $\mathcal{J}$  if



# Def.: $\Sigma$ -Valuation

Let  $\mathcal{J} := (\mathcal{D}, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and  $v : \mathcal{D}_o \rightarrow \{\text{T}, \text{F}\}$ . We say  $v$  is a  $\Sigma$ -valuation w.r.t  $\mathcal{J}$  if  $\mathcal{L}_c((\mathcal{E}(c)))$  holds w.r.t  $v$  for each logical constant  $c \in \Sigma$ .

# Def.: $\Sigma$ -Model



Let  $\mathcal{J} := (D, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a  $\Sigma$ -valuation w.r.t  $\mathcal{J}$

# Def.: $\Sigma$ -Model

Let  $\mathcal{J} := (D, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a  $\Sigma$ -valuation w.r.t  $\mathcal{J}$

We say  $\mathcal{M} = (D, @, \mathcal{E}, v)$  is a  $\Sigma$ -model.

# Def.: $\Sigma$ -Model

Let  $\mathcal{J} := (D, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a  $\Sigma$ -valuation w.r.t  $\mathcal{J}$

We say  $\mathcal{M} = (D, @, \mathcal{E}, v)$  is a  $\Sigma$ -model.

If  $(D, @, \mathcal{E})$  is functional (full, standard), we say  $\mathcal{M}$  is functional (full, standard).

# Def.: $\Sigma$ -Model

Let  $\mathcal{J} := (D, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a  $\Sigma$ -valuation w.r.t  $\mathcal{J}$

We say  $\mathcal{M} = (D, @, \mathcal{E}, v)$  is a  $\Sigma$ -model.

If  $(D, @, \mathcal{E})$  is functional (full, standard), we say  $\mathcal{M}$  is functional (full, standard).

If  $(D, @, \mathcal{E})$  is  $\eta$ -functional, we say  $\mathcal{M}$  is  $\eta$ -functional.

# Def.: $\Sigma$ -Model

Let  $\mathcal{J} := (D, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a  $\Sigma$ -valuation w.r.t  $\mathcal{J}$

We say  $\mathcal{M} = (D, @, \mathcal{E}, v)$  is a  $\Sigma$ -model.

If  $(D, @, \mathcal{E})$  is functional (full, standard), we say  $\mathcal{M}$  is functional (full, standard).

If  $(D, @, \mathcal{E})$  is  $\eta$ -functional, we say  $\mathcal{M}$  is  $\eta$ -functional.

If  $(D, @, \mathcal{E})$  is  $\xi$ -functional, we say  $\mathcal{M}$  is  $\xi$ -functional.

# Some Conventions: Equality



Some important conventions:

- $=$  denotes **primitive equality**

# Some Conventions: Equality



Some important conventions:

- $=$  denotes **primitive equality**
- $\doteq$  denotes **Leibniz equality**:  $A_\alpha \doteq^\alpha B_\alpha := \forall P_{o\alpha}. (PA) \Rightarrow (PB)$



# Some Conventions: Equality



Some important conventions:

- $=$  denotes **primitive equality**
- $\doteq$  denotes **Leibniz equality**:  $A_\alpha \doteq^\alpha B_\alpha := \forall P_{o\alpha}. (PA) \Rightarrow (PB)$
- $\ddot{=}$  ... other definition of equality (e.g., see [Andrews02])

# Some Conventions: Equality



Some important conventions:

- $=$  denotes **primitive equality**
- $\doteq$  denotes **Leibniz equality**:  $A_\alpha \doteq^\alpha B_\alpha := \forall P_{o\alpha}. (PA) \Rightarrow (PB)$
- $\ddot{=}$  ... other definition of equality (e.g., see [Andrews02])

We use  $\stackrel{*}{=}$  in the following to refer to **any** of the above

# Def.: Properties $f, b, \eta, \xi$



Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model. We say,  $\mathbf{M}$  has property

# Def.: Properties $f, b, \eta, \xi$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model. We say,  $M$  has property  $\eta$  if  $M$  is  $\eta$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\eta$ -functional)

# Def.: Properties $f, b, \eta, \xi$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model. We say,  $M$  has property

$\eta$  if  $M$  is  $\eta$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\eta$ -functional)

$\xi$  if  $M$  is  $\xi$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\xi$ -functional)

# Def.: Properties $f, b, \eta, \xi$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model. We say,  $M$  has property

$\eta$  if  $M$  is  $\eta$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\eta$ -functional)

$\xi$  if  $M$  is  $\xi$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\xi$ -functional)

$f$  if  $M$  is functional (respectively  $(D, @, \mathcal{E})$  is functional)

# Def.: Properties $f, b, \eta, \xi$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model. We say,  $M$  has property

$\eta$  if  $M$  is  $\eta$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\eta$ -functional)

$\xi$  if  $M$  is  $\xi$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\xi$ -functional)

$f$  if  $M$  is functional (respectively  $(D, @, \mathcal{E})$  is functional)

$b$  if  $v$  is injective.

# Def.: Properties $f, b, \eta, \xi$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model. We say,  $M$  has property

$\eta$  if  $M$  is  $\eta$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\eta$ -functional)

$\xi$  if  $M$  is  $\xi$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\xi$ -functional)

$f$  if  $M$  is functional (respectively  $(D, @, \mathcal{E})$  is functional)

$b$  if  $v$  is injective.

Note: In the [JSC04]-paper,  $b$  is defined as  $D_o = \{T, F\}$ , but here we are using the injectivity criterion, because we are varying the signature. If the signature is too sparse, we could have a  $D_o$  with two elements which both valuate via  $v$  to  $T$ . Another ill case would be  $D_o$  with just one element.



# Def.: Properties $f, b, \eta, \xi$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model. We say,  $M$  has property

$\eta$  if  $M$  is  $\eta$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\eta$ -functional)

$\xi$  if  $M$  is  $\xi$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\xi$ -functional)

$f$  if  $M$  is functional (respectively  $(D, @, \mathcal{E})$  is functional)

$b$  if  $v$  is injective.

$q$  if for all  $\alpha \in \mathcal{T}$  there is some  $q \in D_{o\alpha\alpha}$  such that  $\mathcal{L}_{=\alpha}(q)$ .

# Def.: Properties $f, b, \eta, \xi$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model. We say,  $M$  has property

$\eta$  if  $M$  is  $\eta$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\eta$ -functional)

$\xi$  if  $M$  is  $\xi$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\xi$ -functional)

$f$  if  $M$  is functional (respectively  $(D, @, \mathcal{E})$  is functional)

$b$  if  $v$  is injective.

$q$  if for all  $\alpha \in \mathcal{T}$  there is some  $q \in D_{o\alpha\alpha}$  such that  $\mathcal{L}_{=\alpha}(q)$ .

Note: This basically says that for each type  $\alpha$  the identity relation over  $\alpha$  is already present in the model. If we require  $=_{o\alpha\alpha} \in \mathcal{C}$  with  $\mathcal{L}_{=\alpha}(\mathcal{E}_{\varphi}(=_{o\alpha\alpha}))$ , then this property is automatically ensured, but not for weaker signatures. See [Andrew71] for a detailed discussion of property  $q$ . Andrews constructs a Henkin model where Leibniz equality  $\doteq$  does not evaluate to the intended identity relation. This is resolved by property  $q$ .

# Lemma: Surjective $v$



Let  $\mathcal{C}$  be a signature and  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model.

# Lemma: Surjective $v$

Let  $\mathcal{C}$  be a signature and  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model.  
 If  $T, F \in \mathcal{C}$  or  $\neg \in \mathcal{C}$  then  $v$  is surjective.

# Lemma: Surjective $v$

Let  $\mathcal{C}$  be a signature and  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model.  
 If  $T, F \in \mathcal{C}$  or  $\neg \in \mathcal{C}$  then  $v$  is surjective.

Proof: Exercise.

# Thm.: Property $\mathfrak{b}$



Let  $\mathcal{C}$  be a signature and  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model.

# Thm.: Property $\mathfrak{b}$



Let  $\mathcal{C}$  be a signature and  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model.  
Suppose  $\mathsf{T}, \mathsf{F} \in \mathcal{C}$  or  $\neg \in \mathcal{C}$ .

# Thm.: Property $\mathfrak{b}$

Let  $\mathcal{C}$  be a signature and  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model.

Suppose  $\mathsf{T}, \mathsf{F} \in \mathcal{C}$  or  $\neg \in \mathcal{C}$ .

Then  $\mathcal{M}$  satisfies property  $\mathfrak{b}$  iff  $|D_o| = 2$ .



# Thm.: Property $\mathfrak{b}$

Let  $\mathcal{C}$  be a signature and  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model.

Suppose  $\mathsf{T}, \mathsf{F} \in \mathcal{C}$  or  $\neg \in \mathcal{C}$ .

Then  $\mathcal{M}$  satisfies property  $\mathfrak{b}$  iff  $|D_o| = 2$ .

Proof: Exercise.



Semantics: HOL-CUBE

# Def. (Reminder): $\Sigma$ -Model

Let  $\mathcal{J} := (D, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a  $\Sigma$ -valuation w.r.t  $\mathcal{J}$

# Def. (Reminder): $\Sigma$ -Model

Let  $\mathcal{J} := (D, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a  $\Sigma$ -valuation w.r.t  $\mathcal{J}$

We say  $\mathcal{M} = (D, @, \mathcal{E}, v)$  is a  $\Sigma$ -model.

# Def. (Reminder): $\Sigma$ -Model

Let  $\mathcal{J} := (D, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a  $\Sigma$ -valuation w.r.t  $\mathcal{J}$

We say  $\mathcal{M} = (D, @, \mathcal{E}, v)$  is a  $\Sigma$ -model.

If  $(D, @, \mathcal{E})$  is functional (full, standard), we say  $\mathcal{M}$  is functional (full, standard).

# Def. (Reminder): $\Sigma$ -Model

Let  $\mathcal{J} := (D, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a  $\Sigma$ -valuation w.r.t  $\mathcal{J}$

We say  $\mathcal{M} = (D, @, \mathcal{E}, v)$  is a  $\Sigma$ -model.

If  $(D, @, \mathcal{E})$  is functional (full, standard), we say  $\mathcal{M}$  is functional (full, standard).

If  $(D, @, \mathcal{E})$  is  $\eta$ -functional, we say  $\mathcal{M}$  is  $\eta$ -functional.

# Def. (Reminder): $\Sigma$ -Model

Let  $\mathcal{J} := (D, @, \mathcal{E})$  be a  $\Sigma$ -evaluation and let  $v : \mathcal{D}_o \rightarrow \{T, F\}$  be a  $\Sigma$ -valuation w.r.t  $\mathcal{J}$

We say  $\mathcal{M} = (D, @, \mathcal{E}, v)$  is a  $\Sigma$ -model.

If  $(D, @, \mathcal{E})$  is functional (full, standard), we say  $\mathcal{M}$  is functional (full, standard).

If  $(D, @, \mathcal{E})$  is  $\eta$ -functional, we say  $\mathcal{M}$  is  $\eta$ -functional.

If  $(D, @, \mathcal{E})$  is  $\xi$ -functional, we say  $\mathcal{M}$  is  $\xi$ -functional.

# Def. (Reminder): Properties $f, b, \eta, \xi$



Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model. We say,  $\mathcal{M}$  has property



# Def. (Reminder): Properties $f, b, \eta, \xi$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model. We say,  $M$  has property  $\eta$  if  $M$  is  $\eta$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\eta$ -functional)

# Def. (Reminder): Properties $f$ , $b$ , $\eta$ , $\xi$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model. We say,  $M$  has property

$\eta$  if  $M$  is  $\eta$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\eta$ -functional)

$\xi$  if  $M$  is  $\xi$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\xi$ -functional)

# Def. (Reminder): Properties $f, b, \eta, \xi$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model. We say,  $M$  has property

$\eta$  if  $M$  is  $\eta$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\eta$ -functional)

$\xi$  if  $M$  is  $\xi$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\xi$ -functional)

$f$  if  $M$  is functional (respectively  $(D, @, \mathcal{E})$  is functional)

# Def. (Reminder): Properties $f, b, \eta, \xi$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model. We say,  $M$  has property

$\eta$  if  $M$  is  $\eta$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\eta$ -functional)

$\xi$  if  $M$  is  $\xi$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\xi$ -functional)

$f$  if  $M$  is functional (respectively  $(D, @, \mathcal{E})$  is functional)

$b$  if  $v$  is injective.

# Def. (Reminder): Properties $f, b, \eta, \xi$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\mathcal{C}$ -model. We say,  $M$  has property

$\eta$  if  $M$  is  $\eta$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\eta$ -functional)

$\xi$  if  $M$  is  $\xi$ -functional (respectively  $(D, @, \mathcal{E})$  is  $\xi$ -functional)

$f$  if  $M$  is functional (respectively  $(D, @, \mathcal{E})$  is functional)

$b$  if  $v$  is injective.

$q$  if for all  $\alpha \in \mathcal{T}$  there is some  $q \in D_{o\alpha\alpha}$  such that  $\mathcal{L}_{=\alpha}(q)$ .

# Def. (Reminder): Different Model Classes



We denote the class of  $\mathcal{C}$ -models by  $\mathfrak{M}_\beta(\Sigma)$ .

# Def. (Reminder): Different Model Classes



We denote the class of  $\mathcal{C}$ -models by  $\mathfrak{M}_\beta(\Sigma)$ . We obtain a hierarchy of subclasses of  $\mathfrak{M}_\beta(\Sigma)$  by adding the properties  $\xi, \eta, f, b$ .

# Def. (Reminder): Different Model Classes



We denote the class of  $\mathcal{C}$ -models by  $\mathfrak{M}_\beta(\Sigma)$ . We obtain a hierarchy of subclasses of  $\mathfrak{M}_\beta(\Sigma)$  by adding the properties  $\xi, \eta, f, b$ . Thus we obtain



# Def. (Reminder): Different Model Classes



We denote the class of  $\mathcal{C}$ -models by  $\mathfrak{M}_\beta(\Sigma)$ . We obtain a hierarchy of subclasses of  $\mathfrak{M}_\beta(\Sigma)$  by adding the properties  $\xi, \eta, f, b$ . Thus we obtain

- $\mathfrak{M}_{\beta\eta}(\Sigma)$

# Def. (Reminder): Different Model Classes



We denote the class of  $\mathcal{C}$ -models by  $\mathfrak{M}_\beta(\Sigma)$ . We obtain a hierarchy of subclasses of  $\mathfrak{M}_\beta(\Sigma)$  by adding the properties  $\xi, \eta, f, b$ . Thus we obtain

- $\mathfrak{M}_{\beta\eta}(\Sigma)$
- $\mathfrak{M}_{\beta\xi}(\Sigma)$

# Def. (Reminder): Different Model Classes



We denote the class of  $\mathcal{C}$ -models by  $\mathfrak{M}_\beta(\Sigma)$ . We obtain a hierarchy of subclasses of  $\mathfrak{M}_\beta(\Sigma)$  by adding the properties  $\xi, \eta, f, b$ . Thus we obtain

- $\mathfrak{M}_{\beta\eta}(\Sigma)$
- $\mathfrak{M}_{\beta\xi}(\Sigma)$
- $\mathfrak{M}_{\beta f}(\Sigma)$

# Def. (Reminder): Different Model Classes



We denote the class of  $\mathcal{C}$ -models by  $\mathfrak{M}_\beta(\Sigma)$ . We obtain a hierarchy of subclasses of  $\mathfrak{M}_\beta(\Sigma)$  by adding the properties  $\xi, \eta, f, b$ . Thus we obtain

- $\mathfrak{M}_{\beta\eta}(\Sigma)$
- $\mathfrak{M}_{\beta\xi}(\Sigma)$
- $\mathfrak{M}_{\beta f}(\Sigma)$
- $\mathfrak{M}_{\beta b}(\Sigma)$

# Def. (Reminder): Different Model Classes



We denote the class of  $\mathcal{C}$ -models by  $\mathfrak{M}_\beta(\Sigma)$ . We obtain a hierarchy of subclasses of  $\mathfrak{M}_\beta(\Sigma)$  by adding the properties  $\xi, \eta, f, b$ . Thus we obtain

- $\mathfrak{M}_{\beta\eta}(\Sigma)$
- $\mathfrak{M}_{\beta\xi}(\Sigma)$
- $\mathfrak{M}_{\beta f}(\Sigma)$
- $\mathfrak{M}_{\beta b}(\Sigma)$
- $\mathfrak{M}_{\beta\eta b}(\Sigma)$

# Def. (Reminder): Different Model Classes



We denote the class of  $\mathcal{C}$ -models by  $\mathfrak{M}_\beta(\Sigma)$ . We obtain a hierarchy of subclasses of  $\mathfrak{M}_\beta(\Sigma)$  by adding the properties  $\xi, \eta, f, b$ . Thus we obtain

- $\mathfrak{M}_{\beta\eta}(\Sigma)$
- $\mathfrak{M}_{\beta\xi}(\Sigma)$
- $\mathfrak{M}_{\beta f}(\Sigma)$
- $\mathfrak{M}_{\beta b}(\Sigma)$
- $\mathfrak{M}_{\beta\eta b}(\Sigma)$
- $\mathfrak{M}_{\beta\xi b}(\Sigma)$

# Def. (Reminder): Different Model Classes



We denote the class of  $\mathcal{C}$ -models by  $\mathfrak{M}_\beta(\Sigma)$ . We obtain a hierarchy of subclasses of  $\mathfrak{M}_\beta(\Sigma)$  by adding the properties  $\xi, \eta, f, b$ . Thus we obtain

- $\mathfrak{M}_{\beta\eta}(\Sigma)$
- $\mathfrak{M}_{\beta\xi}(\Sigma)$
- $\mathfrak{M}_{\beta f}(\Sigma)$
- $\mathfrak{M}_{\beta b}(\Sigma)$
- $\mathfrak{M}_{\beta\eta b}(\Sigma)$
- $\mathfrak{M}_{\beta\xi b}(\Sigma)$
- $\mathfrak{M}_{\beta fb}(\Sigma)$

# Def.: Satisfies, models, and $\models$



Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\Sigma$ -model and let  $\varphi$  be an assignment into  $\mathcal{M}$ .



# Def.: Satisfies, models, and $\models$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\Sigma$ -model and let  $\varphi$  be an assignment into  $\mathcal{M}$ .

We say  $\varphi$  **satisfies** a formula  $A \in wff_o(\Sigma)$  in  $\mathcal{M}$  (we write  $\mathcal{M} \models_{\varphi} A$ ) if  $v(\mathcal{E}_{\varphi}(A)) = T$ .

# Def.: Satisfies, models, and $\models$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\Sigma$ -model and let  $\varphi$  be an assignment into  $\mathcal{M}$ .

We say  $\varphi$  **satisfies** a formula  $A \in wff_o(\Sigma)$  in  $\mathcal{M}$  (we write  $\mathcal{M} \models_{\varphi} A$ ) if  $v(\mathcal{E}_{\varphi}(A)) = T$ .

We say that  $A$  is **valid** in  $\mathcal{M}$  (and write  $\mathcal{M} \models A$ ) if  $\mathcal{M} \models_{\varphi} A$  for all assignments  $\varphi$ .

# Def.: Satisfies, models, and $\models$

Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\Sigma$ -model and let  $\varphi$  be an assignment into  $\mathcal{M}$ .

We say  $\varphi$  **satisfies** a formula  $A \in wff_o(\Sigma)$  in  $\mathcal{M}$  (we write  $\mathcal{M} \models_{\varphi} A$ ) if  $v(\mathcal{E}_{\varphi}(A)) = T$ .

We say that  $A$  is **valid** in  $\mathcal{M}$  (and write  $\mathcal{M} \models A$ ) if  $\mathcal{M} \models_{\varphi} A$  for all assignments  $\varphi$ . When  $A \in cwff_o(\Sigma)$ , we drop the reference to the assignment and use the notation  $\mathcal{M} \models A$ .

# Def.: Satisfies, models, and $\models$


Let  $\mathcal{M} = (D, @, \mathcal{E}, v)$  be a  $\Sigma$ -model and let  $\varphi$  be an assignment into  $\mathcal{M}$ .

We say  $\varphi$  **satisfies** a formula  $A \in wff_o(\Sigma)$  in  $\mathcal{M}$  (we write  $\mathcal{M} \models_{\varphi} A$ ) if  $v(\mathcal{E}_{\varphi}(A)) = T$ .

We say that  $A$  is **valid** in  $\mathcal{M}$  (and write  $\mathcal{M} \models A$ ) if  $\mathcal{M} \models_{\varphi} A$  for all assignments  $\varphi$ . When  $A \in cwff_o(\Sigma)$ , we drop the reference to the assignment and use the notation  $\mathcal{M} \models A$ .

Finally, we say that  $\mathcal{M}$  is a  **$\Sigma$ -model for a set  $\Phi \subseteq cwff_o(\Sigma)$**  (we write  $\mathcal{M} \models \Phi$ ) if  $\mathcal{M} \models A$  for all  $A \in \Phi$ .

# Semantics: HOL-CUBE



## Landscape of HOL model classes


[Kohlhase-PhD-94]

[Benzmüller-PhD-99]

[Brown-PhD-04]

**[Benzm.BrownKohlhase-JSL-04]**

# Semantics: HOL-CUBE



## Landscape of HOL model classes

[Kohlhase-PhD-94]


[Benzmüller-PhD-99]

[Brown-PhD-04]

**[Benzm.BrownKohlhase-JSL-04]**

$\mathfrak{M}_\beta(\Sigma)$  model class for  $\Sigma$ -fragment of  
elementary type theory

# Semantics: HOL-CUBE



## Landscape of HOL model classes

[Kohlhase-PhD-94]

[Benzmüller-PhD-99]


[Brown-PhD-04]

**[Benzm.BrownKohlhase-JSL-04]**

$\mathfrak{M}_\beta(\Sigma)$  model class for  $\Sigma$ -fragment of  
elementary type theory

$\mathfrak{M}_{\beta f b}(\Sigma)$  model class for  $\Sigma$ -fragment of  
extensional type theory (Henkin models)

# Semantics: HOL-CUBE




$\beta$ : models support  $\beta$ -equality

$\eta$ : models provide identity relations

$$\forall \alpha : \text{id} \in \mathcal{D}_{\alpha \rightarrow \alpha \rightarrow o}$$



# Semantics: HOL-CUBE



$\beta$ : models support  $\beta$ -equality

$\eta$ : models provide identity relations

$$\forall \alpha : \text{id} \in \mathcal{D}_{\alpha \rightarrow \alpha \rightarrow o}$$

- [Andrews72]: without property  $\eta$  Leibniz equality  $\doteq$  not necessarily evaluates to identity relation even in Henkin semantics ( $\mathcal{H}(\Sigma)$ )

# Standard Models and Henkin Models



Leon Henkin generalized the class of admissible domains for functional types.

# Standard Models and Henkin Models



Leon Henkin generalized the class of admissible domains for functional types.

Instead of requiring  $\mathcal{D}_{\alpha\beta}$  (and thus in particular,  $\mathcal{D}_{o_i}$ ) to be the full set of functions (predicates), it is sufficient to require that  $\mathcal{D}_{\alpha\beta}$  has enough members that any well-formed formula can be evaluated (in other words, the domains of function types are rich enough to satisfy comprehension).

# Standard Models and Henkin Models




Leon Henkin generalized the class of admissible domains for functional types.

Instead of requiring  $\mathcal{D}_{\alpha\beta}$  (and thus in particular,  $\mathcal{D}_{o_i}$ ) to be the full set of functions (predicates), it is sufficient to require that  $\mathcal{D}_{\alpha\beta}$  has enough members that any well-formed formula can be evaluated (in other words, the domains of function types are rich enough to satisfy comprehension).

Note that with this generalized notion of a model, there are fewer formulae that are valid in all models (intuitively, for any given formula there are more possibilities for counter-models).

# Standard Models and Henkin Models



# Standard Models and Henkin Models



The generalization to Henkin models restricts the set of valid formulae sufficiently so that **all of them can be proven** by a Hilbert-style calculus [Henkin50].

# Standard Models and Henkin Models



The generalization to Henkin models restricts the set of valid formulae sufficiently so that **all of them can be proven** by a Hilbert-style calculus [Henkin50].

Of course our HOL-CUBE is not complete here; we can axiomatically require the existence of particular (classes of) functions, e.g., by assuming the **description or choice operators**.

# Standard Models and Henkin Models



The generalization to Henkin models restricts the set of valid formulae sufficiently so that **all of them can be proven** by a Hilbert-style calculus [Henkin50].

Of course our HOL-CUBE is not complete here; we can axiomatically require the existence of particular (classes of) functions, e.g., by assuming the **description or choice operators**.

We will not pursue this here; for a detailed discussion of the semantic issues raised by the presence of these logical constants see [Andrews72].



# Standard Models and Henkin Models




The generalization to Henkin models restricts the set of valid formulae sufficiently so that **all of them can be proven** by a Hilbert-style calculus [Henkin50].

Of course our HOL-CUBE is not complete here; we can axiomatically require the existence of particular (classes of) functions, e.g., by assuming the **description or choice operators**.


We will not pursue this here; for a detailed discussion of the semantic issues raised by the presence of these logical constants see [Andrews72].

Note that even though we can consider model classes with richer and richer function spaces, **we can never reach standard models where function spaces are full while maintaining complete (recursively axiomatizable) calculi**.

# Standard Models and Henkin Models



# Standard Models and Henkin Models



What has been our motivation for further generalization of Henkin semantics with respect to Boolean and functional extensionality?

# Models without Functional Extensionality



Motivation: modeling programs as (higher-order) functions

- We might be interested in intensional properties like run-time complexity.

# Models without Functional Extensionality



Motivation: modeling programs as (higher-order) functions

- We might be interested in intensional properties like run-time complexity.
- $I := \lambda X.X$  and  $L := \lambda X.\text{rev}(\text{rev}(X))$ , where  $\text{rev}$  is the self-inverse function.

# Models without Functional Extensionality



Motivation: modeling programs as (higher-order) functions

- We might be interested in intensional properties like run-time complexity.
- $I := \lambda X.X$  and  $L := \lambda X.\text{rev}(\text{rev}(X))$ , where  $\text{rev}$  is the self-inverse function.
- The identity function has constant complexity, the function  $\text{rev}$  is linear in the length of its argument.

# Models without Functional Extensionality



How do we account for models without functional extensionality?

- We have generalized the notion of domains at function types and evaluation functions.

# Models without Functional Extensionality



How do we account for models without functional extensionality?

- We have generalized the notion of domains at function types and evaluation functions.
- The usual construction already uses sets of (extensional) functions for the domains of function type and the property of functionality to construct values for  $\lambda$ -terms.



# Models without Functional Extensionality



How do we account for models without functional extensionality?

- We have generalized the notion of domains at function types and evaluation functions.
- The usual construction already uses sets of (extensional) functions for the domains of function type and the property of functionality to construct values for  $\lambda$ -terms.
- We build on the notion of applicative structures to define  $\Sigma$ -evaluations, where the evaluation function is assumed to respect application and  $\beta$ -conversion.


# Models without Functional Extensionality



How do we account for models without functional extensionality?

- We have generalized the notion of domains at function types and evaluation functions.
- The usual construction already uses sets of (extensional) functions for the domains of function type and the property of functionality to construct values for  $\lambda$ -terms.
- We build on the notion of applicative structures to define  $\Sigma$ -evaluations, where the evaluation function is assumed to respect application and  $\beta$ -conversion.
- In such models, a function is not uniquely determined by its behavior on all possible arguments.

# Semantics: HOL-CUBE



$f$ : models are functional

$$\forall f, g \in \mathcal{D}_{\beta\alpha} : \\ f = g \text{ iff } f@a = g@a \ (\forall a \in \mathcal{D}_\alpha)$$

# Models without $\eta$ - or $\xi$ -Functionality



Motivation: in standard literature functional extensionality is often is discussed in terms of

# Models without $\eta$ - or $\xi$ -Functionality



Motivation: in standard literature functional extensionality is often is discussed in terms of

- $\xi$ -functionality

# Models without $\eta$ - or $\xi$ -Functionality



Motivation: in standard literature functional extensionality is often is discussed in terms of

- $\xi$ -functionality
- $\eta$ -functionality


# Models without $\eta$ - or $\xi$ -Functionality



Motivation: in standard literature functional extensionality is often is discussed in terms of

- $\xi$ -functionality
- $\eta$ -functionality
- Therefore, we integrated these two cases in our landscape.

# Semantics: HOL-CUBE




$\eta$ : models are  $\eta$ -functional

$$\mathcal{E}_\varphi(A) = \mathcal{E}_\varphi(A \downarrow_{\beta\eta})$$



# Semantics: HOL-CUBE



$\xi$ : models are  $\xi$ -functional

$$\mathcal{E}_\varphi(\lambda X_\alpha.M_\beta) = \mathcal{E}_\varphi(\lambda X_\alpha.N_\beta) \text{ iff } \mathcal{E}_{\varphi,[a/X]}(M) = \mathcal{E}_{\varphi,[a/X]}(N) \ (\forall a \in \mathcal{D}_\alpha)$$

# Models without Boolean Extensionality



Motivation: Semantics of natural language

# Models without Boolean Extensionality



Motivation: Semantics of natural language

- We may not want to commit to a logic where the sentence  
“John believes that Phil is a woodchuck”

# Models without Boolean Extensionality



Motivation: Semantics of natural language

- We may not want to commit to a logic where the sentence “John believes that Phil is a woodchuck” automatically entails “John believes that Phil is a groundhog”

# Models without Boolean Extensionality



Motivation: Semantics of natural language

- We may not want to commit to a logic where the sentence “John believes that Phil is a woodchuck” automatically entails “John believes that Phil is a groundhog” since John might not know that “woodchuck” is just another word for “groundhog”.

# Models without Boolean Extensionality



Motivation: Semantics of natural language

- We may not want to commit to a logic where the sentence “John believes that Phil is a woodchuck” automatically entails “John believes that Phil is a groundhog” since John might not know that “woodchuck” is just another word for “groundhog”.
- However, Boolean extensionality does just that: **whenever two propositions are equivalent, they must be equal, and can be substituted for each other.**

# Models without Boolean Extensionality



Motivation: Semantics of natural language

- We may not want to commit to a logic where the sentence “John believes that Phil is a woodchuck” automatically entails “John believes that Phil is a groundhog” since John might not know that “woodchuck” is just another word for “groundhog”.
- However, Boolean extensionality does just that: **whenever two propositions are equivalent, they must be equal, and can be substituted for each other.**
- Another example:  $\text{obvious}(\mathbf{O})$  and  $\text{obvious}(\mathbf{F})$  where  $\mathbf{O} := 2 + 2 = 4$  and  $\mathbf{F} := \forall n > 2. x^n + y^n = z^n \Rightarrow x = y = z = 0$  should not be equivalent, even if their arguments are.

# Models without Boolean Extensionality



Motivation: Semantics of natural language

- We may not want to commit to a logic where the sentence “John believes that Phil is a woodchuck” automatically entails “John believes that Phil is a groundhog” since John might not know that “woodchuck” is just another word for “groundhog”.
- However, Boolean extensionality does just that: **whenever two propositions are equivalent, they must be equal, and can be substituted for each other.**
- Another example:  $\text{obvious}(\mathbf{O})$  and  $\text{obvious}(\mathbf{F})$  where  $\mathbf{O} := 2 + 2 = 4$  and  $\mathbf{F} := \forall n > 2. x^n + y^n = z^n \Rightarrow x = y = z = 0$  should not be equivalent, even if their arguments are.
- Such phenomena have been studied under the heading of **“hyper-intensional semantics”** in theoretical semantics.



# Models without Boolean Extensionality



How do we account for models without Boolean extensionality?

- We have weakened the assumption that  $\mathcal{D}_o = \{\mathbf{T}, \mathbf{F}\}$ , since this entails that the values of **O** and **F** are identical.

# Models without Boolean Extensionality



How do we account for models without Boolean extensionality?

- We have weakened the assumption that  $\mathcal{D}_o = \{\mathbf{T}, \mathbf{F}\}$ , since this entails that the values of **O** and **F** are identical.
- In our  $\Sigma$ -models without property **b** we only insist that there is a division of the truth values into “good” and “bad” ones, which we express by insisting on the existence of a valuation  $v$  of  $\mathcal{D}_o$ , i.e., a function  $v: \mathcal{D}_o \rightarrow \{\mathbf{T}, \mathbf{F}\}$  that is coordinated with the interpretations of the logical constants  $\neg$ ,  $\vee$ , and  $\Pi^\alpha$  (for each type  $\alpha$ ).


# Models without Boolean Extensionality



How do we account for models without Boolean extensionality?

- We have weakened the assumption that  $\mathcal{D}_o = \{\mathbf{T}, \mathbf{F}\}$ , since this entails that the values of **O** and **F** are identical.
- In our  $\Sigma$ -models without property **b** we only insist that there is a division of the truth values into “good” and “bad” ones, which we express by insisting on the existence of a valuation  $v$  of  $\mathcal{D}_o$ , i.e., a function  $v: \mathcal{D}_o \rightarrow \{\mathbf{T}, \mathbf{F}\}$  that is coordinated with the interpretations of the logical constants  $\neg$ ,  $\vee$ , and  $\Pi^\alpha$  (for each type  $\alpha$ ).
- Notion of validity: we call a sentence **A** valid in such a model if  $v(a) = \mathbf{T}$ , where  $a \in \mathcal{D}_o$  is the denotation of the sentence **A**.


# Semantics: HOL-CUBE



$b$ : models are Boolean extensional

$v$  is injective

# Semantics: HOL-CUBE



$\mathfrak{b}$ : models are Boolean extensional

$\mathfrak{v}$  is injective

If  $\Sigma$  contains sufficiently many logical constants:

$\mathcal{D}_o = \{\perp, \top\}$



# Semantics and Theorem Proving: Test Problems for Theorem Provers

# Test Problems for Theorem Provers



- Test problems for FOL theorem provers

# Test Problems for Theorem Provers



- Test problems for FOL theorem provers
  - ▶ [McCharenOverbeekWos76], [WilsonMinker79], [Pelletier86], etc.



# Test Problems for Theorem Provers



- Test problems for FOL theorem provers
  - ▶ [McCharenOverbeekWos76], [WilsonMinker79], [Pelletier86], etc.
  - ▶ TPTP [PelletierSutcliffeSuttner02]

# Test Problems for Theorem Provers



- Test problems for FOL theorem provers
  - ▶ [McCharenOverbeekWos76], [WilsonMinker79], [Pelletier86], etc.
  - ▶ TPTP [PelletierSutcliffeSuttner02]
  - ▶ significantly fostered the development of FOL ATPs

# Test Problems for Theorem Provers



- Test problems for FOL theorem provers
  - ▶ [McCharenOverbeekWos76], [WilsonMinker79], [Pelletier86], etc.
  - ▶ TPTP [PelletierSutcliffeSuttner02]
  - ▶ significantly fostered the development of FOL ATPs
- Test problems for HOL theorem provers

# Test Problems for Theorem Provers



- Test problems for FOL theorem provers
  - ▶ [McCharenOverbeekWos76], [WilsonMinker79], [Pelletier86], etc.
  - ▶ TPTP [PelletierSutcliffeSuttner02]
  - ▶ significantly fostered the development of FOL ATPs
- Test problems for HOL theorem provers
  - ▶ common library missing

# Test Problems for Theorem Provers



- Test problems for FOL theorem provers
  - ▶ [McCharenOverbeekWos76], [WilsonMinker79], [Pelletier86], etc.
  - ▶ TPTP [PelletierSutcliffeSuttner02]
  - ▶ significantly fostered the development of FOL ATPs
- Test problems for HOL theorem provers
  - ▶ common library missing
- Following slides: example problems from our paper [TPHOLS-05]

# Test Problems for Theorem Provers



- Test problems for FOL theorem provers
  - ▶ [McCharenOverbeekWos76], [WilsonMinker79], [Pelletier86], etc.
  - ▶ TPTP [PelletierSutcliffeSuttner02]
  - ▶ significantly fostered the development of FOL ATPs
- Test problems for HOL theorem provers
  - ▶ common library missing
- Following slides: example problems from our paper [TPHOLS-05]
- Are we proposing challenging HOL benchmark problems?

# Test Problems for Theorem Provers



- Test problems for FOL theorem provers
  - ▶ [McCharenOverbeekWos76], [WilsonMinker79], [Pelletier86], etc.
  - ▶ TPTP [PelletierSutcliffeSuttner02]
  - ▶ significantly fostered the development of FOL ATPs
- Test problems for HOL theorem provers
  - ▶ common library missing
- Following slides: example problems from our paper [TPHOLS-05]
- Are we proposing challenging HOL benchmark problems?
  - ▶ **No!!!**

# Test Problems for Theorem Provers



- Examples are simple



# Test Problems for Theorem Provers



- Examples are simple
  - ▶ highlight the essence of some semantical or technical point

# Test Problems for Theorem Provers



- Examples are simple
  - ▶ highlight the essence of some semantical or technical point
  - ▶ easy to understand and easy to encode

# Test Problems for Theorem Provers



- Examples are simple
  - ▶ highlight the essence of some semantical or technical point
  - ▶ easy to understand and easy to encode
  - ▶ relevant for both: automated and interactive TP

# Test Problems for Theorem Provers



- Examples are simple
  - ▶ highlight the essence of some semantical or technical point
  - ▶ easy to understand and easy to encode
  - ▶ relevant for both: automated and interactive TP
- Examples are structured

# Test Problems for Theorem Provers



- Examples are simple
  - ▶ highlight the essence of some semantical or technical point
  - ▶ easy to understand and easy to encode
  - ▶ relevant for both: automated and interactive TP
- Examples are structured
  - ▶ quick indicators for completeness and soundness wrt to HOL model classes from [Benzm.BrownKohlhase-JSL-04]

# Test Problems for Theorem Provers



- Examples are simple
  - ▶ highlight the essence of some semantical or technical point
  - ▶ easy to understand and easy to encode
  - ▶ relevant for both: automated and interactive TP
- Examples are structured
  - ▶ quick indicators for completeness and soundness wrt to HOL model classes from [Benzm.BrownKohlhase-JSL-04]
  - ▶ shall precede formal soundness / completeness analysis

# Test Problems for Theorem Provers



- Examples are simple

- ▶ highlight the essence of some semantical or technical point
- ▶ easy to understand and easy to encode
- ▶ relevant for both: automated and interactive TP

- Examples are structured

- ▶ quick indicators for completeness and soundness wrt to HOL model classes from [Benzm.BrownKohlhase-JSL-04]
- ▶ shall precede formal soundness / completeness analysis
- ▶ many are collected from experience with LEO and TPS

# Test Problems for Theorem Provers



- Examples are simple
  - ▶ highlight the essence of some semantical or technical point
  - ▶ easy to understand and easy to encode
  - ▶ relevant for both: automated and interactive TP
- Examples are structured
  - ▶ quick indicators for completeness and soundness wrt to HOL model classes from [Benzm.BrownKohlhase-JSL-04]
  - ▶ shall precede formal soundness / completeness analysis
  - ▶ many are collected from experience with LEO and TPS
- (Some more challenging examples are also added in [TPHOLS-05])




# Remark: Signature

Unless stated otherwise we assume on the following slides that our signature  $\Sigma$  contains the following logical connectives:

$$\{\top, \perp, \neg, \wedge, \vee, \supset, \Leftrightarrow\} \cup \{\Pi^\alpha, \Sigma^\alpha, =^\alpha\}$$

(less logical connectives are possible)


# HOL-Problems: $\beta$



$\simeq^*$  is equivalence relation

- $\forall X_\alpha. X \simeq^* X$
- $\forall X_\alpha, Y_\alpha. X \simeq^* Y \supset Y \simeq^* X$
- $\forall X_\alpha, Y_\alpha, Z_\alpha. (X \simeq^* Y \wedge Y \simeq^* Z) \supset X \simeq^* Z$

# HOL-Problems: $\beta$




$\equiv^*$  is equivalence relation

- $\forall X_\alpha. X \equiv^* X$
- $\forall X_\alpha, Y_\alpha. X \equiv^* Y \supset Y \equiv^* X$
- $\forall X_\alpha, Y_\alpha, Z_\alpha. (X \equiv^* Y \wedge Y \equiv^* Z) \supset X \equiv^* Z$

$\equiv^*$  is congruence relation

- $\forall X_\alpha, Y_\alpha, F_{\alpha\alpha}. X \equiv^* Y \supset (FX) \equiv^* (FY)$
- $\forall X_\alpha, Y_\alpha, P_{o\alpha}. X \equiv^* Y \wedge (PX) \supset (PY)$

# HOL-Problems: $\beta$



$\equiv^*$  is equivalence relation

- $\forall X_\alpha. X \equiv^* X$
- $\forall X_\alpha, Y_\alpha. X \equiv^* Y \supset Y \equiv^* X$
- $\forall X_\alpha, Y_\alpha, Z_\alpha. (X \equiv^* Y \wedge Y \equiv^* Z) \supset X \equiv^* Z$


$\equiv^*$  is congruence relation

- $\forall X_\alpha, Y_\alpha, F_{\alpha\alpha}. X \equiv^* Y \supset (FX) \equiv^* (FY)$
- $\forall X_\alpha, Y_\alpha, P_{o\alpha}. X \equiv^* Y \wedge (PX) \supset (PY)$

Trivial directions of Boolean and functional extensionality

- $\forall A_o, B_o. A \equiv^* B \supset (A \Leftrightarrow B)$
- $\forall F_{\beta\alpha}, G_{\beta\alpha}. F \equiv^* G \supset (\forall X_\alpha. FX \equiv^* GX)$


# HOL-Problems: $\mathfrak{b}$



Non-trivial direction of Boolean extensionality

■  $\forall A_o, B_o. (A \Leftrightarrow B) \supset A \stackrel{*}{=} B$


# HOL-Problems: $f$



Non-trivial direct. of functional extensionality

$$\blacksquare \quad \forall F_{\beta\alpha}, G_{\beta\alpha}. (\forall X_{\alpha}. FX \stackrel{*}{=} GX) \supset F \stackrel{*}{=} G$$


# HOL-Problems: $\eta$



Example requiring property  $\eta$

$$\blacksquare \quad (p_{o(\iota\iota)}(\lambda X_{\iota}.f_{\iota\iota}X)) \supset (p \ f)$$

# HOL-Problems: $\xi$




Example requiring property  $\xi$  (and  $q$ !)

- $(\forall X_{\iota}. (f_{\iota\iota} X) \stackrel{*}{=} X) \wedge p_{o(\iota\iota)}(\lambda X_{\iota}. X)$   
 $\supset p(\lambda X_{\iota}. f X)$




# HOL-Problems: f



### Example requiring property $f$ (and $q$ !)


- $(\forall X_{\iota}. (f_{\iota\iota} X) \stackrel{*}{=} X) \wedge p_{o(\iota\iota)}(\lambda X_{\iota}. X)$   
 $\supset (p\ f)$



Examples requiring property  $\flat$

- $(p \circ \circ a) \wedge (p \circ b) \Rightarrow (p (a \wedge b))$
- $\neg(a \stackrel{*}{=} \neg a)$  (in particular  $\neg(a = \neg a)$ )
- $(h_{\iota o}((h \top) \stackrel{*}{=} (h \perp))) \stackrel{*}{=} (h \perp)$

# HOL-Problems: Other Examples




Playing with DeMorgan's Law:

■  $\forall X, Y. X \wedge Y \Leftrightarrow \neg(\neg X \vee \neg Y)$

'Ok' for all model classes

# HOL-Problems: DeMorgan's Law




Playing with DeMorgan's Law:

- $\forall X, Y. X \wedge Y \Leftrightarrow \neg(\neg X \vee \neg Y)$
- $\forall X, Y. X \wedge Y \stackrel{*}{=} \neg(\neg X \vee \neg Y)$

requires b

# HOL-Problems: DeMorgan's Law




Playing with DeMorgan's Law:

- $\forall X, Y. X \wedge Y \Leftrightarrow \neg(\neg X \vee \neg Y)$
- $\forall X, Y. X \wedge Y \stackrel{*}{=} \neg(\neg X \vee \neg Y)$
- $(\lambda U \lambda V. U \wedge V) \stackrel{*}{=} (\lambda X \lambda Y. \neg(\neg X \vee \neg Y))$

requires b and  $\xi$

# HOL-Problems: DeMorgan's Law




Playing with DeMorgan's Law:

- $\forall X, Y. X \wedge Y \Leftrightarrow \neg(\neg X \vee \neg Y)$
- $\forall X, Y. X \wedge Y \stackrel{*}{=} \neg(\neg X \vee \neg Y)$
- $(\lambda U \lambda V. U \wedge V) \stackrel{*}{=} (\lambda X \lambda Y. \neg(\neg X \vee \neg Y))$
- $\wedge \stackrel{*}{=} (\lambda X \lambda Y. \neg(\neg X \vee \neg Y))$

requires  $b$  and  $f$


# HOL-Problems: Set Comprehension



## Set comprehension

- big challenge for automation
- [Benzm.BrownKohlhase-Draft-05] set instantiations can be used to simulate cut-rule if one of the following axioms is given: comprehension, induction, extensionality, choice, description
- depend on logical constants in  $\Sigma$

# HOL-Problems: Set Comprehension



## Set comprehension


- big challenge for automation
- [Benzm.BrownKohlhase-Draft-05] set instantiations can be used to simulate cut-rule if one of the following axioms is given: comprehension, induction, extensionality, choice, description
- depend on logical constants in  $\Sigma$

## On the following slides emphasis on:

- signature  $\Sigma$  varying
- no property  $q$  assumed



# HOL-Problems: Set Comprehension



## Set comprehension

- big challenge for automation
- [Benzm.BrownKohlhase-Draft-05] set instantiations can be used to simulate cut-rule if one of the following axioms is given: comprehension, induction, extensionality, choice, description
- depend on logical constants in  $\Sigma$


## On the following slides emphasis on:

- signature  $\Sigma$  varying
- no property  $q$  assumed

## External vs. internal logical constants

- if  $\neg \notin \Sigma$ :  
 $\neg$  refers to 'external' symbol  
 $\mathcal{M} \models \neg A$  means  $\mathcal{M} \not\models A$


# HOL-Problems: Set Comprehension



## Set comprehension

- $\exists N_{oo} \forall P_{oo}. NP \Leftrightarrow \neg P$ 
  - ▶ if  $\neg \in \Sigma$  or  $\{\perp, \supset\} \subseteq \Sigma$  or  $\{\perp, \Leftrightarrow\} \subseteq \Sigma$
  - ▶ e.g.:  $N_{oo} \leftarrow \lambda X_{oo}. \neg X$
  - ▶ e.g.:  $N_{oo} \leftarrow \lambda X_{oo}. X \supset \perp$


# HOL-Problems: Set Comprehension



## Set comprehension

- $\exists N_{oo} \forall P_{oo} . NP \Leftrightarrow \neg P$
- ▶ if  $\neg \notin \Sigma$  and  $\{\perp, \supset\} \not\subseteq \Sigma$  or  $\{\perp, \Leftrightarrow\} \not\subseteq \Sigma$

# HOL-Problems: Set Comprehension



## Set comprehension

- $\exists N_{\circ\circ} \forall P_{\circ\circ}. NP \Leftrightarrow \neg P$ 
  - ▶ if  $\neg \notin \Sigma$  and  $\{\perp, \supset\} \not\subseteq \Sigma$  or  $\{\perp, \Leftrightarrow\} \not\subseteq \Sigma$

## Other examples from [Brown-PhD-04]

- Surjective Cantor Theorem
- Injective Cantor Theorem



# Semantics: Examples of $\Sigma$ -Models

# Examples of $\Sigma$ -Models



We now sketch the construction of models in the model classes  $\mathfrak{M}_*(\Sigma)$  to demonstrate concretely how properties for Boolean, strong and weak functional extensionality can fail.

# Examples of $\Sigma$ -Models



We now sketch the construction of models in the model classes  $\mathfrak{M}_*(\Sigma)$  to demonstrate concretely how properties for Boolean, strong and weak functional extensionality can fail.

We need this to show that the inclusions of the model classes in our landscape are proper, and we indeed need all of them.

# Ex.: Singleton Model



- We choose  $(\mathcal{D}, @)$  as the full frame with  $\mathcal{D}_o := \{T, F\}$  and  $\mathcal{D}_\iota := \{*\}$ .



# Ex.: Singleton Model

- We choose  $(\mathcal{D}, @)$  as the full frame with  $\mathcal{D}_o := \{\mathbf{T}, \mathbf{F}\}$  and  $\mathcal{D}_\iota := \{*\}$ .
- Easy to define an evaluation function  $\mathcal{E}$  for this frame by induction on terms, using functions to interpret  $\lambda$ -abstractions.

# Ex.: Singleton Model

- We choose  $(\mathcal{D}, @)$  as the full frame with  $\mathcal{D}_o := \{T, F\}$  and  $\mathcal{D}_\iota := \{*\}$ .
- Easy to define an evaluation function  $\mathcal{E}$  for this frame by induction on terms, using functions to interpret  $\lambda$ -abstractions.
- The identity function  $v: \mathcal{D}_o \longrightarrow \{T, F\}$  is a valuation, assuming the logical constants are interpreted in the standard way.

# Ex.: Singleton Model

- We choose  $(\mathcal{D}, @)$  as the full frame with  $\mathcal{D}_o := \{\mathbf{T}, \mathbf{F}\}$  and  $\mathcal{D}_l := \{*\}$ .
- Easy to define an evaluation function  $\mathcal{E}$  for this frame by induction on terms, using functions to interpret  $\lambda$ -abstractions.
- The identity function  $v: \mathcal{D}_o \longrightarrow \{\mathbf{T}, \mathbf{F}\}$  is a valuation, assuming the logical constants are interpreted in the standard way.
- Thus,  $\mathcal{M}^{\beta\text{fb}} := (\mathcal{D}, @, \mathcal{E}, v)$  defines a  $\Sigma$ -model.


# Ex.: Singleton Model

- We choose  $(\mathcal{D}, @)$  as the full frame with  $\mathcal{D}_o := \{T, F\}$  and  $\mathcal{D}_i := \{*\}$ .
- Easy to define an evaluation function  $\mathcal{E}$  for this frame by induction on terms, using functions to interpret  $\lambda$ -abstractions.
- The identity function  $v: \mathcal{D}_o \longrightarrow \{T, F\}$  is a valuation, assuming the logical constants are interpreted in the standard way.
- Thus,  $\mathcal{M}^{\beta\mathfrak{fb}} := (\mathcal{D}, @, \mathcal{E}, v)$  defines a  $\Sigma$ -model.
- This model satisfies properties  $\mathfrak{b}$ ,  $\mathfrak{f}$  (hence  $\eta$  and  $\xi$ ) and  $\mathfrak{q}$  (since the frame is full).

# Ex.: Singleton Model

- We choose  $(\mathcal{D}, @)$  as the full frame with  $\mathcal{D}_o := \{T, F\}$  and  $\mathcal{D}_i := \{*\}$ .
- Easy to define an evaluation function  $\mathcal{E}$  for this frame by induction on terms, using functions to interpret  $\lambda$ -abstractions.
- The identity function  $v: \mathcal{D}_o \longrightarrow \{T, F\}$  is a valuation, assuming the logical constants are interpreted in the standard way.
- Thus,  $\mathcal{M}^{\beta_{fb}} := (\mathcal{D}, @, \mathcal{E}, v)$  defines a  $\Sigma$ -model.
- This model satisfies properties  $b$ ,  $f$  (hence  $\eta$  and  $\xi$ ) and  $q$  (since the frame is full).
- So,  $\mathcal{M}^{\beta_{fb}} \in \mathfrak{ST}(\Sigma) \subseteq \mathfrak{H}(\Sigma) \subseteq \mathfrak{M}_{\beta_{fb}}(\Sigma) \subseteq \dots$

# Ex.: Singleton Model



# Ex.: Model without Boolean Extensionality



- Assume  $\Sigma$  contains only the connectives  $\neg, \vee, \Pi^\alpha$ ; other connectives defined as usual, e.g.,  $\forall X, Y. X \wedge Y \Leftrightarrow \neg(\neg X \vee \neg Y)$ .

# Ex.: Model without Boolean Extensionality



- Assume  $\Sigma$  contains only the connectives  $\neg, \vee, \Pi^\alpha$ ; other connectives defined as usual, e.g.,  $\forall X, Y. X \wedge Y \Leftrightarrow \neg(\neg X \vee \neg Y)$ .
- Choose  $(\mathcal{D}, @)$  as full frame with  $\mathcal{D}_o = \{a, b, c\}$  and  $\mathcal{D}_i = \{0, 1\}$ .



# Ex.: Model without Boolean Extensionality



- Assume  $\Sigma$  contains only the connectives  $\neg, \vee, \Pi^\alpha$ ; other connectives defined as usual, e.g.,  $\forall X, Y. X \wedge Y \Leftrightarrow \neg(\neg X \vee \neg Y)$ .
- Choose  $(\mathcal{D}, @)$  as full frame with  $\mathcal{D}_o = \{a, b, c\}$  and  $\mathcal{D}_\iota = \{0, 1\}$ .
- We define evaluation function  $\mathcal{E}$  for this frame by defining  $\mathcal{E}(\neg)$ ,  $\mathcal{E}(\vee)$ , and  $\mathcal{E}(\Pi^\alpha)$ :

$\mathcal{E}(\neg)$	a	b	c
	c	c	a

$\mathcal{E}(\vee)$	a	b	c
a	a	a	a
b	a	a	a
c	a	a	c

$$\mathcal{E}(\Pi^\alpha)@f = \begin{cases} a, & \text{if } f@g \in \{a, b\} \text{ for all } g \in \mathcal{D}_\alpha \\ c, & \text{if } f@g = c \text{ for some } g \in \mathcal{D}_\alpha \end{cases}$$

# Ex.: Model without Boolean Extensionality



- Assume  $\Sigma$  contains only the connectives  $\neg, \vee, \Pi^\alpha$ ; other connectives defined as usual, e.g.,  $\forall X, Y. X \wedge Y \Leftrightarrow \neg(\neg X \vee \neg Y)$ .
- Choose  $(\mathcal{D}, @)$  as full frame with  $\mathcal{D}_o = \{a, b, c\}$  and  $\mathcal{D}_\iota = \{0, 1\}$ .
- We define evaluation function  $\mathcal{E}$  for this frame by defining  $\mathcal{E}(\neg)$ ,  $\mathcal{E}(\vee)$ , and  $\mathcal{E}(\Pi^\alpha)$ :

$\mathcal{E}(\neg)$	a	b	c
	c	c	a

$\mathcal{E}(\vee)$	a	b	c
a	a	a	a
b	a	a	a
c	a	a	c

$$\mathcal{E}(\Pi^\alpha)@f = \begin{cases} a, & \text{if } f@g \in \{a, b\} \text{ for all } g \in \mathcal{D}_\alpha \\ c, & \text{if } f@g = c \text{ for some } g \in \mathcal{D}_\alpha \end{cases}$$

- We can choose  $\mathcal{E}(w)$  to be arbitrary for parameters  $w \in \Sigma$ .

# Ex.: Model without Boolean Extensionality



- Since  $(\mathcal{D}, @)$  is a frame, hence functional, this uniquely determines  $\mathcal{E}$  on all formulae.

# Ex.: Model without Boolean Extensionality



- Since  $(\mathcal{D}, @)$  is a frame, hence functional, this uniquely determines  $\mathcal{E}$  on all formulae.
- Since the frame is full, we are guaranteed that there will be enough functions to interpret  $\lambda$ -abstractions.

# Ex.: Model without Boolean Extensionality



- Since  $(\mathcal{D}, @)$  is a frame, hence functional, this uniquely determines  $\mathcal{E}$  on all formulae.
- Since the frame is full, we are guaranteed that there will be enough functions to interpret  $\lambda$ -abstractions.
- Let  $v: \mathcal{D}_o \longrightarrow \{T, F\}$  be defined by  $v(a) := T$ ,  $v(b) := T$  and  $v(c) := F$ .

# Ex.: Model without Boolean Extensionality



- Since  $(\mathcal{D}, @)$  is a frame, hence functional, this uniquely determines  $\mathcal{E}$  on all formulae.
- Since the frame is full, we are guaranteed that there will be enough functions to interpret  $\lambda$ -abstractions.
- Let  $v: \mathcal{D}_o \longrightarrow \{T, F\}$  be defined by  $v(a) := T$ ,  $v(b) := T$  and  $v(c) := F$ .
- Easy to check that  $\mathcal{M}^{\beta f} := (\mathcal{D}, @, \mathcal{E}, v)$  is indeed a  $\Sigma$ -model.

# Ex.: Model without Boolean Extensionality



- Since  $(\mathcal{D}, @)$  is a frame, hence functional, this uniquely determines  $\mathcal{E}$  on all formulae.
- Since the frame is full, we are guaranteed that there will be enough functions to interpret  $\lambda$ -abstractions.
- Let  $v: \mathcal{D}_o \longrightarrow \{T, F\}$  be defined by  $v(a) := T$ ,  $v(b) := T$  and  $v(c) := F$ .
- Easy to check that  $\mathcal{M}^{\beta f} := (\mathcal{D}, @, \mathcal{E}, v)$  is indeed a  $\Sigma$ -model.
- Since  $\mathcal{M}^{\beta f}$  is a model over a frame it satisfies property  $f$ .

# Ex.: Model without Boolean Extensionality



- Since  $(\mathcal{D}, @)$  is a frame, hence functional, this uniquely determines  $\mathcal{E}$  on all formulae.
- Since the frame is full, we are guaranteed that there will be enough functions to interpret  $\lambda$ -abstractions.
- Let  $v: \mathcal{D}_o \longrightarrow \{T, F\}$  be defined by  $v(a) := T$ ,  $v(b) := T$  and  $v(c) := F$ .
- Easy to check that  $\mathcal{M}^{\beta f} := (\mathcal{D}, @, \mathcal{E}, v)$  is indeed a  $\Sigma$ -model.
- Since  $\mathcal{M}^{\beta f}$  is a model over a frame it satisfies property  $f$ .
- Since this frame is full, we know property  $q$  holds.



# Ex.: Model without Boolean Extensionality




- Since  $(\mathcal{D}, @)$  is a frame, hence functional, this uniquely determines  $\mathcal{E}$  on all formulae.
- Since the frame is full, we are guaranteed that there will be enough functions to interpret  $\lambda$ -abstractions.
- Let  $v: \mathcal{D}_o \longrightarrow \{T, F\}$  be defined by  $v(a) := T$ ,  $v(b) := T$  and  $v(c) := F$ .
- Easy to check that  $\mathcal{M}^{\beta f} := (\mathcal{D}, @, \mathcal{E}, v)$  is indeed a  $\Sigma$ -model.
- Since  $\mathcal{M}^{\beta f}$  is a model over a frame it satisfies property  $f$ .
- Since this frame is full, we know property  $q$  holds.
- Clearly property  $b$  fails.

# Ex.: Model without Boolean Extensionality



- Since  $(\mathcal{D}, @)$  is a frame, hence functional, this uniquely determines  $\mathcal{E}$  on all formulae.
- Since the frame is full, we are guaranteed that there will be enough functions to interpret  $\lambda$ -abstractions.
- Let  $v: \mathcal{D}_o \longrightarrow \{\mathbf{T}, \mathbf{F}\}$  be defined by  $v(\mathbf{a}) := \mathbf{T}$ ,  $v(\mathbf{b}) := \mathbf{T}$  and  $v(\mathbf{c}) := \mathbf{F}$ .
- Easy to check that  $\mathcal{M}^{\beta\mathbf{f}} := (\mathcal{D}, @, \mathcal{E}, v)$  is indeed a  $\Sigma$ -model.
- Since  $\mathcal{M}^{\beta\mathbf{f}}$  is a model over a frame it satisfies property  $\mathbf{f}$ .
- Since this frame is full, we know property  $\mathbf{q}$  holds.
- Clearly property  $\mathbf{b}$  fails.
- So,  $\mathcal{M}^{\beta\mathbf{f}} \in \mathfrak{M}_{\beta\mathbf{f}}(\Sigma) \setminus \mathfrak{M}_{\beta\mathbf{fb}}(\Sigma)$ .

# Ex.: Model without Boolean Extensionality



# Ex.: Model without Boolean Extensionality



In the previous model one can easily verify, if  $d := \mathcal{E}_\varphi(\mathbf{D}_o)$  and  $e := \mathcal{E}_\varphi(\mathbf{E}_o)$ , then the values  $\mathcal{E}_\varphi(\mathbf{D} \wedge \mathbf{E})$ ,  $\mathcal{E}_\varphi(\mathbf{D} \Rightarrow \mathbf{E})$ , and  $\mathcal{E}_\varphi(\mathbf{D} \Leftrightarrow \mathbf{E})$  are given by the following tables:

$\mathcal{E}(\mathbf{D} \wedge \mathbf{E})$	e:				$\mathcal{E}(\mathbf{D} \Rightarrow \mathbf{E})$	e:				$\mathcal{E}(\mathbf{D} \Leftrightarrow \mathbf{E})$	e:			
	a	b	c			a	b	c			a	b	c	
d: a	a	a	c		d: a	a	a	c		d: a	a	a	c	
b	a	a	c		b	a	a	c		b	a	a	c	
c	c	c	c		c	a	a	a		c	c	c	a	

Now we show that one can properly model the **woodchuck/groundhog** example.

# Ex.: Groundhogs and Woodchucks



- Let  $\mathcal{M}^{\beta f}$  be given as above and suppose  $\text{woodchuck}_{\iota \rightarrow o}$ ,  $\text{groundhog}_{\iota \rightarrow o}$ ,  $\text{john}_{\iota}$ , and  $\text{phil}_{\iota}$  are in the signature  $\Sigma$ .

# Ex.: Groundhogs and Woodchucks



- Let  $\mathcal{M}^{\beta f}$  be given as above and suppose  $\text{woodchuck}_{\iota \rightarrow o}$ ,  $\text{groundhog}_{\iota \rightarrow o}$ ,  $\text{john}_{\iota}$ , and  $\text{phil}_{\iota}$  are in the signature  $\Sigma$ .
- Let  $\mathcal{E}(\text{phil}) := 0$  and  $\mathcal{E}(\text{john}) := 1$ .

# Ex.: Groundhogs and Woodchucks



- Let  $\mathcal{M}^{\beta f}$  be given as above and suppose  $\text{woodchuck}_{\iota \rightarrow o}$ ,  $\text{groundhog}_{\iota \rightarrow o}$ ,  $\text{john}_{\iota}$ , and  $\text{phil}_{\iota}$  are in the signature  $\Sigma$ .
- Let  $\mathcal{E}(\text{phil}) := 0$  and  $\mathcal{E}(\text{john}) := 1$ .
- Let  $\mathcal{E}(\text{woodchuck})$  be the function  $w \in \mathcal{D}_{\iota \rightarrow o}$  with  $w(0) = b$  and  $w(1) = c$ .

# Ex.: Groundhogs and Woodchucks



- Let  $\mathcal{M}^{\beta f}$  be given as above and suppose  $\text{woodchuck}_{\iota \rightarrow o}$ ,  $\text{groundhog}_{\iota \rightarrow o}$ ,  $\text{john}_{\iota}$ , and  $\text{phil}_{\iota}$  are in the signature  $\Sigma$ .
- Let  $\mathcal{E}(\text{phil}) := 0$  and  $\mathcal{E}(\text{john}) := 1$ .
- Let  $\mathcal{E}(\text{woodchuck})$  be the function  $w \in \mathcal{D}_{\iota \rightarrow o}$  with  $w(0) = b$  and  $w(1) = c$ .
- Let  $\mathcal{E}(\text{groundhog})$  be the function  $g \in \mathcal{D}_{\iota \rightarrow o}$  with  $g(0) = a$  and  $g(1) = c$ .



# Ex.: Groundhogs and Woodchucks

- Let  $\mathcal{M}^{\beta f}$  be given as above and suppose  $\text{woodchuck}_{\iota \rightarrow o}$ ,  $\text{groundhog}_{\iota \rightarrow o}$ ,  $\text{john}_{\iota}$ , and  $\text{phil}_{\iota}$  are in the signature  $\Sigma$ .
- Let  $\mathcal{E}(\text{phil}) := 0$  and  $\mathcal{E}(\text{john}) := 1$ .
- Let  $\mathcal{E}(\text{woodchuck})$  be the function  $w \in \mathcal{D}_{\iota \rightarrow o}$  with  $w(0) = b$  and  $w(1) = c$ .
- Let  $\mathcal{E}(\text{groundhog})$  be the function  $g \in \mathcal{D}_{\iota \rightarrow o}$  with  $g(0) = a$  and  $g(1) = c$ .
- One can show that the sentence  $\forall X_{\iota}. (\text{woodchuck } X) \Leftrightarrow (\text{groundhog } X)$  is valid.

# Ex.: Groundhogs and Woodchucks

- Let  $\mathcal{M}^{\beta f}$  be given as above and suppose  $\text{woodchuck}_{\iota \rightarrow o}$ ,  $\text{groundhog}_{\iota \rightarrow o}$ ,  $\text{john}_{\iota}$ , and  $\text{phil}_{\iota}$  are in the signature  $\Sigma$ .
- Let  $\mathcal{E}(\text{phil}) := 0$  and  $\mathcal{E}(\text{john}) := 1$ .
- Let  $\mathcal{E}(\text{woodchuck})$  be the function  $w \in \mathcal{D}_{\iota \rightarrow o}$  with  $w(0) = b$  and  $w(1) = c$ .
- Let  $\mathcal{E}(\text{groundhog})$  be the function  $g \in \mathcal{D}_{\iota \rightarrow o}$  with  $g(0) = a$  and  $g(1) = c$ .
- One can show that the sentence  $\forall X_{\iota}. (\text{woodchuck } X) \Leftrightarrow (\text{groundhog } X)$  is valid.
- Also,  $\mathcal{E}(\text{woodchuck phil}) = b$  and  $\mathcal{E}(\text{groundhog phil}) = a$ , so the propositions  $(\text{woodchuck phil})$  and  $(\text{groundhog phil})$  are valid.

# Ex.: Groundhogs and Woodchucks



- Suppose  $\text{believe}_{\iota \rightarrow o \rightarrow o} \in \Sigma$  and  $\mathcal{E}(\text{believe})$  is the (Curried) function  $\text{bel} \in \mathcal{D}_{\iota \rightarrow o \rightarrow o}$  such that  $\text{bel}(1)(b) = b$  and  $\text{bel}(1)(a) = \text{bel}(1)(c) = \text{bel}(0)(a) = \text{bel}(0)(b) = \text{bel}(0)(c) = c$ .

# Ex.: Groundhogs and Woodchucks



- Suppose  $\text{believe}_{\iota \rightarrow o \rightarrow o} \in \Sigma$  and  $\mathcal{E}(\text{believe})$  is the (Curried) function  $\text{bel} \in \mathcal{D}_{\iota \rightarrow o \rightarrow o}$  such that  $\text{bel}(1)(b) = b$  and  $\text{bel}(1)(a) = \text{bel}(1)(c) = \text{bel}(0)(a) = \text{bel}(0)(b) = \text{bel}(0)(c) = c$ .
- Intuitively, John believes propositions with value  $b$ , but not those with value  $a$  or  $c$ .

# Ex.: Groundhogs and Woodchucks



- Suppose  $\text{believe}_{\iota \rightarrow o \rightarrow o} \in \Sigma$  and  $\mathcal{E}(\text{believe})$  is the (Curried) function  $\text{bel} \in \mathcal{D}_{\iota \rightarrow o \rightarrow o}$  such that  $\text{bel}(1)(b) = b$  and  $\text{bel}(1)(a) = \text{bel}(1)(c) = \text{bel}(0)(a) = \text{bel}(0)(b) = \text{bel}(0)(c) = c$ .
- Intuitively, John believes propositions with value  $b$ , but not those with value  $a$  or  $c$ .
- So,  $\text{believes john}(\text{woodchuck phil})$  is valid, while  $\text{believes john}(\text{groundhog phil})$  is not.

# Generalizing the Previous Model



As we have seen, Boolean extensionality fails when one has more than two values in  $\mathcal{D}_o$ .

# Generalizing the Previous Model

As we have seen, Boolean extensionality fails when one has more than two values in  $\mathcal{D}_o$ . We can generalize the construction defining  $\mathcal{D}_o := \{F\} \cup \mathcal{B}$ , where  $\mathcal{B}$  is any set with  $T \in \mathcal{B}$  and  $F \notin \mathcal{B}$ .

# Generalizing the Previous Model



As we have seen, Boolean extensionality fails when one has more than two values in  $\mathcal{D}_o$ . We can generalize the construction defining  $\mathcal{D}_o := \{F\} \cup \mathcal{B}$ , where  $\mathcal{B}$  is any set with  $T \in \mathcal{B}$  and  $F \notin \mathcal{B}$ . The model will satisfy Boolean extensionality iff  $\mathcal{B} = \{T\}$ .



# Generalizing the Previous Model

As we have seen, Boolean extensionality fails when one has more than two values in  $\mathcal{D}_o$ . We can generalize the construction defining  $\mathcal{D}_o := \{F\} \cup \mathcal{B}$ , where  $\mathcal{B}$  is any set with  $T \in \mathcal{B}$  and  $F \notin \mathcal{B}$ . The model will satisfy Boolean extensionality iff  $\mathcal{B} = \{T\}$ . In this way, we can easily construct models for the case with property  $b$  and the case without property  $b$  simultaneously.

# Generalizing the Previous Model

As we have seen, Boolean extensionality fails when one has more than two values in  $\mathcal{D}_o$ . We can generalize the construction defining  $\mathcal{D}_o := \{F\} \cup \mathcal{B}$ , where  $\mathcal{B}$  is any set with  $T \in \mathcal{B}$  and  $F \notin \mathcal{B}$ . The model will satisfy Boolean extensionality iff  $\mathcal{B} = \{T\}$ . In this way, we can easily construct models for the case with property  $b$  and the case without property  $b$  simultaneously. We will use this idea to parameterize the remaining model constructions by  $\mathcal{B}$ .

# Generalizing the Previous Model

As we have seen, Boolean extensionality fails when one has more than two values in  $\mathcal{D}_o$ . We can generalize the construction defining  $\mathcal{D}_o := \{F\} \cup \mathcal{B}$ , where  $\mathcal{B}$  is any set with  $T \in \mathcal{B}$  and  $F \notin \mathcal{B}$ . The model will satisfy Boolean extensionality iff  $\mathcal{B} = \{T\}$ . In this way, we can easily construct models for the case with property  $b$  and the case without property  $b$  simultaneously. We will use this idea to parameterize the remaining model constructions by  $\mathcal{B}$ .

These semantic constructions are similar to those in multi-valued logics.

# Generalizing the Previous Model

As we have seen, Boolean extensionality fails when one has more than two values in  $\mathcal{D}_o$ . We can generalize the construction defining  $\mathcal{D}_o := \{F\} \cup \mathcal{B}$ , where  $\mathcal{B}$  is any set with  $T \in \mathcal{B}$  and  $F \notin \mathcal{B}$ . The model will satisfy Boolean extensionality iff  $\mathcal{B} = \{T\}$ . In this way, we can easily construct models for the case with property  $b$  and the case without property  $b$  simultaneously. We will use this idea to parameterize the remaining model constructions by  $\mathcal{B}$ .

These semantic constructions are similar to those in multi-valued logics. In contrast to these logics where the logical connectives are adapted to talk about multiple truth values, in our setting we are mainly interested in multiple truth values as diverse  $v$ -pre-images of  $T$  and  $F$ .



## Semantics: Examples of $\Sigma$ -Models (Contd.)

# Ex.: Models without Funct. Extensionality



- Idea: attach distinguishing labels to functions without changing their applicative behavior

# Ex.: Models without Funct. Extensionality



- Idea: attach distinguishing labels to functions without changing their applicative behavior
- Let  $\mathcal{B}$  be any set with  $\mathsf{T} \in \mathcal{B}$  and  $\mathsf{F} \notin \mathcal{B}$

# Ex.: Models without Funct. Extensionality



- Idea: attach distinguishing labels to functions without changing their applicative behavior
- Let  $\mathcal{B}$  be any set with  $\mathsf{T} \in \mathcal{B}$  and  $\mathsf{F} \notin \mathcal{B}$
- Let  $\mathcal{D}_o := \{\mathsf{F}\} \cup \mathcal{B}$  and  $\mathcal{D}_i := \{*\}$



# Ex.: Models without Funct. Extensionality



- Idea: attach distinguishing labels to functions without changing their applicative behavior
- Let  $\mathcal{B}$  be any set with  $\top \in \mathcal{B}$  and  $\bot \notin \mathcal{B}$
- Let  $\mathcal{D}_\circ := \{\bot\} \cup \mathcal{B}$  and  $\mathcal{D}_\iota := \{*\}$
- For each function type  $\beta\alpha$ , let

$$\mathcal{D}_{\beta\alpha} := \{(i, f) \mid i \in \{0, 1\} \text{ and } f: \mathcal{D}_\alpha \longrightarrow \mathcal{D}_\beta\}$$

# Ex.: Models without Funct. Extensionality



- Idea: attach distinguishing labels to functions without changing their applicative behavior
- Let  $\mathcal{B}$  be any set with  $\mathbf{T} \in \mathcal{B}$  and  $\mathbf{F} \notin \mathcal{B}$
- Let  $\mathcal{D}_o := \{\mathbf{F}\} \cup \mathcal{B}$  and  $\mathcal{D}_\iota := \{*\}$
- For each function type  $\beta\alpha$ , let

$$\mathcal{D}_{\beta\alpha} := \{(i, f) \mid i \in \{0, 1\} \text{ and } f: \mathcal{D}_\alpha \longrightarrow \mathcal{D}_\beta\}$$

- We define application by

$$(i, f)@a := f(a)$$

whenever  $(i, f) \in \mathcal{D}_{\beta\alpha}$  and  $a \in \mathcal{D}_\alpha$

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Easy to check that  $(\mathcal{D}, @)$  is an applicative structure:

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Easy to check that  $(\mathcal{D}, @)$  is an applicative structure:
- Evaluation function defined by induction on terms

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Easy to check that  $(\mathcal{D}, @)$  is an applicative structure:
- Evaluation function defined by induction on terms
  - ▶  $\mathcal{E}(\neg) := (0, n)$  where  $n(b) := \mathbf{F}$  for every  $b \in \mathcal{B}$  and  $n(\mathbf{F}) := \mathbf{T}$

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Easy to check that  $(\mathcal{D}, @)$  is an applicative structure:
- Evaluation function defined by induction on terms
  - ▶  $\mathcal{E}(\neg) := (0, n)$  where  $n(b) := \mathbf{F}$  for every  $b \in \mathcal{B}$  and  $n(\mathbf{F}) := \mathbf{T}$
  - ▶  $\mathcal{E}(\vee) := (0, d)$  where
    - $d(b) := (0, k^{\mathbf{T}})$  for every  $b \in \mathcal{B}$  and
    - $d(\mathbf{F}) := (0, \text{id})$
    - $(k^{\mathbf{T}}$  is the constant  $\mathbf{T}$  function)
    - $(\text{id}$  is the identity function from  $\mathcal{D}_o$  to  $\mathcal{D}_o$ )

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Easy to check that  $(\mathcal{D}, @)$  is an applicative structure:
- Evaluation function defined by induction on terms
  - ▶  $\mathcal{E}(\neg) := (0, n)$  where  $n(b) := \mathbf{F}$  for every  $b \in \mathcal{B}$  and  $n(\mathbf{F}) := \mathbf{T}$
  - ▶  $\mathcal{E}(\vee) := (0, d)$  where
    - $d(b) := (0, k^{\mathbf{T}})$  for every  $b \in \mathcal{B}$  and
    - $d(\mathbf{F}) := (0, \text{id})$
    - $(k^{\mathbf{T}}$  is the constant  $\mathbf{T}$  function)
    - $(\text{id}$  is the identity function from  $\mathcal{D}_o$  to  $\mathcal{D}_o$ )
  - ▶  $\mathcal{E}(\Pi^\alpha) := (0, \pi^\alpha)$  where for each  $(i, f) \in \mathcal{D}_{o\alpha}$ ,  $\pi^\alpha((i, f)) := \mathbf{T}$  if  $f(a) \in \mathcal{B}$  for all  $a \in \mathcal{D}_\alpha$  and  $\pi^\alpha(i, f) := \mathbf{F}$  otherwise

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Easy to check that  $(\mathcal{D}, @)$  is an applicative structure:
- Evaluation function defined by induction on terms
  - ▶  $\mathcal{E}(\neg) := (0, n)$  where  $n(b) := \mathbf{F}$  for every  $b \in \mathcal{B}$  and  $n(\mathbf{F}) := \mathbf{T}$
  - ▶  $\mathcal{E}(\vee) := (0, d)$  where
    - $d(b) := (0, k^{\mathbf{T}})$  for every  $b \in \mathcal{B}$  and
    - $d(\mathbf{F}) := (0, \text{id})$
    - $(k^{\mathbf{T}}$  is the constant  $\mathbf{T}$  function)
    - $(\text{id}$  is the identity function from  $\mathcal{D}_o$  to  $\mathcal{D}_o$ )
  - ▶  $\mathcal{E}(\Pi^\alpha) := (0, \pi^\alpha)$  where for each  $(i, f) \in \mathcal{D}_{o\alpha}$ ,  $\pi^\alpha((i, f)) := \mathbf{T}$  if  $f(a) \in \mathcal{B}$  for all  $a \in \mathcal{D}_\alpha$  and  $\pi^\alpha(i, f) := \mathbf{F}$  otherwise
  - ▶  $q^\alpha := (0, q^\alpha) \in \mathcal{D}_{o\alpha\alpha}$  where  $q^\alpha(a) := (0, s^a)$  and  $s^a(b) := \mathbf{T}$  if  $a = b$  and  $s^a(b) := \mathbf{F}$  otherwise



# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Easy to check that  $(\mathcal{D}, @)$  is an applicative structure:
- Evaluation function defined by induction on terms
  - ▶  $\mathcal{E}(\neg) := (0, n)$  where  $n(b) := \mathbf{F}$  for every  $b \in \mathcal{B}$  and  $n(\mathbf{F}) := \mathbf{T}$
  - ▶  $\mathcal{E}(\vee) := (0, d)$  where
    - $d(b) := (0, k^T)$  for every  $b \in \mathcal{B}$  and
    - $d(\mathbf{F}) := (0, \text{id})$
    - $(k^T$  is the constant  $\mathbf{T}$  function)
    - $(\text{id}$  is the identity function from  $\mathcal{D}_o$  to  $\mathcal{D}_o$ )
  - ▶  $\mathcal{E}(\Pi^\alpha) := (0, \pi^\alpha)$  where for each  $(i, f) \in \mathcal{D}_{o\alpha}$ ,  $\pi^\alpha((i, f)) := \mathbf{T}$  if  $f(a) \in \mathcal{B}$  for all  $a \in \mathcal{D}_\alpha$  and  $\pi^\alpha(i, f) := \mathbf{F}$  otherwise
  - ▶  $q^\alpha := (0, q^\alpha) \in \mathcal{D}_{o\alpha\alpha}$  where  $q^\alpha(a) := (0, s^a)$  and  $s^a(b) := \mathbf{T}$  if  $a = b$  and  $s^a(b) := \mathbf{F}$  otherwise
  - ▶  $\mathcal{E}(w) \in \mathcal{D}_\alpha$  arbitrary for parameters  $w \in \Sigma_\alpha$ .

# Ex.: Models without $\eta$ and $\mathfrak{f}$



# Ex.: Models without $\eta$ and $\mathfrak{f}$



- ▶ For variables, we define  $\mathcal{E}_\varphi(X) := \varphi(X)$

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- ▶ For variables, we define  $\mathcal{E}_\varphi(X) := \varphi(X)$
- ▶ For application, we define  $\mathcal{E}_\varphi(\mathbf{FA}) := \mathcal{E}_\varphi(\mathbf{F}) @ \mathcal{E}_\varphi(\mathbf{A})$

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- ▶ For variables, we define  $\mathcal{E}_\varphi(X) := \varphi(X)$
- ▶ For application, we define  $\mathcal{E}_\varphi(\mathbf{FA}) := \mathcal{E}_\varphi(\mathbf{F}) @ \mathcal{E}_\varphi(\mathbf{A})$
- ▶ For  $\lambda$ -abstractions, we define  $\mathcal{E}_\varphi(\lambda X_\alpha. \mathbf{B}_\beta) := (0, f)$  where  $f: \mathcal{D}_\alpha \longrightarrow \mathcal{D}_\beta$  is the function such that  $f(a) = \mathcal{E}_{\varphi, [a/X]}(\mathbf{B})$  for all  $a \in \mathcal{D}_\alpha$

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- ▶ For variables, we define  $\mathcal{E}_\varphi(X) := \varphi(X)$
- ▶ For application, we define  $\mathcal{E}_\varphi(\mathbf{FA}) := \mathcal{E}_\varphi(\mathbf{F}) @ \mathcal{E}_\varphi(\mathbf{A})$
- ▶ For  $\lambda$ -abstractions, we define  $\mathcal{E}_\varphi(\lambda X_\alpha. \mathbf{B}_\beta) := (0, f)$  where  $f: \mathcal{D}_\alpha \longrightarrow \mathcal{D}_\beta$  is the function such that  $f(a) = \mathcal{E}_{\varphi, [a/X]}(\mathbf{B})$  for all  $a \in \mathcal{D}_\alpha$
- With some work (which we omit), one can show that this  $\mathcal{E}$  is an evaluation function

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- ▶ For variables, we define  $\mathcal{E}_\varphi(X) := \varphi(X)$
  - ▶ For application, we define  $\mathcal{E}_\varphi(\mathbf{FA}) := \mathcal{E}_\varphi(\mathbf{F}) @ \mathcal{E}_\varphi(\mathbf{A})$
  - ▶ For  $\lambda$ -abstractions, we define  $\mathcal{E}_\varphi(\lambda X_\alpha. \mathbf{B}_\beta) := (0, f)$  where  $f: \mathcal{D}_\alpha \longrightarrow \mathcal{D}_\beta$  is the function such that  $f(a) = \mathcal{E}_{\varphi, [a/X]}(\mathbf{B})$  for all  $a \in \mathcal{D}_\alpha$
- With some work (which we omit), one can show that this  $\mathcal{E}$  is an evaluation function
  - Taking  $v$  to be the function such that  $v(b) := \mathbf{T}$  for every  $b \in \mathcal{B}$  and  $v(\mathbf{F}) := \mathbf{F}$ , one can easily show that this is a valuation

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- ▶ For variables, we define  $\mathcal{E}_\varphi(X) := \varphi(X)$
- ▶ For application, we define  $\mathcal{E}_\varphi(\mathbf{FA}) := \mathcal{E}_\varphi(\mathbf{F}) @ \mathcal{E}_\varphi(\mathbf{A})$
- ▶ For  $\lambda$ -abstractions, we define  $\mathcal{E}_\varphi(\lambda X_\alpha. \mathbf{B}_\beta) := (0, f)$  where  $f: \mathcal{D}_\alpha \longrightarrow \mathcal{D}_\beta$  is the function such that  $f(a) = \mathcal{E}_{\varphi, [a/X]}(\mathbf{B})$  for all  $a \in \mathcal{D}_\alpha$
- With some work (which we omit), one can show that this  $\mathcal{E}$  is an evaluation function
- Taking  $v$  to be the function such that  $v(b) := \mathbf{T}$  for every  $b \in \mathcal{B}$  and  $v(\mathbf{F}) := \mathbf{F}$ , one can easily show that this is a valuation
- Hence,  $\mathcal{M}^\mathcal{B} := (\mathcal{D}, @, \mathcal{E}, v)$  is a  $\Sigma$ -model



# Ex.: Models without $\eta$ and $\mathfrak{f}$



- The objects  $q^\alpha := (0, q^\alpha)$  witness property  $q$  for  $\mathcal{M}^B$

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- The objects  $q^\alpha := (0, q^\alpha)$  witness property  $q$  for  $\mathcal{M}^B$
- The objects  $(1, q^\alpha)$  also witness property  $q$  (so, in the non-functional case such witnesses are not unique)

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- The objects  $q^\alpha := (0, q^\alpha)$  witness property  $q$  for  $\mathcal{M}^\mathcal{B}$
- The objects  $(1, q^\alpha)$  also witness property  $q$  (so, in the non-functional case such witnesses are not unique)
- Hence,  $\mathcal{M}^\mathcal{B} := (\mathcal{D}, @, \mathcal{E}, v)$  is a  $\Sigma$ -model with property  $q$

# Ex.: Models without $\eta$ and $\mathfrak{f}$



- Property  $\mathfrak{f}$  fails for  $\mathcal{M}^B$ , since the applicative structure  $(\mathcal{D}, @)$  is not functional:

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Property  $\mathfrak{f}$  fails for  $\mathcal{M}^B$ , since the applicative structure  $(\mathcal{D}, @)$  is not functional:
  - ▶ Consider  $u: \mathcal{D}_\iota \longrightarrow \mathcal{D}_\iota$ .

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Property  $\mathfrak{f}$  fails for  $\mathcal{M}^{\mathcal{B}}$ , since the applicative structure  $(\mathcal{D}, @)$  is not functional:
  - ▶ Consider  $u: \mathcal{D}_\iota \longrightarrow \mathcal{D}_\iota$ .
  - ▶ For both  $(0, u), (1, u) \in \mathcal{D}_u$  we have

$$(i, u)@* = *$$

although  $(0, u) \neq (1, u)$

# Ex.: Models without $\eta$ and $\mathfrak{f}$



- Does  $\eta$  hold?

# Ex.: Models without $\eta$ and $\mathfrak{f}$



- Does  $\eta$  hold?
- No!



# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Does  $\eta$  hold?
- No!
- Compute, for example,  $\mathcal{E}(\lambda F_{\beta\alpha}.F)$  and  $\mathcal{E}(\lambda F_{\beta\alpha}.\lambda X_{\alpha}.FX)$

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Does  $\eta$  hold?
- No!
- Compute, for example,  $\mathcal{E}(\lambda F_{\beta\alpha}.F)$  and  $\mathcal{E}(\lambda F_{\beta\alpha}.\lambda X_{\alpha}.FX)$ 
  - ▶  $\mathcal{E}(\lambda F_{\beta\alpha}.F) = (0, \text{id})$  where  $\text{id}$  is the identity function from  $\mathcal{D}_{\beta\alpha}$  to  $\mathcal{D}_{\beta\alpha}$

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Does  $\eta$  hold?
- No!
- Compute, for example,  $\mathcal{E}(\lambda F_{\beta\alpha}.F)$  and  $\mathcal{E}(\lambda F_{\beta\alpha}.\lambda X_{\alpha}.FX)$ 
  - ▶  $\mathcal{E}(\lambda F_{\beta\alpha}.F) = (0, \text{id})$  where  $\text{id}$  is the identity function from  $\mathcal{D}_{\beta\alpha}$  to  $\mathcal{D}_{\beta\alpha}$
  - ▶  $\mathcal{E}(\lambda F_{\beta\alpha}.\lambda X_{\alpha}.FX) = (0, p)$  where  $p$  is the function from  $\mathcal{D}_{\beta\alpha}$  to  $\mathcal{D}_{\beta\alpha}$  such that  $p((i, f)) = (0, f)$  for each  $f: \mathcal{D}_{\alpha} \longrightarrow \mathcal{D}_{\beta}$

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Does  $\eta$  hold?
- No!
- Compute, for example,  $\mathcal{E}(\lambda F_{\beta\alpha}.F)$  and  $\mathcal{E}(\lambda F_{\beta\alpha}.\lambda X_{\alpha}.FX)$ 
  - ▶  $\mathcal{E}(\lambda F_{\beta\alpha}.F) = (0, \text{id})$  where  $\text{id}$  is the identity function from  $\mathcal{D}_{\beta\alpha}$  to  $\mathcal{D}_{\beta\alpha}$
  - ▶  $\mathcal{E}(\lambda F_{\beta\alpha}.\lambda X_{\alpha}.FX) = (0, p)$  where  $p$  is the function from  $\mathcal{D}_{\beta\alpha}$  to  $\mathcal{D}_{\beta\alpha}$  such that  $p((i, f)) = (0, f)$  for each  $f: \mathcal{D}_{\alpha} \longrightarrow \mathcal{D}_{\beta}$
- Hence  $\mathcal{E}(\lambda F_{\beta\alpha}.F) \neq \mathcal{E}(\lambda F_{\beta\alpha}.\lambda X_{\alpha}.FX)$

# Ex.: Models without $\eta$ and $\mathfrak{f}$



- Does  $\xi$  hold?

# Ex.: Models without $\eta$ and $\mathfrak{f}$



- Does  $\xi$  hold?
- Yes!

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Does  $\xi$  hold?
- Yes!
- If

$$\mathcal{E}_{\varphi, [a/X]}(\mathbf{M}) = \mathcal{E}_{\varphi, [a/X]}(\mathbf{N})$$

for every  $a \in \mathcal{D}_{\alpha}$ , then

$$\mathcal{E}_{\varphi}(\lambda X_{\alpha}.\mathbf{M}) = (0, f) = \mathcal{E}_{\varphi}(\lambda X.\mathbf{N})$$

where  $f(a) = \mathcal{E}_{\varphi, [a/X]}(\mathbf{M}) = \mathcal{E}_{\varphi, [a/X]}(\mathbf{N})$  for every  $a \in \mathcal{D}_{\alpha}$ .

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- If  $\mathcal{B} = \{\mathbf{T}\}$ , then the model  $\mathcal{M}^{\beta\xi\mathfrak{b}} := \mathcal{M}^{\{\mathbf{T}\}}$  satisfies property  $\mathfrak{b}$ .



# Ex.: Models without $\eta$ and $\mathfrak{f}$

- If  $\mathcal{B} = \{\mathsf{T}\}$ , then the model  $\mathcal{M}^{\beta\xi\mathfrak{b}} := \mathcal{M}^{\{\mathsf{T}\}}$  satisfies property  $\mathfrak{b}$ .
- So, we know  $\mathcal{M}^{\beta\xi\mathfrak{b}} \in \mathfrak{M}_{\beta\xi\mathfrak{b}}(\Sigma) \setminus \mathfrak{M}_{\beta\mathfrak{fb}}(\Sigma)$ .


# Ex.: Models without $\eta$ and $\mathfrak{f}$

- If  $\mathcal{B} = \{\mathsf{T}\}$ , then the model  $\mathcal{M}^{\beta\xi\mathfrak{b}} := \mathcal{M}^{\{\mathsf{T}\}}$  satisfies property  $\mathfrak{b}$ .
- So, we know  $\mathcal{M}^{\beta\xi\mathfrak{b}} \in \mathfrak{M}_{\beta\xi\mathfrak{b}}(\Sigma) \setminus \mathfrak{M}_{\beta\mathfrak{f}\mathfrak{b}}(\Sigma)$ .
- On the other hand, if  $\mathfrak{b}$  is any value with  $\mathfrak{b} \notin \{\mathsf{T}, \mathsf{F}\}$ , and  $\mathcal{B} = \{\mathsf{T}, \mathfrak{b}\}$ , then the model  $\mathcal{M}^{\beta\xi} := \mathcal{M}^{\{\mathsf{T}, \mathfrak{b}\}}$  does not satisfy property  $\mathfrak{b}$ .


# Ex.: Models without $\eta$ and $\mathfrak{f}$

- If  $\mathcal{B} = \{\mathbf{T}\}$ , then the model  $\mathcal{M}^{\beta\xi\mathbf{b}} := \mathcal{M}^{\{\mathbf{T}\}}$  satisfies property  $\mathbf{b}$ .
- So, we know  $\mathcal{M}^{\beta\xi\mathbf{b}} \in \mathfrak{M}_{\beta\xi\mathbf{b}}(\Sigma) \setminus \mathfrak{M}_{\beta\mathfrak{f}\mathbf{b}}(\Sigma)$ .
- On the other hand, if  $\mathbf{b}$  is any value with  $\mathbf{b} \notin \{\mathbf{T}, \mathbf{F}\}$ , and  $\mathcal{B} = \{\mathbf{T}, \mathbf{b}\}$ , then the model  $\mathcal{M}^{\beta\xi} := \mathcal{M}^{\{\mathbf{T}, \mathbf{b}\}}$  does not satisfy property  $\mathbf{b}$ .
- In this case, we know  $\mathcal{M}^{\beta\xi} \in \mathfrak{M}_{\beta\xi}(\Sigma) \setminus (\mathfrak{M}_{\beta\mathfrak{f}}(\Sigma) \cup \mathfrak{M}_{\beta\xi\mathbf{b}}(\Sigma))$ .

# Ex.: Models without $\eta$ and $f$



# Ex.: Models without $\eta$ and $f$



# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Let  $\mathcal{M}^{\mathcal{B}}$  be the  $\Sigma$ -model  $(\mathcal{D}, @, \mathcal{E}, v)$  as constructed before

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Let  $\mathcal{M}^{\mathcal{B}}$  be the  $\Sigma$ -model  $(\mathcal{D}, @, \mathcal{E}, v)$  as constructed before
- Define an alternative evaluation function  $\mathcal{E}'$  by induction:

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Let  $\mathcal{M}^{\mathcal{B}}$  be the  $\Sigma$ -model  $(\mathcal{D}, @, \mathcal{E}, v)$  as constructed before
- Define an alternative evaluation function  $\mathcal{E}'$  by induction:
  - ▶ For all  $w \in \Sigma$ , let  $\mathcal{E}'(w) := \mathcal{E}(w)$



# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Let  $\mathcal{M}^{\mathcal{B}}$  be the  $\Sigma$ -model  $(\mathcal{D}, @, \mathcal{E}, v)$  as constructed before
- Define an alternative evaluation function  $\mathcal{E}'$  by induction:
  - ▶ For all  $w \in \Sigma$ , let  $\mathcal{E}'(w) := \mathcal{E}(w)$
  - ▶ For variables we define  $\mathcal{E}'_{\varphi}(X) := \varphi(X)$

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Let  $\mathcal{M}^{\mathcal{B}}$  be the  $\Sigma$ -model  $(\mathcal{D}, @, \mathcal{E}, v)$  as constructed before
- Define an alternative evaluation function  $\mathcal{E}'$  by induction:
  - ▶ For all  $w \in \Sigma$ , let  $\mathcal{E}'(w) := \mathcal{E}(w)$
  - ▶ For variables we define  $\mathcal{E}'_{\varphi}(X) := \varphi(X)$
  - ▶ We must define  $\mathcal{E}'_{\varphi}(\mathbf{FA}) := \mathcal{E}'_{\varphi}(\mathbf{F}) @ \mathcal{E}'_{\varphi}(\mathbf{A})$

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Let  $\mathcal{M}^{\mathcal{B}}$  be the  $\Sigma$ -model  $(\mathcal{D}, @, \mathcal{E}, v)$  as constructed before
- Define an alternative evaluation function  $\mathcal{E}'$  by induction:
  - ▶ For all  $w \in \Sigma$ , let  $\mathcal{E}'(w) := \mathcal{E}(w)$
  - ▶ For variables we define  $\mathcal{E}'_{\varphi}(X) := \varphi(X)$
  - ▶ We must define  $\mathcal{E}'_{\varphi}(\mathbf{FA}) := \mathcal{E}'_{\varphi}(\mathbf{F}) @ \mathcal{E}'_{\varphi}(\mathbf{A})$
  - ▶ We choose  $\mathcal{E}'_{\varphi}(\lambda X_{\alpha}.\mathbf{B}_{\beta}) := (1, f)$  where  $f: \mathcal{D}_{\alpha} \longrightarrow \mathcal{D}_{\beta}$  is the function such that  $f(a) = \mathcal{E}_{\varphi, [a/X]}(\mathbf{B})$  for all  $a \in \mathcal{D}_{\alpha}$

# Ex.: Models without $\eta$ and $\mathfrak{f}$

- Let  $\mathcal{M}^B$  be the  $\Sigma$ -model  $(\mathcal{D}, @, \mathcal{E}, v)$  as constructed before
- Define an alternative evaluation function  $\mathcal{E}'$  by induction:
  - ▶ For all  $w \in \Sigma$ , let  $\mathcal{E}'(w) := \mathcal{E}(w)$
  - ▶ For variables we define  $\mathcal{E}'_{\varphi}(X) := \varphi(X)$
  - ▶ We must define  $\mathcal{E}'_{\varphi}(\mathbf{FA}) := \mathcal{E}'_{\varphi}(\mathbf{F}) @ \mathcal{E}'_{\varphi}(\mathbf{A})$
  - ▶ We choose  $\mathcal{E}'_{\varphi}(\lambda X_{\alpha}. \mathbf{B}_{\beta}) := (1, f)$  where  $f: \mathcal{D}_{\alpha} \longrightarrow \mathcal{D}_{\beta}$  is the function such that  $f(a) = \mathcal{E}_{\varphi, [a/X]}(\mathbf{B})$  for all  $a \in \mathcal{D}_{\alpha}$
- $\mathcal{E}$  and  $\mathcal{E}'$  agree on all constants, they are different though:

$$\mathcal{E}(\lambda X_{\iota}. X) = (0, \text{id}) \neq (1, \text{id}) = \mathcal{E}'(\lambda X_{\iota}. X)$$

where  $\text{id} : \mathcal{D}_{\iota} \longrightarrow \mathcal{D}_{\iota}$  is the identity function

# Ex.: Models without $\eta$ and $\mathfrak{f}$


- Let  $\mathcal{M}^B$  be the  $\Sigma$ -model  $(\mathcal{D}, @, \mathcal{E}, v)$  as constructed before
- Define an alternative evaluation function  $\mathcal{E}'$  by induction:
  - ▶ For all  $w \in \Sigma$ , let  $\mathcal{E}'(w) := \mathcal{E}(w)$
  - ▶ For variables we define  $\mathcal{E}'_{\varphi}(X) := \varphi(X)$
  - ▶ We must define  $\mathcal{E}'_{\varphi}(\mathbf{FA}) := \mathcal{E}'_{\varphi}(\mathbf{F}) @ \mathcal{E}'_{\varphi}(\mathbf{A})$
  - ▶ We choose  $\mathcal{E}'_{\varphi}(\lambda X_{\alpha}. \mathbf{B}_{\beta}) := (1, f)$  where  $f: \mathcal{D}_{\alpha} \longrightarrow \mathcal{D}_{\beta}$  is the function such that  $f(a) = \mathcal{E}_{\varphi, [a/X]}(\mathbf{B})$  for all  $a \in \mathcal{D}_{\alpha}$
- $\mathcal{E}$  and  $\mathcal{E}'$  agree on all constants, they are different though:

$$\mathcal{E}(\lambda X_{\iota}. X) = (0, \text{id}) \neq (1, \text{id}) = \mathcal{E}'(\lambda X_{\iota}. X)$$


where  $\text{id} : \mathcal{D}_{\iota} \longrightarrow \mathcal{D}_{\iota}$  is the identity function

- Thus, in non-functional models evaluation functions are not uniquely determined by their values on constants

# Ex.: Models without $\xi$




# Ex.: Models without $\xi$



Not here!


See [JSL-04]

# Ex.: Models without $\xi$





# Ex.: Models without $\xi$



Not here!

See [JSL-04]