

Examples of Diophantine relations:

1. $a \equiv b \pmod{c}$: $\exists x: a = cx + b$ or $b = cx + a$. (over \mathbb{N} !)
2. $a \geq b$: $a = b + x$ (\exists is always implicit)
3. $a > b$: (Exercise)
4. $a = b \pmod{c}$: $a \equiv b \pmod{c}$ and $0 \leq a < c$.
5. $\{(a, b, c) \mid a = b^c\}$?
6. $\{2^k\}$? Tarski believed not Diophantine

$G_b(0) = 0, G_b(1) = 1, \begin{cases} G_n(n+1) = b \cdot G_b(n) - G_b(n-1) \\ G_b(n-1) + G_n(n+1) = b \cdot G_b(n) \end{cases}$ close to the recursion for b^{n-1}
 \Leftrightarrow symmetric between forward and backward

$b = 4$: $[\dots, -15, -4, -1,] 0, 1, 4, 15, 56, \dots$

$b = 3$: $0, 1, 3, 8, 21, \dots$

$b = 2$: $0, 1, 2, 3, 4, 5, \dots$ (useless?)

WE SHOW: $a = G_b(c)$ for (fixed) $b \geq 3$ is Diophantine. ($b \geq 4$ simplifies some arguments.)

Missing link! [Yuri Matiyasevich 1970, Julia Robinson, Martin Davis, Hilary Putnam 1961]

The points $\begin{pmatrix} x_n \\ y_n \end{pmatrix} = \begin{pmatrix} G_b(n+1) \\ G_b(n) \end{pmatrix}$ lie on the hyperbola $h_b(x, y) := x^2 - bxy + y^2 - 1 = 0$. (\rightarrow picture)

Lemma 1. *The only integer solutions of $h_b(x, y) = 0$ with $x > y \geq 0$ are those points.*

Proof: The hyperbola is invariant under the shift $\begin{pmatrix} G_b(n) \\ G_b(n-1) \end{pmatrix} \leftrightarrow \begin{pmatrix} G_b(n+1) \\ G_b(n) \end{pmatrix}$: $\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} bx-y \\ x \end{pmatrix}$ or $\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} y \\ by-x \end{pmatrix}$, and the shift preserves $y < x$. \square

Now we can generate $\{G_b(n)\} = \{x \mid \exists y: h_b(x, y) = 0\}$ but we don't know n .

Lemma 2. $b \equiv b' \pmod{u} \implies G_b(n) \equiv G_{b'}(n) \pmod{u}$ (Induction. Easy.) \square

IDEA: Choose two appropriate moduli M and m to coordinate $G_b(n)$ with n :

$$\begin{array}{ll} G_w(0), G_w(1), \dots, G_w(n), \dots \pmod{M} & w \equiv b \pmod{M} \quad \boxed{\rightarrow G_b(n)} \\ G_w(0), G_w(1), \dots, G_w(n), \dots \pmod{m} & w \equiv 2 \pmod{m} \quad \boxed{\rightarrow G_2(n) = n} \end{array}$$

As long as n is small and $G_w(n) \leq M$, we have $G_w(n) \pmod{M} = G_b(n)$ (and $G_w(n) \pmod{m} = n$), but for larger n , $G_w(n) \pmod{M}$ gets out of control, and there will be extra solutions. (\rightarrow picture)

- Make $G_w(n) \pmod{M}$ mirror-symmetric after reaching a peak at $G_w(p) = G_b(p)$: (\rightarrow picture)
 $G_b(p-1) \equiv G_b(p+1) \pmod{M} \implies M := G_b(p+1) - G_b(p-1)$ ($p = \text{peak} = \text{period}$)
- Avoid “negative” values by using the absmod operation instead of mod. (\rightarrow picture)
 $x \text{ absmod } M = a \iff x = qM \pm a$ and $0 \leq a \leq M/2$.
- The period m of “ $G_w(n) \text{ absmod } m = n$ ” should divide the period $2p$ of “ $G_w(n) \text{ absmod } M$ ”: $m \mid p$.
- Choose m (and M) larger than twice the (supposed) value a of $G_b(c)$, so that absmod does no harm.

1. $m > 2a$.
2. p should be a multiple of m
3. $M := G_b(p+1) - G_b(p-1)$
4. Choose $w > 2$ with $w \equiv b \pmod{M}$
 $w \equiv 2 \pmod{m}$
5. $h_w(x, y) = 0$ [$\implies x = G_w(n)$ for some n]
6. $a = x \text{ absmod } M$ [$a = G_b(n)$]
 $c = x \text{ absmod } m$ [$c = G_2(n) = n$]

Lemma 3. $G_b(k)^2 \mid G_b(p) \implies G_b(k) \mid p$

(Cf. Fibonacci numbers: $F_k \mid F_p \iff k \mid p$.)

Application: Choose m of the form $m = G_b(k)$ for some k , by requiring $h_b(m, m') = 0$.

Then $m^2 \mid G_b(p) \implies m \mid p$.

Implementation of Conditions 2 and 3.

$$\begin{aligned} \boxed{h_b(r, s) = 0, r < s} \quad & G_b(p-1) = r, \text{ for some } p \\ & G_b(p) = s \\ & G_b(p+1) = bs - r \end{aligned}$$

$$\boxed{M = (bs - r) - r} \quad [= G_b(p+1) - G_b(p-1). \text{ Also } G_b(p) < M/2.]$$

$$\boxed{m^2 \mid s} \quad [\implies m \mid p.]$$

The conditions are enough to ensure that every solution (a, c) satisfies $a = G_b(c)$. (The condition $a < m/2$ cuts off extra solutions.)

Converse direction:

We need to show that m, M with $\gcd(m, M) = 1$ exist (then w satisfying (4.) exists, by the Chinese Remainder Theorem), and that $m^2 \mid s$ can be fulfilled.

- Choose $m = G_b(k)$ odd and set $s = G_b(p)$ for $p = k \cdot m$:

Then it can be shown that $\gcd(m, M) = 1$ and $m^2 \mid s$.

Getting to the relation $a = b^c$:

$$(b-1)^n \leq G_b(n+1) \leq b^n$$

$$b^c = \lim_{x \rightarrow \infty} \frac{G_{bx+4}(c+1)}{G_x(c+1)} \approx \frac{(bx \pm \text{const})^c}{(x \pm \text{const})^c} \rightarrow b^c, \text{ for all } b, c \geq 0$$

$$b^c = \left\lfloor \frac{G_{bx+4}(c+1)}{G_x(c+1)} \right\rfloor \text{ for } x > 16(c+1)G_{b+4}(c+1).$$

(The “+4” term ensures that this works even for $b = 0$.)