



Completeness (of  $\mathcal{N}_*$ )

# Completeness (of $\mathcal{NK}_*$ )



- How to prove completeness?

# Completeness (of $\mathcal{NK}_*$ )



- How to prove completeness?
- Completeness can be proven rather easily for propositional logic calculi.

# Completeness (of $\mathcal{NK}_*$ )



- How to prove completeness?
- Completeness can be proven rather easily for propositional logic calculi.
- For first-order and especially higher-order logic completeness proofs become increasingly difficult and technical.

# Completeness (of $\mathcal{NK}_*$ )



- How to prove completeness?
- Completeness can be proven rather easily for propositional logic calculi.
- For first-order and especially higher-order logic completeness proofs become increasingly difficult and technical.
- Here we will introduce a **strong proof tool** that uniformly supports completeness proofs (and many other things): **abstract consistency**.

# Completeness (of $\mathcal{NK}_*$ )



- How to prove completeness?
- Completeness can be proven rather easily for propositional logic calculi.
- For first-order and especially higher-order logic completeness proofs become increasingly difficult and technical.
- Here we will introduce a **strong proof tool** that uniformly supports completeness proofs (and many other things): **abstract consistency**.
- This proof tool is based on a strong theorem which connects syntax and semantics: **model existence theorem**.



Abstract Consistency

# Abstract Consistency: History



- Technique was developed for first-order logic by Jaakko Hintikka and Raymond Smullyan [Hintikka55,Smullyan63,Smullyan68]. It is well explained in Fitting's textbook [Fitting96].



# Abstract Consistency: History



- Technique was developed for first-order logic by Jaakko Hintikka and Raymond Smullyan [Hintikka55,Smullyan63,Smullyan68]. It is well explained in Fitting's textbook [Fitting96].
- The technique has been (partly) extended to higher-order logic by Peter Andrews' in [Andrews71]; Peter Andrews only achieves a generalization for his rather weak semantical  $v$ -complexes (corresponding to our  $\mathfrak{M}_\beta(\Sigma)$ ) and not, for instance, for Henkin Semantics. This extension is well explained in Peter Andrews's textbook [Andrews02].

# Abstract Consistency: History



- Technique was developed for first-order logic by Jaakko Hintikka and Raymond Smullyan [Hintikka55,Smullyan63,Smullyan68]. It is well explained in Fitting's textbook [Fitting96].
- The technique has been (partly) extended to higher-order logic by Peter Andrews' in [Andrews71]; Peter Andrews only achieves a generalization for his rather weak semantical  $v$ -complexes (corresponding to our  $\mathfrak{M}_\beta(\Sigma)$ ) and not, for instance, for Henkin Semantics. This extension is well explained in Peter Andrews's textbook [Andrews02].
- The technique has been extended to our landscape of HOL model classes in [Benzmueller-PhD-99,JSL04].

# Abstract Consistency: Idea



- A **model existence theorem** for a logical system (i.e., a logical language  $L$  together with a consequence relation  $\models$ ) is a theorem of the form:

# Abstract Consistency: Idea



- A **model existence theorem** for a logical system (i.e., a logical language  $L$  together with a consequence relation  $\models$ ) is a theorem of the form:

*If a set of sentences  $\Phi$  of  $L$  is a member of an (saturated) abstract consistency class  $\Gamma$ , then there exists a model for  $\Phi$ .*

# Abstract Consistency: Idea

- Employing the model existence theorem we can prove completeness of a calculus  $\mathcal{C}$  (i.e., the derivability rel.  $\vdash_{\mathcal{C}}$ ) by

# Abstract Consistency: Idea

- Employing the model existence theorem we can prove completeness of a calculus  $\mathcal{C}$  (i.e., the derivability rel.  $\vdash_{\mathcal{C}}$ ) by proving that the class  $\Gamma$  of sets of sentences  $\Phi$  that are  $\mathcal{C}$ -consistent (i.e., cannot be refuted in  $\mathcal{C}$ ) is an (saturated) abstract consistency class.

# Abstract Consistency: Idea

- Employing the model existence theorem we can prove completeness of a calculus  $\mathcal{C}$  (i.e., the derivability rel.  $\vdash_{\mathcal{C}}$ ) by proving that the class  $\Gamma$  of sets of sentences  $\Phi$  that are  $\mathcal{C}$ -consistent (i.e., cannot be refuted in  $\mathcal{C}$ ) is an (saturated) abstract consistency class.
- Why does this work?

# Abstract Consistency: Idea

- Employing the model existence theorem we can prove completeness of a calculus  $\mathcal{C}$  (i.e., the derivability rel.  $\vdash_{\mathcal{C}}$ ) by proving that the class  $\Gamma$  of sets of sentences  $\Phi$  that are  $\mathcal{C}$ -consistent (i.e., cannot be refuted in  $\mathcal{C}$ ) is an (saturated) abstract consistency class.
- Why does this work?
  - ▶ The model existence theorem tells us that  $\mathcal{C}$ -consistent sets of sentences are satisfiable.



# Abstract Consistency: Idea

- Employing the model existence theorem we can prove completeness of a calculus  $\mathcal{C}$  (i.e., the derivability rel.  $\vdash_{\mathcal{C}}$ ) by proving that the class  $\Gamma$  of sets of sentences  $\Phi$  that are  $\mathcal{C}$ -consistent (i.e., cannot be refuted in  $\mathcal{C}$ ) is an (saturated) abstract consistency class.
- Why does this work?
  - ▶ The model existence theorem tells us that  $\mathcal{C}$ -consistent sets of sentences are satisfiable.
  - ▶ Now we assume that a sentence  $A$  is valid, so  $\neg A$  does not have a model and is therefore  $\mathcal{C}$ -inconsistent.

# Abstract Consistency: Idea

- Employing the model existence theorem we can prove completeness of a calculus  $\mathcal{C}$  (i.e., the derivability rel.  $\vdash_{\mathcal{C}}$ ) by proving that the class  $\Gamma$  of sets of sentences  $\Phi$  that are  $\mathcal{C}$ -consistent (i.e., cannot be refuted in  $\mathcal{C}$ ) is an (saturated) abstract consistency class.
- Why does this work?
  - ▶ The model existence theorem tells us that  $\mathcal{C}$ -consistent sets of sentences are satisfiable.
  - ▶ Now we assume that a sentence  $A$  is valid, so  $\neg A$  does not have a model and is therefore  $\mathcal{C}$ -inconsistent.
  - ▶ Hence,  $\neg A$  is refutable in  $\mathcal{C}$ .

# Abstract Consistency: Idea

- Employing the model existence theorem we can prove completeness of a calculus  $\mathcal{C}$  (i.e., the derivability rel.  $\vdash_{\mathcal{C}}$ ) by proving that the class  $\Gamma$  of sets of sentences  $\Phi$  that are  $\mathcal{C}$ -consistent (i.e., cannot be refuted in  $\mathcal{C}$ ) is an (saturated) abstract consistency class.
- Why does this work?
  - ▶ The model existence theorem tells us that  $\mathcal{C}$ -consistent sets of sentences are satisfiable.
  - ▶ Now we assume that a sentence  $A$  is valid, so  $\neg A$  does not have a model and is therefore  $\mathcal{C}$ -inconsistent.
  - ▶ Hence,  $\neg A$  is refutable in  $\mathcal{C}$ .
  - ▶ This shows refutation completeness of  $\mathcal{C}$ .

# Abstract Consistency: Idea

- Employing the model existence theorem we can prove completeness of a calculus  $\mathcal{C}$  (i.e., the derivability rel.  $\vdash_{\mathcal{C}}$ ) by proving that the class  $\Gamma$  of sets of sentences  $\Phi$  that are  $\mathcal{C}$ -consistent (i.e., cannot be refuted in  $\mathcal{C}$ ) is an (saturated) abstract consistency class.
- Why does this work?
  - ▶ The model existence theorem tells us that  $\mathcal{C}$ -consistent sets of sentences are satisfiable.
  - ▶ Now we assume that a sentence  $A$  is valid, so  $\neg A$  does not have a model and is therefore  $\mathcal{C}$ -inconsistent.
  - ▶ Hence,  $\neg A$  is refutable in  $\mathcal{C}$ .
  - ▶ This shows refutation completeness of  $\mathcal{C}$ .
  - ▶ For many calculi  $\mathcal{C}$ , this also shows  $A$  is provable, thus establishing completeness of  $\mathcal{C}$ .

# Def.: Closed under Subsets / Compact



Let  $\mathcal{C}$  be a class of sets then  $\mathcal{C}$  is called **closed under subset** if for all sets  $S$  and  $T$  it holds that

# Def.: Closed under Subsets / Compact



Let  $\mathcal{C}$  be a class of sets then  $\mathcal{C}$  is called **closed under subset** if for all sets  $S$  and  $T$  it holds that

from  $S \subseteq T$  and  $T \in \mathcal{C}$  it follows that  $S \in \mathcal{C}$ .

# Def.: Closed under Subsets / Compact



Let  $C$  be a class of sets then  $C$  is called **closed under subset** if for all sets  $S$  and  $T$  it holds that

from  $S \subseteq T$  and  $T \in C$  it follows that  $S \in C$ .

Let  $C$  be a class of sets.  $C$  is called **compact** or **of finite character** if and only if for every set  $S$  holds:

# Def.: Closed under Subsets / Compact



Let  $C$  be a class of sets then  $C$  is called **closed under subset** if for all sets  $S$  and  $T$  it holds that

from  $S \subseteq T$  and  $T \in C$  it follows that  $S \in C$ .

Let  $C$  be a class of sets.  $C$  is called **compact** or **of finite character** if and only if for every set  $S$  holds:

$S \in C$  if and only if every finite subset of  $S$  is a member of  $C$ .



# Ex.: Closed under Subsets / Compact



- not closed under subsets:  $\{\{\neg(A \vee B), \neg A, C\}, \{\neg A\}\}$

# Ex.: Closed under Subsets / Compact



- not closed under subsets:  $\{\{\neg(A \vee B), \neg A, C\}, \{\neg A\}\}$
- closed under subsets:  $\{\{\neg(A \vee B), \neg A, C\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B), C\}, \{\neg A, C\}, \{\neg(A \vee B)\}, \{\neg A\}, \{C\}, \{\}\}$

# Ex.: Closed under Subsets / Compact



- not closed under subsets:  $\{\{\neg(A \vee B), \neg A, C\}, \{\neg A\}\}$
- closed under subsets:  $\{\{\neg(A \vee B), \neg A, C\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B), C\}, \{\neg A, C\}, \{\neg(A \vee B)\}, \{\neg A\}, \{C\}, \{\}\}$
- We define two classes of sets

# Ex.: Closed under Subsets / Compact



- not closed under subsets:  $\{\{\neg(A \vee B), \neg A, C\}, \{\neg A\}\}$
- closed under subsets:  $\{\{\neg(A \vee B), \neg A, C\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B), C\}, \{\neg A, C\}, \{\neg(A \vee B)\}, \{\neg A\}, \{C\}, \{\}\}$
- We define two classes of sets
  - ▶  $C := \{\varphi \mid \varphi \text{ is finite subset of } \mathbb{N}\}$

# Ex.: Closed under Subsets / Compact



- not closed under subsets:  $\{\{\neg(A \vee B), \neg A, C\}, \{\neg A\}\}$
- closed under subsets:  $\{\{\neg(A \vee B), \neg A, C\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B), C\}, \{\neg A, C\}, \{\neg(A \vee B)\}, \{\neg A\}, \{C\}, \{\}\}$
- We define two classes of sets
  - ▶  $C := \{\varphi \mid \varphi \text{ is finite subset of } \mathbb{N}\}$
  - ▶  $D := 2^{\mathbb{N}}$

# Ex.: Closed under Subsets / Compact



- not closed under subsets:  $\{\{\neg(A \vee B), \neg A, C\}, \{\neg A\}\}$
- closed under subsets:  $\{\{\neg(A \vee B), \neg A, C\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B), C\}, \{\neg A, C\}, \{\neg(A \vee B)\}, \{\neg A\}, \{C\}, \{\}\}$
- We define two classes of sets
  - ▶  $C := \{\varphi \mid \varphi \text{ is finite subset of } \mathbb{N}\}$
  - ▶  $D := 2^{\mathbb{N}}$
  - ▶  $C$  is closed under subsets but **not** compact.

# Ex.: Closed under Subsets / Compact



- not closed under subsets:  $\{\{\neg(A \vee B), \neg A, C\}, \{\neg A\}\}$
- closed under subsets:  $\{\{\neg(A \vee B), \neg A, C\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B), C\}, \{\neg A, C\}, \{\neg(A \vee B)\}, \{\neg A\}, \{C\}, \{\}\}$
- We define two classes of sets
  - ▶  $C := \{\varphi \mid \varphi \text{ is finite subset of } \mathbb{N}\}$
  - ▶  $D := 2^{\mathbb{N}}$
  - ▶  $C$  is closed under subsets but **not** compact.
  - ▶  $D$  is closed under subsets **and** compact.

# Lemma: Closed under Subsets / Compact



Lemma:

If  $C$  is compact then  $C$  is closed under subsets.



# Lemma: Closed under Subsets / Compact



Lemma:

If  $C$  is compact then  $C$  is closed under subsets.

Proof:

# Lemma: Closed under Subsets / Compact



Lemma:

If  $C$  is compact then  $C$  is closed under subsets.

Proof:

Let  $T \in C$  and  $S \subseteq T$ .

# Lemma: Closed under Subsets / Compact



Lemma:

If  $C$  is compact then  $C$  is closed under subsets.

Proof:

Let  $T \in C$  and  $S \subseteq T$ .

We have to show that  $S \in C$ .

# Lemma: Closed under Subsets / Compact



Lemma:

If  $C$  is compact then  $C$  is closed under subsets.

Proof:

Let  $T \in C$  and  $S \subseteq T$ .

We have to show that  $S \in C$ .

Every finite subset  $A$  of  $S$  is also a finite subset of  $T$ .

# Lemma: Closed under Subsets / Compact



Lemma:

If  $C$  is compact then  $C$  is closed under subsets.

Proof:

Let  $T \in C$  and  $S \subseteq T$ .

We have to show that  $S \in C$ .

Every finite subset  $A$  of  $S$  is also a finite subset of  $T$ .

Since  $C$  is compact and  $T \in C$  we get that all  $A \in C$ .

# Lemma: Closed under Subsets / Compact



Lemma:

If  $C$  is compact then  $C$  is closed under subsets.

Proof:

Let  $T \in C$  and  $S \subseteq T$ .

We have to show that  $S \in C$ .

Every finite subset  $A$  of  $S$  is also a finite subset of  $T$ .

Since  $C$  is compact and  $T \in C$  we get that all  $A \in C$ .

Thus,  $S \in C$  by compactness.

# Def.: Sufficiently $\Sigma$ -Pure



We introduce a technical side-condition that ensures that we always have enough witness constants.

# Def.: Sufficiently $\Sigma$ -Pure

We introduce a technical side-condition that ensures that we always have enough witness constants.

Let  $\Sigma$  be a signature and  $\Phi$  be a set of  $\Sigma$ -sentences.  $\Phi$  is called **sufficiently  $\Sigma$ -pure** if for each type  $\alpha$  there is a set  $\mathcal{P}_\alpha \subseteq \Sigma_\alpha$  of parameters with equal cardinality to  $wff_\alpha(\Sigma)$ , such that the elements of  $\mathcal{P}_\alpha$  do not occur in the sentences of  $\Phi$ .



# Def.: Sufficiently $\Sigma$ -Pure

We introduce a technical side-condition that ensures that we always have enough witness constants.

Let  $\Sigma$  be a signature and  $\Phi$  be a set of  $\Sigma$ -sentences.  $\Phi$  is called **sufficiently  $\Sigma$ -pure** if for each type  $\alpha$  there is a set  $\mathcal{P}_\alpha \subseteq \Sigma_\alpha$  of parameters with equal cardinality to  $wff_\alpha(\Sigma)$ , such that the elements of  $\mathcal{P}_\alpha$  do not occur in the sentences of  $\Phi$ .

This can be obtained in practice by enriching the signature with spurious parameters.

# Abstract Consistency: Conventions



Remember the conventions for this part of the lecture:

- signature  $\Sigma$  contains only the logical constants  $\neg, \vee, \Pi^\alpha$  unless stated otherwise

# Abstract Consistency: Conventions



Remember the conventions for this part of the lecture:

- signature  $\Sigma$  contains only the logical constants  $\neg, \vee, \Pi^\alpha$  unless stated otherwise
- as a matter of convenience we will write  $\varphi * A$  for  $\varphi \cup \{A\}$ .

# Def.: Abstract Consistency Properties



Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $A, B \in \text{cwff}_o(\Sigma)$ ,  $F \in \text{cwff}_{\alpha \rightarrow o}(\Sigma)$  are arbitrary):

# Def.: Abstract Consistency Properties

Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $A, B \in \text{cwff}_o(\Sigma)$ ,  $F \in \text{cwff}_{\alpha \rightarrow o}(\Sigma)$  are arbitrary):

$\nabla_c$  If  $A$  is atomic, then  $A \notin \Phi$  or  $\neg A \notin \Phi$ .

# Def.: Abstract Consistency Properties

Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $A, B \in \text{cwff}_o(\Sigma)$ ,  $F \in \text{cwff}_{\alpha \rightarrow o}(\Sigma)$  are arbitrary):

$\nabla_c$  If  $A$  is atomic, then  $A \notin \Phi$  or  $\neg A \notin \Phi$ .

$\nabla_\neg$  If  $\neg\neg A \in \Phi$ , then  $\Phi * A \in \Gamma_\Sigma$ .

# Def.: Abstract Consistency Properties

Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{A}, \mathbf{B} \in \text{cwff}_o(\Sigma)$ ,  $\mathbf{F} \in \text{cwff}_{\alpha \rightarrow o}(\Sigma)$  are arbitrary):

$\nabla_c$  If  $\mathbf{A}$  is atomic, then  $\mathbf{A} \notin \Phi$  or  $\neg \mathbf{A} \notin \Phi$ .

$\nabla_\neg$  If  $\neg \neg \mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{A} \in \Gamma_\Sigma$ .

$\nabla_\beta$  If  $\mathbf{A} =_\beta \mathbf{B}$  and  $\mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .

# Def.: Abstract Consistency Properties

Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{A}, \mathbf{B} \in \text{cwff}_o(\Sigma)$ ,  $\mathbf{F} \in \text{cwff}_{\alpha \rightarrow o}(\Sigma)$  are arbitrary):

$\nabla_c$  If  $\mathbf{A}$  is atomic, then  $\mathbf{A} \notin \Phi$  or  $\neg \mathbf{A} \notin \Phi$ .

$\nabla_\neg$  If  $\neg \neg \mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{A} \in \Gamma_\Sigma$ .

$\nabla_\beta$  If  $\mathbf{A} =_\beta \mathbf{B}$  and  $\mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_\vee$  If  $\mathbf{A} \vee \mathbf{B} \in \Phi$ , then  $\Phi * \mathbf{A} \in \Gamma_\Sigma$  or  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .



# Def.: Abstract Consistency Properties

Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{A}, \mathbf{B} \in \text{cwff}_o(\Sigma)$ ,  $\mathbf{F} \in \text{cwff}_{\alpha \rightarrow o}(\Sigma)$  are arbitrary):

$\nabla_c$  If  $\mathbf{A}$  is atomic, then  $\mathbf{A} \notin \Phi$  or  $\neg \mathbf{A} \notin \Phi$ .

$\nabla_{\neg}$  If  $\neg \neg \mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{A} \in \Gamma_\Sigma$ .

$\nabla_\beta$  If  $\mathbf{A} =_\beta \mathbf{B}$  and  $\mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_\vee$  If  $\mathbf{A} \vee \mathbf{B} \in \Phi$ , then  $\Phi * \mathbf{A} \in \Gamma_\Sigma$  or  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_\wedge$  If  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , then  $\Phi * \neg \mathbf{A} * \neg \mathbf{B} \in \Gamma_\Sigma$ .

# Def.: Abstract Consistency Properties

Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{A}, \mathbf{B} \in \text{cwff}_o(\Sigma)$ ,  $\mathbf{F} \in \text{cwff}_{\alpha \rightarrow o}(\Sigma)$  are arbitrary):

$\nabla_c$  If  $\mathbf{A}$  is atomic, then  $\mathbf{A} \notin \Phi$  or  $\neg \mathbf{A} \notin \Phi$ .

$\nabla_{\neg}$  If  $\neg \neg \mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{A} \in \Gamma_\Sigma$ .

$\nabla_\beta$  If  $\mathbf{A} =_\beta \mathbf{B}$  and  $\mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_{\vee}$  If  $\mathbf{A} \vee \mathbf{B} \in \Phi$ , then  $\Phi * \mathbf{A} \in \Gamma_\Sigma$  or  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_{\wedge}$  If  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , then  $\Phi * \neg \mathbf{A} * \neg \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_{\forall}$  If  $\Pi^\alpha \mathbf{F} \in \Phi$ , then  $\Phi * \mathbf{F}\mathbf{W} \in \Gamma_\Sigma$  for each  $\mathbf{W} \in \text{cwff}_\alpha(\Sigma)$ .

# Def.: Abstract Consistency Properties

Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{A}, \mathbf{B} \in \text{cwff}_0(\Sigma)$ ,  $\mathbf{F} \in \text{cwff}_{\alpha \rightarrow o}(\Sigma)$  are arbitrary):

$\nabla_c$  If  $\mathbf{A}$  is atomic, then  $\mathbf{A} \notin \Phi$  or  $\neg \mathbf{A} \notin \Phi$ .

$\nabla_\neg$  If  $\neg \neg \mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{A} \in \Gamma_\Sigma$ .

$\nabla_\beta$  If  $\mathbf{A} =_\beta \mathbf{B}$  and  $\mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_\vee$  If  $\mathbf{A} \vee \mathbf{B} \in \Phi$ , then  $\Phi * \mathbf{A} \in \Gamma_\Sigma$  or  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_\wedge$  If  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , then  $\Phi * \neg \mathbf{A} * \neg \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_\forall$  If  $\Pi^\alpha \mathbf{F} \in \Phi$ , then  $\Phi * \mathbf{F}\mathbf{W} \in \Gamma_\Sigma$  for each  $\mathbf{W} \in \text{cwff}_\alpha(\Sigma)$ .

$\nabla_\exists$  If  $\neg \Pi^\alpha \mathbf{F} \in \Phi$ , then  $\Phi * \neg(\mathbf{F}\mathbf{w}) \in \Gamma_\Sigma$  for any parameter  $w_\alpha \in \Sigma_\alpha$  which does not occur in any sentence of  $\Phi$ .

# Def.: Abstract Consistency Properties

Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{A}, \mathbf{B} \in \text{cwff}_0(\Sigma)$ ,  $\mathbf{F} \in \text{cwff}_{\alpha \rightarrow o}(\Sigma)$  are arbitrary):

$\nabla_c$  If  $\mathbf{A}$  is atomic, then  $\mathbf{A} \notin \Phi$  or  $\neg \mathbf{A} \notin \Phi$ .

$\nabla_{\neg}$  If  $\neg \neg \mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{A} \in \Gamma_\Sigma$ .

$\nabla_\beta$  If  $\mathbf{A} =_\beta \mathbf{B}$  and  $\mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_{\vee}$  If  $\mathbf{A} \vee \mathbf{B} \in \Phi$ , then  $\Phi * \mathbf{A} \in \Gamma_\Sigma$  or  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_{\wedge}$  If  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , then  $\Phi * \neg \mathbf{A} * \neg \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_{\forall}$  If  $\Pi^\alpha \mathbf{F} \in \Phi$ , then  $\Phi * \mathbf{F}\mathbf{W} \in \Gamma_\Sigma$  for each  $\mathbf{W} \in \text{cwff}_\alpha(\Sigma)$ .

$\nabla_{\exists}$  If  $\neg \Pi^\alpha \mathbf{F} \in \Phi$ , then  $\Phi * \neg(\mathbf{F}\mathbf{w}) \in \Gamma_\Sigma$  for any parameter  $w_\alpha \in \Sigma_\alpha$  which does not occur in any sentence of  $\Phi$ .

(These properties are going back to Hintikka, Smullyan, and Andrews)

# Def.: Abstract Consistency Properties



Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{A}, \mathbf{B} \in \text{cwff}_0(\Sigma)$ ,  $\mathbf{G}, \mathbf{H}, (\lambda X_\alpha.\mathbf{M}), (\lambda X_\alpha.\mathbf{N}) \in \text{cwff}_{\alpha \rightarrow \beta}(\Sigma)$  are arbitrary):

# Def.: Abstract Consistency Properties



Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{A}, \mathbf{B} \in \text{cwff}_0(\Sigma)$ ,  $\mathbf{G}, \mathbf{H}, (\lambda X_\alpha.\mathbf{M}), (\lambda X_\alpha.\mathbf{N}) \in \text{cwff}_{\alpha \rightarrow \beta}(\Sigma)$  are arbitrary):

$\nabla_b$  If  $\neg(\mathbf{A} \doteq^\circ \mathbf{B}) \in \Phi$ , then  $\Phi * \mathbf{A} * \neg\mathbf{B} \in \Gamma_\Sigma$  or  $\Phi * \neg\mathbf{A} * \mathbf{B} \in \Gamma_\Sigma$ .

# Def.: Abstract Consistency Properties



Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{A}, \mathbf{B} \in \text{cwff}_0(\Sigma)$ ,  $\mathbf{G}, \mathbf{H}, (\lambda X_\alpha.\mathbf{M}), (\lambda X_\alpha.\mathbf{N}) \in \text{cwff}_{\alpha \rightarrow \beta}(\Sigma)$  are arbitrary):

$\nabla_b$  If  $\neg(\mathbf{A} \doteq^\circ \mathbf{B}) \in \Phi$ , then  $\Phi * \mathbf{A} * \neg \mathbf{B} \in \Gamma_\Sigma$  or  $\Phi * \neg \mathbf{A} * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_\eta$  If  $\mathbf{A} \stackrel{\beta\eta}{=} \mathbf{B}$  and  $\mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .

# Def.: Abstract Consistency Properties

Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{A}, \mathbf{B} \in \text{cwff}_0(\Sigma)$ ,  $\mathbf{G}, \mathbf{H}, (\lambda X_\alpha.\mathbf{M}), (\lambda X_\alpha.\mathbf{N}) \in \text{cwff}_{\alpha \rightarrow \beta}(\Sigma)$  are arbitrary):

$\nabla_b$  If  $\neg(\mathbf{A} \dot{=}^o \mathbf{B}) \in \Phi$ , then  $\Phi * \mathbf{A} * \neg \mathbf{B} \in \Gamma_\Sigma$  or  $\Phi * \neg \mathbf{A} * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_\eta$  If  $\mathbf{A} \stackrel{\beta\eta}{=} \mathbf{B}$  and  $\mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_\xi$  If  $\neg(\lambda X_\alpha.\mathbf{M} \dot{=}^{\alpha \rightarrow \beta} \lambda X_\alpha.\mathbf{N}) \in \Phi$ , then  
 $\Phi * \neg([\mathbf{w}/\mathbf{X}]\mathbf{M} \dot{=}^\beta [\mathbf{w}/\mathbf{X}]\mathbf{N}) \in \Gamma_\Sigma$  for any parameter  $w_\alpha \in \Sigma_\alpha$   
 which does not occur in any sentence of  $\Phi$ .



# Def.: Abstract Consistency Properties

Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{A}, \mathbf{B} \in \text{cwff}_0(\Sigma)$ ,  $\mathbf{G}, \mathbf{H}, (\lambda X_\alpha.\mathbf{M}), (\lambda X_\alpha.\mathbf{N}) \in \text{cwff}_{\alpha \rightarrow \beta}(\Sigma)$  are arbitrary):

$\nabla_b$  If  $\neg(\mathbf{A} \dot{=}^\circ \mathbf{B}) \in \Phi$ , then  $\Phi * \mathbf{A} * \neg \mathbf{B} \in \Gamma_\Sigma$  or  $\Phi * \neg \mathbf{A} * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_\eta$  If  $\mathbf{A} \stackrel{\beta\eta}{=} \mathbf{B}$  and  $\mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_\xi$  If  $\neg(\lambda X_\alpha.\mathbf{M} \dot{=}^{\alpha \rightarrow \beta} \lambda X_\alpha.\mathbf{N}) \in \Phi$ , then  $\Phi * \neg([\mathbf{w}/\mathbf{X}]\mathbf{M} \dot{=}^\beta [\mathbf{w}/\mathbf{X}]\mathbf{N}) \in \Gamma_\Sigma$  for any parameter  $\mathbf{w}_\alpha \in \Sigma_\alpha$  which does not occur in any sentence of  $\Phi$ .

$\nabla_f$  If  $\neg(\mathbf{G} \dot{=}^{\alpha \rightarrow \beta} \mathbf{H}) \in \Phi$ , then  $\Phi * \neg(\mathbf{G}\mathbf{w} \dot{=}^\beta \mathbf{H}\mathbf{w}) \in \Gamma_\Sigma$  for any parameter  $\mathbf{w}_\alpha \in \Sigma_\alpha$  which does not occur in any sentence of  $\Phi$ .

# Def.: Abstract Consistency Properties

Let  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences. We define (where  $\Phi \in \Gamma_\Sigma$ ,  $\alpha, \beta \in \mathcal{T}$ ,  $\mathbf{A}, \mathbf{B} \in \text{cwff}_0(\Sigma)$ ,  $\mathbf{G}, \mathbf{H}, (\lambda X_\alpha.\mathbf{M}), (\lambda X_\alpha.\mathbf{N}) \in \text{cwff}_{\alpha \rightarrow \beta}(\Sigma)$  are arbitrary):

$\nabla_b$  If  $\neg(\mathbf{A} \dot{=}^\circ \mathbf{B}) \in \Phi$ , then  $\Phi * \mathbf{A} * \neg \mathbf{B} \in \Gamma_\Sigma$  or  $\Phi * \neg \mathbf{A} * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_\eta$  If  $\mathbf{A} \stackrel{\beta\eta}{=} \mathbf{B}$  and  $\mathbf{A} \in \Phi$ , then  $\Phi * \mathbf{B} \in \Gamma_\Sigma$ .

$\nabla_\xi$  If  $\neg(\lambda X_\alpha.\mathbf{M} \dot{=}^{\alpha \rightarrow \beta} \lambda X_\alpha.\mathbf{N}) \in \Phi$ , then  $\Phi * \neg([\mathbf{w}/\mathbf{X}]\mathbf{M} \dot{=}^\beta [\mathbf{w}/\mathbf{X}]\mathbf{N}) \in \Gamma_\Sigma$  for any parameter  $\mathbf{w}_\alpha \in \Sigma_\alpha$  which does not occur in any sentence of  $\Phi$ .

$\nabla_f$  If  $\neg(\mathbf{G} \dot{=}^{\alpha \rightarrow \beta} \mathbf{H}) \in \Phi$ , then  $\Phi * \neg(\mathbf{G}\mathbf{w} \dot{=}^\beta \mathbf{H}\mathbf{w}) \in \Gamma_\Sigma$  for any parameter  $\mathbf{w}_\alpha \in \Sigma_\alpha$  which does not occur in any sentence of  $\Phi$ .

(These properties are new in [Benzmüller-PhD-99,JSL04])

# Def.: Abstract Consistency Classes



Let  $\Sigma$  be a signature and  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences that is closed under subsets.

# Def.: Abstract Consistency Classes



Let  $\Sigma$  be a signature and  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences that is closed under subsets.

If  $\nabla_c, \nabla_{\neg}, \nabla_{\beta}, \nabla_{\vee}, \nabla_{\wedge}, \nabla_{\forall}$  and  $\nabla_{\exists}$  are valid for  $\Gamma_\Sigma$ , then  $\Gamma_\Sigma$  is called an **abstract consistency class** for  $\Sigma$ -models.

# Def.: Abstract Consistency Classes

Let  $\Sigma$  be a signature and  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences that is closed under subsets.

If  $\nabla_c, \nabla_{\neg}, \nabla_\beta, \nabla_{\forall}, \nabla_{\wedge}, \nabla_{\vee}$  and  $\nabla_{\exists}$  are valid for  $\Gamma_\Sigma$ , then  $\Gamma_\Sigma$  is called an **abstract consistency class** for  $\Sigma$ -models.

We will denote the collection of abstract consistency classes by  $\mathcal{Acc}_\beta$ .

# Def.: Abstract Consistency Classes


Let  $\Sigma$  be a signature and  $\Gamma_\Sigma$  be a class of sets of  $\Sigma$ -sentences that is closed under subsets.

If  $\nabla_c, \nabla_{\neg}, \nabla_\beta, \nabla_{\forall}, \nabla_{\wedge}, \nabla_{\vee}$  and  $\nabla_{\exists}$  are valid for  $\Gamma_\Sigma$ , then  $\Gamma_\Sigma$  is called an **abstract consistency class** for  $\Sigma$ -models.

We will denote the collection of abstract consistency classes by  $\mathcal{Acc}_\beta$ .

Similarly, we introduce the following collections of specialized abstract consistency classes (with primitive equality):  $\mathcal{Acc}_{\beta\eta}, \mathcal{Acc}_{\beta\xi}, \mathcal{Acc}_{\beta f}, \mathcal{Acc}_{\beta b}, \mathcal{Acc}_{\beta\eta b}, \mathcal{Acc}_{\beta\xi b}, \mathcal{Acc}_{\beta fb}$ , where we indicate by indices which additional properties from  $\{\nabla_\eta, \nabla_\xi, \nabla_f, \nabla_b\}$  are required.

# Abstract Consistency Classes



# Ex.: Abstract Consistency Class



- not an abstract consistency class:

$\{\{\neg(A \vee B), \neg A\}, \{\neg(A \vee B)\}, \{\neg A\}, \{\}\}$



# Ex.: Abstract Consistency Class

- not an abstract consistency class:

$\{\{\neg(A \vee B), \neg A\}, \{\neg(A \vee B)\}, \{\neg A\}, \{\}\}$

- still not:

$\{\{\neg(A \vee B), \neg A, \neg B\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B)\}, \{\neg A\}, \{\}\}$

# Ex.: Abstract Consistency Class

- not an abstract consistency class:

$$\{\{\neg(A \vee B), \neg A\}, \{\neg(A \vee B)\}, \{\neg A\}, \{\}\}$$

- still not:

$$\{\{\neg(A \vee B), \neg A, \neg B\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B)\}, \{\neg A\}, \{\}\}$$

- how about this one:

$$\Gamma := \{\{\neg(A \vee B), \neg A, \neg B\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B), \neg B\}, \{\neg A, \neg B\}, \{\neg(A \vee B)\}, \{\neg A\}, \{\neg B\}, \{\}\}$$

# Ex.: Abstract Consistency Class

- not an abstract consistency class:

$$\{\{\neg(A \vee B), \neg A\}, \{\neg(A \vee B)\}, \{\neg A\}, \{\}\}$$

- still not:

$$\{\{\neg(A \vee B), \neg A, \neg B\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B)\}, \{\neg A\}, \{\}\}$$

- how about this one:

$$\Gamma := \{\{\neg(A \vee B), \neg A, \neg B\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B), \neg B\}, \{\neg A, \neg B\}, \{\neg(A \vee B)\}, \{\neg A\}, \{\neg B\}, \{\}\}$$

- and how about this:

$$\Gamma_0 := \Gamma$$

$$\Phi \in \Gamma_i \wedge A \in \Phi \wedge B =_{\beta\eta} A \wedge B \neq A \wedge (\Phi * B) \notin \Gamma_i \longrightarrow$$

$$\Gamma_{i+1} := \text{close-under-subsets}(\Gamma_i * (\Phi * B))$$

$$\Gamma^* := \Gamma_\infty$$

# Rem.: Possible Generalization

The work presented here is based on the choice of the primitive logical connectives  $\neg$ ,  $\vee$  and  $\Pi^\alpha$ .

# Rem.: Possible Generalization

The work presented here is based on the choice of the primitive logical connectives  $\neg$ ,  $\vee$  and  $\Box^\alpha$ . A means to generalize the framework over the concrete choice of logical primitives is provided by the uniform notation approach as, for instance, given in [Fitting96].

# Rem.: Possible Generalization

The work presented here is based on the choice of the primitive logical connectives  $\neg$ ,  $\vee$  and  $\Box^\alpha$ . A means to generalize the framework over the concrete choice of logical primitives is provided by the uniform notation approach as, for instance, given in [Fitting96]. This can be done in straightforward manner:  $\nabla_\wedge$  becomes an  $\alpha$ -property,  $\nabla_\vee$  becomes a  $\beta$ -property,  $\nabla_\forall$  becomes a  $\gamma$ -property, and  $\nabla_\exists$  becomes a  $\delta$ -property.

# Rem.: Possible Generalization

The work presented here is based on the choice of the primitive logical connectives  $\neg$ ,  $\vee$  and  $\Box^\alpha$ . A means to generalize the framework over the concrete choice of logical primitives is provided by the uniform notation approach as, for instance, given in [Fitting96]. This can be done in straightforward manner:  $\nabla_\wedge$  becomes an  $\alpha$ -property,  $\nabla_\vee$  becomes a  $\beta$ -property,  $\nabla_\forall$  becomes a  $\gamma$ -property, and  $\nabla_\exists$  becomes a  $\delta$ -property. Thus they will have the following form:

# Rem.: Possible Generalization

The work presented here is based on the choice of the primitive logical connectives  $\neg$ ,  $\vee$  and  $\Box^\alpha$ . A means to generalize the framework over the concrete choice of logical primitives is provided by the uniform notation approach as, for instance, given in [Fitting96]. This can be done in straightforward manner:  $\nabla_\wedge$  becomes an  $\alpha$ -property,  $\nabla_\vee$  becomes a  $\beta$ -property,  $\nabla_\forall$  becomes a  $\gamma$ -property, and  $\nabla_\exists$  becomes a  $\delta$ -property. Thus they will have the following form:

**$\alpha$ -case** If  $\alpha \in \Phi$ , then  $\Phi * \alpha_1 * \alpha_2 \in \Gamma_\Sigma$ .



# Rem.: Possible Generalization

The work presented here is based on the choice of the primitive logical connectives  $\neg$ ,  $\vee$  and  $\Box^\alpha$ . A means to generalize the framework over the concrete choice of logical primitives is provided by the uniform notation approach as, for instance, given in [Fitting96]. This can be done in straightforward manner:  $\nabla_\wedge$  becomes an  $\alpha$ -property,  $\nabla_\vee$  becomes a  $\beta$ -property,  $\nabla_\forall$  becomes a  $\gamma$ -property, and  $\nabla_\exists$  becomes a  $\delta$ -property. Thus they will have the following form:

**$\alpha$ -case** If  $\alpha \in \Phi$ , then  $\Phi * \alpha_1 * \alpha_2 \in \Gamma_\Sigma$ .

**$\beta$ -case** If  $\beta \in \Phi$ , then  $\Phi * \beta_1 \in \Gamma_\Sigma$  or  $\Phi * \beta_2 \in \Gamma_\Sigma$ .

# Rem.: Possible Generalization

The work presented here is based on the choice of the primitive logical connectives  $\neg$ ,  $\vee$  and  $\Pi^\alpha$ . A means to generalize the framework over the concrete choice of logical primitives is provided by the uniform notation approach as, for instance, given in [Fitting96]. This can be done in straightforward manner:  $\nabla_\wedge$  becomes an  $\alpha$ -property,  $\nabla_\vee$  becomes a  $\beta$ -property,  $\nabla_\forall$  becomes a  $\gamma$ -property, and  $\nabla_\exists$  becomes a  $\delta$ -property. Thus they will have the following form:

**$\alpha$ -case** If  $\alpha \in \Phi$ , then  $\Phi * \alpha_1 * \alpha_2 \in \Gamma_\Sigma$ .

**$\beta$ -case** If  $\beta \in \Phi$ , then  $\Phi * \beta_1 \in \Gamma_\Sigma$  or  $\Phi * \beta_2 \in \Gamma_\Sigma$ .

**$\gamma$ -case** If  $\gamma \in \Phi$ , then  $\Phi * \gamma \mathbf{W} \in \Gamma_\Sigma$  for each  $\mathbf{W} \in cwff_\alpha(\Sigma)$ .

# Rem.: Possible Generalization

The work presented here is based on the choice of the primitive logical connectives  $\neg$ ,  $\vee$  and  $\prod^\alpha$ . A means to generalize the framework over the concrete choice of logical primitives is provided by the uniform notation approach as, for instance, given in [Fitting96]. This can be done in straightforward manner:  $\nabla_\wedge$  becomes an  $\alpha$ -property,  $\nabla_\vee$  becomes a  $\beta$ -property,  $\nabla_\forall$  becomes a  $\gamma$ -property, and  $\nabla_\exists$  becomes a  $\delta$ -property. Thus they will have the following form:

**$\alpha$ -case** If  $\alpha \in \Phi$ , then  $\Phi * \alpha_1 * \alpha_2 \in \Gamma_\Sigma$ .

**$\beta$ -case** If  $\beta \in \Phi$ , then  $\Phi * \beta_1 \in \Gamma_\Sigma$  or  $\Phi * \beta_2 \in \Gamma_\Sigma$ .

**$\gamma$ -case** If  $\gamma \in \Phi$ , then  $\Phi * \gamma \mathbf{W} \in \Gamma_\Sigma$  for each  $\mathbf{W} \in \text{cwff}_\alpha(\Sigma)$ .

**$\delta$ -case** If  $\delta \in \Phi$ , then  $\Phi * \delta w \in \Gamma_\Sigma$  for any parameter  $w_\alpha \in \Sigma$  which does not occur in any sentence of  $\Phi$ .

# Def.: Saturated

---



Consider the following property (where  $\phi \in \Gamma_\Sigma$ ,  $A \in \text{cwff}_o(\Sigma)$ ):

# Def.: Saturated

Consider the following property (where  $\Phi \in \Gamma_\Sigma$ ,  $A \in \text{cwff}_o(\Sigma)$ ):

$\nabla_{\text{sat}}$  Either  $\Phi * A \in \Gamma_\Sigma$  or  $\Phi * \neg A \in \Gamma_\Sigma$ .

# Def.: Saturated

Consider the following property (where  $\Phi \in \Gamma_\Sigma$ ,  $A \in \text{cwff}_0(\Sigma)$ ):

$\nabla_{\text{sat}}$  Either  $\Phi * A \in \Gamma_\Sigma$  or  $\Phi * \neg A \in \Gamma_\Sigma$ .

We call an abstract consistency class  $\Gamma_\Sigma$  **atomically saturated** if  $\nabla_{\text{sat}}$  holds for all atomic  $A$ .

# Def.: Saturated

Consider the following property (where  $\Phi \in \Gamma_\Sigma$ ,  $A \in \text{cwff}_o(\Sigma)$ ):

$\nabla_{\text{sat}}$  Either  $\Phi * A \in \Gamma_\Sigma$  or  $\Phi * \neg A \in \Gamma_\Sigma$ .

We call an abstract consistency class  $\Gamma_\Sigma$  **atomically saturated** if  $\nabla_{\text{sat}}$  holds for all atomic  $A$ .

We call an abstract consistency class  $\Gamma_\Sigma$  **saturated** if  $\nabla_{\text{sat}}$  holds for all  $A$ .

# Ex.: Saturated



- consider  $\Gamma$  (and  $\Gamma^*$ ) from before:

$\{\{\neg(A \vee B), \neg A, \neg B\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B), \neg B\}, \{\neg A, \neg B\}, \{\neg(A \vee B)\}, \{\neg A\}, \{\neg B\}, \{\}\}$



# Ex.: Saturated

- consider  $\Gamma$  (and  $\Gamma^*$ ) from before:  
 $\{\{\neg(A \vee B), \neg A, \neg B\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B), \neg B\}, \{\neg A, \neg B\}, \{\neg(A \vee B)\}, \{\neg A\}, \{\neg B\}, \{\}\}$
- $\Gamma$  (and  $\Gamma^*$ ) is atomically saturated in case our signature contains no further constants besides  $A_0$  and  $B_0$  and the logical connectives.

# Ex.: Saturated

- consider  $\Gamma$  (and  $\Gamma^*$ ) from before:  
 $\{\{\neg(A \vee B), \neg A, \neg B\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B), \neg B\}, \{\neg A, \neg B\}, \{\neg(A \vee B)\}, \{\neg A\}, \{\neg B\}, \{\}\}$
- $\Gamma$  (and  $\Gamma^*$ ) is atomically saturated in case our signature contains no further constants besides  $A_0$  and  $B_0$  and the logical connectives.
- if there is another symbol  $C_0$  in the signature, then  $\Gamma$  (and  $\Gamma^*$ ) is not atomically saturated anymore

# Ex.: Saturated

- consider  $\Gamma$  (and  $\Gamma^*$ ) from before:  
 $\{\{\neg(A \vee B), \neg A, \neg B\}, \{\neg(A \vee B), \neg A\}, \{\neg(A \vee B), \neg B\}, \{\neg A, \neg B\}, \{\neg(A \vee B)\}, \{\neg A\}, \{\neg B\}, \{\}\}$
- $\Gamma$  (and  $\Gamma^*$ ) is atomically saturated in case our signature contains no further constants besides  $A_0$  and  $B_0$  and the logical connectives.
- if there is another symbol  $C_0$  in the signature, then  $\Gamma$  (and  $\Gamma^*$ ) is not atomically saturated anymore
- $\Gamma$  (and  $\Gamma^*$ ) is not saturated: for instance, it does not provide information on the formulas  $(\neg A \vee B) \vee A$  and  $\Pi^0(\lambda X_0.X)$

# Thm.: Model Existence Theorem



Let  $\Gamma_\Sigma$  be a saturated abstract consistency class and let  $\Phi \in \Gamma_\Sigma$  be a sufficiently  $\Sigma$ -pure set of sentences.

For all  $*$   $\in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$  we have:

# Thm.: Model Existence Theorem

Let  $\Gamma_\Sigma$  be a saturated abstract consistency class and let  $\Phi \in \Gamma_\Sigma$  be a sufficiently  $\Sigma$ -pure set of sentences.

For all  $* \in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$  we have:

If  $\Gamma_\Sigma$  is an  $\mathcal{Acc}_*$ , then there exists a model  $\mathcal{M} \in \mathfrak{M}_*$  that satisfies  $\Phi$ .

# Thm.: Model Existence Theorem

Let  $\Gamma_\Sigma$  be a saturated abstract consistency class and let  $\Phi \in \Gamma_\Sigma$  be a sufficiently  $\Sigma$ -pure set of sentences.

For all  $*$   $\in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$  we have:

If  $\Gamma_\Sigma$  is an  $\mathcal{Acc}_*$ , then there exists a model  $\mathcal{M} \in \mathfrak{M}_*$  that satisfies  $\Phi$ .

Furthermore, each domain of  $\mathcal{M}$  has cardinality at most  $\aleph_s$ .

# Thm.: Model Existence Theorem

Let  $\Gamma_\Sigma$  be a saturated abstract consistency class and let  $\Phi \in \Gamma_\Sigma$  be a sufficiently  $\Sigma$ -pure set of sentences.

For all  $* \in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$  we have:

If  $\Gamma_\Sigma$  is an  $\mathcal{Acc}_*$ , then there exists a model  $\mathcal{M} \in \mathfrak{M}_*$  that satisfies  $\Phi$ .

Furthermore, each domain of  $\mathcal{M}$  has cardinality at most  $\aleph_s$ .

Proof: ... we are not yet ready for this ...

# Thm.: Model Existence for Henkin Models



Let  $\Gamma_\Sigma$  be a saturated abstract consistency class in  $\mathcal{Acc}_{\beta fb}$  and let  $\Phi \in \Gamma_\Sigma$  be a sufficiently  $\Sigma$ -pure set of sentences.



# Thm.: Model Existence for Henkin Models



Let  $\Gamma_\Sigma$  be a saturated abstract consistency class in  $\mathcal{ACC}_{\beta\text{fb}}$  and let  $\Phi \in \Gamma_\Sigma$  be a sufficiently  $\Sigma$ -pure set of sentences.

Then there is a Henkin Model that satisfies  $\Phi$ .

# Thm.: Model Existence for Henkin Models



Let  $\Gamma_\Sigma$  be a saturated abstract consistency class in  $\mathcal{Acc}_{\beta fb}$  and let  $\Phi \in \Gamma_\Sigma$  be a sufficiently  $\Sigma$ -pure set of sentences.

Then there is a Henkin Model that satisfies  $\Phi$ .

Furthermore, each domain of the model has cardinality at most  $\aleph_s$ .

# Thm.: Model Existence for Henkin Models



Let  $\Gamma_\Sigma$  be a saturated abstract consistency class in  $\mathcal{Acc}_{\beta\text{fb}}$  and let  $\Phi \in \Gamma_\Sigma$  be a sufficiently  $\Sigma$ -pure set of sentences.

Then there is a Henkin Model that satisfies  $\Phi$ .

Furthermore, each domain of the model has cardinality at most  $\aleph_s$ .

Proof: ... we are not yet ready for this ...



Completeness of  $\mathcal{N}_*$  via  
Abstract Consistency

# Def.: $\mathcal{N}_*$ -Consistent/Inconsistent



A set of sentences  $\Phi$  is  $\mathcal{N}_*$ -inconsistent if  $\Phi \Vdash_{\mathcal{N}_*} \mathbf{F}_o$ , and  $\mathcal{N}_*$ -consistent otherwise.

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_0(\Sigma) \mid \Phi \text{ is } \mathcal{N}\mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_0(\Sigma) \mid \Phi \text{ is } \mathcal{N}\mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{N}\mathcal{K}_*$ -consistent set is  $\mathcal{N}\mathcal{K}_*$ -consistent.

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_0(\Sigma) \mid \Phi \text{ is } \mathcal{N}\mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{N}\mathcal{K}_*$ -consistent set is  $\mathcal{N}\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.



# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_0(\Sigma) \mid \Phi \text{ is } \mathcal{N}\mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{N}\mathcal{K}_*$ -consistent set is  $\mathcal{N}\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_0(\Sigma) \mid \Phi \text{ is } \mathcal{N}\mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{N}\mathcal{K}_*$ -consistent set is  $\mathcal{N}\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.

$\nabla_c$  Suppose  $A, \neg A \in \Phi$ .

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_o(\Sigma) \mid \Phi \text{ is } \mathcal{N}\mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{N}\mathcal{K}_*$ -consistent set is  $\mathcal{N}\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.

$\nabla_c$  Suppose  $\mathbf{A}, \neg\mathbf{A} \in \Phi$ . We have  $\Phi \Vdash \mathbf{F}_o$  by  $\mathcal{N}\mathcal{K}(\text{Hyp})$  and  $\mathcal{N}\mathcal{K}(\neg E)$ .

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_0(\Sigma) \mid \Phi \text{ is } \mathcal{N}\mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{N}\mathcal{K}_*$ -consistent set is  $\mathcal{N}\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.

$\nabla_c$  Suppose  $\mathbf{A}, \neg\mathbf{A} \in \Phi$ . We have  $\Phi \Vdash \mathbf{F}_0$  by  $\mathcal{N}\mathcal{K}(\text{Hyp})$  and  $\mathcal{N}\mathcal{K}(\neg E)$ .  
Hence  $\Phi$  is  $\mathcal{N}\mathcal{K}_*$ -inconsistent.

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_0(\Sigma) \mid \Phi \text{ is } \mathcal{N}\mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{N}\mathcal{K}_*$ -consistent set is  $\mathcal{N}\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.

$\nabla_c$  Suppose  $\mathbf{A}, \neg\mathbf{A} \in \Phi$ . We have  $\Phi \Vdash \mathbf{F}_0$  by  $\mathcal{N}\mathcal{K}(\text{Hyp})$  and  $\mathcal{N}\mathcal{K}(\neg E)$ .  
Hence  $\Phi$  is  $\mathcal{N}\mathcal{K}_*$ -inconsistent.

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_0(\Sigma) \mid \Phi \text{ is } \mathcal{N}\mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{N}\mathcal{K}_*$ -consistent set is  $\mathcal{N}\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.

$\nabla_c$  Suppose  $\mathbf{A}, \neg\mathbf{A} \in \Phi$ . We have  $\Phi \Vdash \mathbf{F}_0$  by  $\mathcal{N}\mathcal{K}(\text{Hyp})$  and  $\mathcal{N}\mathcal{K}(\neg E)$ .  
Hence  $\Phi$  is  $\mathcal{N}\mathcal{K}_*$ -inconsistent.

$\nabla_\beta$  Let  $\mathbf{A} \in \Phi$ ,  $\mathbf{A} =_\beta \mathbf{B}$  and  $\Phi * \mathbf{B}$  be  $\mathcal{N}\mathcal{K}_*$ -inconsistent.

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_o(\Sigma) \mid \Phi \text{ is } \mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{K}_*$ -consistent set is  $\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.

$\nabla_c$  Suppose  $\mathbf{A}, \neg\mathbf{A} \in \Phi$ . We have  $\Phi \Vdash \mathbf{F}_o$  by  $\mathcal{K}(\text{Hyp})$  and  $\mathcal{K}(\neg E)$ .  
Hence  $\Phi$  is  $\mathcal{K}_*$ -inconsistent.

$\nabla_\beta$  Let  $\mathbf{A} \in \Phi$ ,  $\mathbf{A} =_\beta \mathbf{B}$  and  $\Phi * \mathbf{B}$  be  $\mathcal{K}_*$ -inconsistent. That is,  
 $\Phi * \mathbf{B} \Vdash \mathbf{F}_o$ .

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_o(\Sigma) \mid \Phi \text{ is } \mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{K}_*$ -consistent set is  $\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.

$\nabla_c$  Suppose  $\mathbf{A}, \neg\mathbf{A} \in \Phi$ . We have  $\Phi \Vdash \mathbf{F}_o$  by  $\mathcal{K}(\text{Hyp})$  and  $\mathcal{K}(\neg E)$ .  
Hence  $\Phi$  is  $\mathcal{K}_*$ -inconsistent.

$\nabla_\beta$  Let  $\mathbf{A} \in \Phi$ ,  $\mathbf{A} =_\beta \mathbf{B}$  and  $\Phi * \mathbf{B}$  be  $\mathcal{K}_*$ -inconsistent. That is,  
 $\Phi * \mathbf{B} \Vdash \mathbf{F}_o$ . By  $\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg\mathbf{B}$ .



# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_0(\Sigma) \mid \Phi \text{ is } \mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{K}_*$ -consistent set is  $\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.

$\nabla_c$  Suppose  $\mathbf{A}, \neg\mathbf{A} \in \Phi$ . We have  $\Phi \Vdash \mathbf{F}_o$  by  $\mathcal{K}(\text{Hyp})$  and  $\mathcal{K}(\neg E)$ .  
Hence  $\Phi$  is  $\mathcal{K}_*$ -inconsistent.

$\nabla_\beta$  Let  $\mathbf{A} \in \Phi$ ,  $\mathbf{A} =_\beta \mathbf{B}$  and  $\Phi * \mathbf{B}$  be  $\mathcal{K}_*$ -inconsistent. That is,  
 $\Phi * \mathbf{B} \Vdash \mathbf{F}_o$ . By  $\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg\mathbf{B}$ . Since  $\mathbf{A} \in \Phi$ , we know  
 $\Phi \Vdash \mathbf{B}$  by  $\mathcal{K}(\text{Hyp})$  and  $\mathcal{K}(\beta)$ .

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_0(\Sigma) \mid \Phi \text{ is } \mathcal{N}\mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{N}\mathcal{K}_*$ -consistent set is  $\mathcal{N}\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.

$\nabla_c$  Suppose  $\mathbf{A}, \neg\mathbf{A} \in \Phi$ . We have  $\Phi \Vdash \mathbf{F}_0$  by  $\mathcal{N}\mathcal{K}(\text{Hyp})$  and  $\mathcal{N}\mathcal{K}(\neg E)$ .  
Hence  $\Phi$  is  $\mathcal{N}\mathcal{K}_*$ -inconsistent.

$\nabla_\beta$  Let  $\mathbf{A} \in \Phi$ ,  $\mathbf{A} =_\beta \mathbf{B}$  and  $\Phi * \mathbf{B}$  be  $\mathcal{N}\mathcal{K}_*$ -inconsistent. That is,  
 $\Phi * \mathbf{B} \Vdash \mathbf{F}_0$ . By  $\mathcal{N}\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg\mathbf{B}$ . Since  $\mathbf{A} \in \Phi$ , we know  
 $\Phi \Vdash \mathbf{B}$  by  $\mathcal{N}\mathcal{K}(\text{Hyp})$  and  $\mathcal{N}\mathcal{K}(\beta)$ . Using  $\mathcal{N}\mathcal{K}(\neg E)$ , we know  $\Phi \Vdash \mathbf{F}_0$ .

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_o(\Sigma) \mid \Phi \text{ is } \mathcal{K}_* \text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{K}_*$ -consistent set is  $\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.

$\nabla_c$  Suppose  $\mathbf{A}, \neg \mathbf{A} \in \Phi$ . We have  $\Phi \Vdash \mathbf{F}_o$  by  $\mathcal{K}(\text{Hyp})$  and  $\mathcal{K}(\neg E)$ .  
Hence  $\Phi$  is  $\mathcal{K}_*$ -inconsistent.

$\nabla_\beta$  Let  $\mathbf{A} \in \Phi$ ,  $\mathbf{A} =_\beta \mathbf{B}$  and  $\Phi * \mathbf{B}$  be  $\mathcal{K}_*$ -inconsistent. That is,  
 $\Phi * \mathbf{B} \Vdash \mathbf{F}_o$ . By  $\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg \mathbf{B}$ . Since  $\mathbf{A} \in \Phi$ , we know  
 $\Phi \Vdash \mathbf{B}$  by  $\mathcal{K}(\text{Hyp})$  and  $\mathcal{K}(\beta)$ . Using  $\mathcal{K}(\neg E)$ , we know  $\Phi \Vdash \mathbf{F}_o$ .

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_o(\Sigma) \mid \Phi \text{ is } \mathcal{K}_* \text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{K}_*$ -consistent set is  $\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.

$\nabla_c$  Suppose  $\mathbf{A}, \neg \mathbf{A} \in \Phi$ . We have  $\Phi \Vdash \mathbf{F}_o$  by  $\mathcal{K}(\text{Hyp})$  and  $\mathcal{K}(\neg E)$ .  
Hence  $\Phi$  is  $\mathcal{K}_*$ -inconsistent.

$\nabla_\beta$  Let  $\mathbf{A} \in \Phi$ ,  $\mathbf{A} =_\beta \mathbf{B}$  and  $\Phi * \mathbf{B}$  be  $\mathcal{K}_*$ -inconsistent. That is,  
 $\Phi * \mathbf{B} \Vdash \mathbf{F}_o$ . By  $\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg \mathbf{B}$ . Since  $\mathbf{A} \in \Phi$ , we know  
 $\Phi \Vdash \mathbf{B}$  by  $\mathcal{K}(\text{Hyp})$  and  $\mathcal{K}(\beta)$ . Using  $\mathcal{K}(\neg E)$ , we know  $\Phi \Vdash \mathbf{F}_o$ .

$\nabla_{\neg}$  Suppose  $\neg \neg \mathbf{A} \in \Phi$  and  $\Phi * \mathbf{A}$  is  $\mathcal{K}_*$ -inconsistent.

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_o(\Sigma) \mid \Phi \text{ is } \mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{K}_*$ -consistent set is  $\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.

$\nabla_c$  Suppose  $\mathbf{A}, \neg\mathbf{A} \in \Phi$ . We have  $\Phi \Vdash \mathbf{F}_o$  by  $\mathcal{K}(\text{Hyp})$  and  $\mathcal{K}(\neg E)$ .  
Hence  $\Phi$  is  $\mathcal{K}_*$ -inconsistent.

$\nabla_\beta$  Let  $\mathbf{A} \in \Phi$ ,  $\mathbf{A} =_\beta \mathbf{B}$  and  $\Phi * \mathbf{B}$  be  $\mathcal{K}_*$ -inconsistent. That is,  
 $\Phi * \mathbf{B} \Vdash \mathbf{F}_o$ . By  $\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg\mathbf{B}$ . Since  $\mathbf{A} \in \Phi$ , we know  
 $\Phi \Vdash \mathbf{B}$  by  $\mathcal{K}(\text{Hyp})$  and  $\mathcal{K}(\beta)$ . Using  $\mathcal{K}(\neg E)$ , we know  $\Phi \Vdash \mathbf{F}_o$ .

$\nabla_\neg$  Suppose  $\neg\neg\mathbf{A} \in \Phi$  and  $\Phi * \mathbf{A}$  is  $\mathcal{K}_*$ -inconsistent. From  $\Phi * \mathbf{A} \Vdash \mathbf{F}_o$   
and  $\mathcal{K}(\neg I)$ , we have  $\Phi \Vdash \neg\mathbf{A}$ .

# Lemma: Saturated $\mathcal{Acc}_*$

The class  $\Gamma_\Sigma^* := \{\Phi \subseteq \text{cwff}_0(\Sigma) \mid \Phi \text{ is } \mathcal{K}_*\text{-consistent}\}$  is a saturated  $\mathcal{Acc}_*$ .

Proof: Obviously  $\Gamma_\Sigma^*$  is closed under subsets, since any subset of an  $\mathcal{K}_*$ -consistent set is  $\mathcal{K}_*$ -consistent. We now check the remaining conditions. We prove all the properties by proving their contrapositive.

$\nabla_c$  Suppose  $\mathbf{A}, \neg\mathbf{A} \in \Phi$ . We have  $\Phi \Vdash \mathbf{F}_0$  by  $\mathcal{K}(\text{Hyp})$  and  $\mathcal{K}(\neg E)$ .  
Hence  $\Phi$  is  $\mathcal{K}_*$ -inconsistent.

$\nabla_\beta$  Let  $\mathbf{A} \in \Phi$ ,  $\mathbf{A} =_\beta \mathbf{B}$  and  $\Phi * \mathbf{B}$  be  $\mathcal{K}_*$ -inconsistent. That is,  
 $\Phi * \mathbf{B} \Vdash \mathbf{F}_0$ . By  $\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg\mathbf{B}$ . Since  $\mathbf{A} \in \Phi$ , we know  
 $\Phi \Vdash \mathbf{B}$  by  $\mathcal{K}(\text{Hyp})$  and  $\mathcal{K}(\beta)$ . Using  $\mathcal{K}(\neg E)$ , we know  $\Phi \Vdash \mathbf{F}_0$ .

$\nabla_\neg$  Suppose  $\neg\neg\mathbf{A} \in \Phi$  and  $\Phi * \mathbf{A}$  is  $\mathcal{K}_*$ -inconsistent. From  $\Phi * \mathbf{A} \Vdash \mathbf{F}_0$   
and  $\mathcal{K}(\neg I)$ , we have  $\Phi \Vdash \neg\mathbf{A}$ . Since  $\neg\neg\mathbf{A} \in \Phi$ , we can apply  
 $\mathcal{K}(\text{Hyp})$  and  $\mathcal{K}(\neg E)$  to obtain  $\Phi \Vdash \mathbf{F}_0$ .

# Lemma: Saturated $\mathcal{Acc}_*$



# Lemma: Saturated $\mathcal{Acc}_*$



$\nabla_V$  Suppose  $(A \vee B) \in \Phi$  and both  $\Phi * A$  and  $\Phi * B$  are  $\mathcal{Acc}_*$ -inconsistent.



# Lemma: Saturated $\mathcal{Acc}_*$



$\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathcal{N}\mathcal{K}_*$ -inconsistent.  
By  $\mathcal{N}\mathcal{K}(Hyp)$  and  $\mathcal{N}\mathcal{K}(\vee E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .

# Lemma: Saturated $\mathcal{Acc}_*$



$\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathcal{N}\mathcal{K}_*$ -inconsistent.  
By  $\mathcal{N}\mathcal{K}(Hyp)$  and  $\mathcal{N}\mathcal{K}(\vee E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .

# Lemma: Saturated $\mathcal{Acc}_*$

- $\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathcal{N}\mathcal{K}_*$ -inconsistent.  
By  $\mathcal{N}\mathcal{K}(Hyp)$  and  $\mathcal{N}\mathcal{K}(\vee E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .
- $\nabla_{\wedge}$  Suppose  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and  $\Phi * \neg\mathbf{A} * \neg\mathbf{B}$  is  $\mathcal{N}\mathcal{K}_*$ -inconsistent.

# Lemma: Saturated $\mathcal{Acc}_*$

- $\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathcal{N}\mathcal{K}_*$ -inconsistent. By  $\mathcal{N}\mathcal{K}(Hyp)$  and  $\mathcal{N}\mathcal{K}(\vee E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .
- $\nabla_{\wedge}$  Suppose  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and  $\Phi * \neg\mathbf{A} * \neg\mathbf{B}$  is  $\mathcal{N}\mathcal{K}_*$ -inconsistent. By  $\mathcal{N}\mathcal{K}(Contr)$  and  $\mathcal{N}\mathcal{K}(\vee I_R)$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{A} \vee \mathbf{B}$ .

# Lemma: Saturated $\mathcal{Acc}_*$

- $\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\vee E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .
- $\nabla_{\wedge}$  Suppose  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and  $\Phi * \neg\mathbf{A} * \neg\mathbf{B}$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_R)$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{F}_o$ .

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\vee E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .
- $\nabla_{\wedge}$  Suppose  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and  $\Phi * \neg\mathbf{A} * \neg\mathbf{B}$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_R)$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{F}_o$ . By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_L)$ , we have  $\Phi \Vdash \mathbf{A} \vee \mathbf{B}$ .

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\vee E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .
- $\nabla_{\wedge}$  Suppose  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and  $\Phi * \neg\mathbf{A} * \neg\mathbf{B}$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_R)$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{F}_o$ . By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_L)$ , we have  $\Phi \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ ,  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\vee E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .
- $\nabla_{\wedge}$  Suppose  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and  $\Phi * \neg\mathbf{A} * \neg\mathbf{B}$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_R)$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{F}_o$ . By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_L)$ , we have  $\Phi \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ ,  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.



# Lemma: Saturated $\mathcal{Acc}_*$

- $\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\vee E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .
- $\nabla_{\wedge}$  Suppose  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and  $\Phi * \neg\mathbf{A} * \neg\mathbf{B}$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_R)$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{F}_o$ . By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_L)$ , we have  $\Phi \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ ,  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.
- $\nabla_{\forall}$  Suppose  $(\Pi^\alpha \mathbf{G}) \in \Phi$  and  $\Phi * (\mathbf{G}\mathbf{A})$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\vee E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .
- $\nabla_{\wedge}$  Suppose  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and  $\Phi * \neg\mathbf{A} * \neg\mathbf{B}$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_R)$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{F}_o$ . By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_L)$ , we have  $\Phi \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ ,  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.
- $\nabla_{\forall}$  Suppose  $(\Pi^\alpha \mathbf{G}) \in \Phi$  and  $\Phi * (\mathbf{G}\mathbf{A})$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(\neg I)$ ,  $\Phi \Vdash \neg(\mathbf{G}\mathbf{A})$ .

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\vee E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .
- $\nabla_{\wedge}$  Suppose  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and  $\Phi * \neg\mathbf{A} * \neg\mathbf{B}$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_R)$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{F}_o$ . By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_L)$ , we have  $\Phi \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ ,  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.
- $\nabla_{\forall}$  Suppose  $(\Pi^\alpha \mathbf{G}) \in \Phi$  and  $\Phi * (\mathbf{G}\mathbf{A})$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(\neg I)$ ,  $\Phi \Vdash \neg(\mathbf{G}\mathbf{A})$ . By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\Pi E)$ ,  $\Phi \Vdash \mathbf{G}\mathbf{A}$ .

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\vee E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .
- $\nabla_{\wedge}$  Suppose  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and  $\Phi * \neg\mathbf{A} * \neg\mathbf{B}$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_R)$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{F}_o$ . By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_L)$ , we have  $\Phi \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ ,  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.
- $\nabla_{\forall}$  Suppose  $(\Pi^\alpha \mathbf{G}) \in \Phi$  and  $\Phi * (\mathbf{G}\mathbf{A})$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(\neg I)$ ,  $\Phi \Vdash \neg(\mathbf{G}\mathbf{A})$ . By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\Pi E)$ ,  $\Phi \Vdash \mathbf{G}\mathbf{A}$ . Finally,  $\mathfrak{N}\mathcal{K}(\neg E)$  implies  $\Phi \Vdash \mathbf{F}_o$ .

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\vee E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .
- $\nabla_{\wedge}$  Suppose  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and  $\Phi * \neg\mathbf{A} * \neg\mathbf{B}$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_R)$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{F}_o$ . By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_L)$ , we have  $\Phi \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ ,  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.
- $\nabla_{\neg}$  Suppose  $(\Pi^\alpha \mathbf{G}) \in \Phi$  and  $\Phi * (\mathbf{G}\mathbf{A})$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(\neg I)$ ,  $\Phi \Vdash \neg(\mathbf{G}\mathbf{A})$ . By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\Pi E)$ ,  $\Phi \Vdash \mathbf{G}\mathbf{A}$ . Finally,  $\mathfrak{N}\mathcal{K}(\neg E)$  implies  $\Phi \Vdash \mathbf{F}_o$ .
- $\nabla_{\exists}$  Suppose  $\neg(\Pi^\alpha \mathbf{G}) \in \Phi$ ,  $w_\alpha$  is a parameter which does not occur in  $\Phi$ , and  $\Phi * \neg(\mathbf{G}w)$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\vee E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .
- $\nabla_{\wedge}$  Suppose  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and  $\Phi * \neg\mathbf{A} * \neg\mathbf{B}$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_R)$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , we have  $\Phi, \neg\mathbf{A} \Vdash \mathbf{F}_o$ . By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_L)$ , we have  $\Phi \Vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ ,  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.
- $\nabla_{\forall}$  Suppose  $(\Pi^\alpha \mathbf{G}) \in \Phi$  and  $\Phi * (\mathbf{G}\mathbf{A})$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(\neg I)$ ,  $\Phi \Vdash \neg(\mathbf{G}\mathbf{A})$ . By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\Pi E)$ ,  $\Phi \Vdash \mathbf{G}\mathbf{A}$ . Finally,  $\mathfrak{N}\mathcal{K}(\neg E)$  implies  $\Phi \Vdash \mathbf{F}_o$ .
- $\nabla_{\exists}$  Suppose  $\neg(\Pi^\alpha \mathbf{G}) \in \Phi$ ,  $w_\alpha$  is a parameter which does not occur in  $\Phi$ , and  $\Phi * \neg(\mathbf{G}w)$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$ ,  $\Phi \Vdash \mathbf{G}w$ .

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\vee E)$ , we have  $\Phi \vdash \mathbf{F}_o$ .
- $\nabla_{\wedge}$  Suppose  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and  $\Phi * \neg\mathbf{A} * \neg\mathbf{B}$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_R)$ , we have  $\Phi, \neg\mathbf{A} \vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , we have  $\Phi, \neg\mathbf{A} \vdash \mathbf{F}_o$ . By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_L)$ , we have  $\Phi \vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ ,  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.
- $\nabla_{\neg}$  Suppose  $(\Pi^\alpha \mathbf{G}) \in \Phi$  and  $\Phi * (\mathbf{G}\mathbf{A})$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(\neg I)$ ,  $\Phi \vdash \neg(\mathbf{G}\mathbf{A})$ . By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\Pi E)$ ,  $\Phi \vdash \mathbf{G}\mathbf{A}$ . Finally,  $\mathfrak{N}\mathcal{K}(\neg E)$  implies  $\Phi \vdash \mathbf{F}_o$ .
- $\nabla_{\exists}$  Suppose  $\neg(\Pi^\alpha \mathbf{G}) \in \Phi$ ,  $w_\alpha$  is a parameter which does not occur in  $\Phi$ , and  $\Phi * \neg(\mathbf{G}w)$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$ ,  $\Phi \vdash \mathbf{G}w$ . By  $\mathfrak{N}\mathcal{K}(III)^w$ ,  $\Phi \vdash (\Pi^\alpha \mathbf{G})$ .

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_{\vee}$  Suppose  $(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and both  $\Phi * \mathbf{A}$  and  $\Phi * \mathbf{B}$  are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\vee E)$ , we have  $\Phi \vdash \mathbf{F}_o$ .
- $\nabla_{\wedge}$  Suppose  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$  and  $\Phi * \neg\mathbf{A} * \neg\mathbf{B}$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_R)$ , we have  $\Phi, \neg\mathbf{A} \vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ , we have  $\Phi, \neg\mathbf{A} \vdash \mathbf{F}_o$ . By  $\mathfrak{N}\mathcal{K}(Contr)$  and  $\mathfrak{N}\mathcal{K}(\vee I_L)$ , we have  $\Phi \vdash \mathbf{A} \vee \mathbf{B}$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\mathbf{A} \vee \mathbf{B}) \in \Phi$ ,  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.
- $\nabla_{\neg}$  Suppose  $(\Pi^\alpha \mathbf{G}) \in \Phi$  and  $\Phi * (\mathbf{G}\mathbf{A})$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(\neg I)$ ,  $\Phi \vdash \neg(\mathbf{G}\mathbf{A})$ . By  $\mathfrak{N}\mathcal{K}(Hyp)$  and  $\mathfrak{N}\mathcal{K}(\Pi E)$ ,  $\Phi \vdash \mathbf{G}\mathbf{A}$ . Finally,  $\mathfrak{N}\mathcal{K}(\neg E)$  implies  $\Phi \vdash \mathbf{F}_o$ .
- $\nabla_{\exists}$  Suppose  $\neg(\Pi^\alpha \mathbf{G}) \in \Phi$ ,  $w_\alpha$  is a parameter which does not occur in  $\Phi$ , and  $\Phi * \neg(\mathbf{G}w)$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. By  $\mathfrak{N}\mathcal{K}(Contr)$ ,  $\Phi \vdash \mathbf{G}w$ . By  $\mathfrak{N}\mathcal{K}(III)^w$ ,  $\Phi \vdash (\Pi^\alpha \mathbf{G})$ . Using  $\mathfrak{N}\mathcal{K}(\neg E)$  with  $\neg(\Pi^\alpha \mathbf{G}) \in \Phi$ ,  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.



# Lemma: Saturated $\mathcal{Acc}_*$



# Lemma: Saturated $\mathcal{A}cc_*$



$\nabla_{\text{sat}}$  Let  $\Phi * A$  and  $\Phi * \neg A$  be  $\mathcal{N}K_*$ -inconsistent.

# Lemma: Saturated $\mathcal{Acc}_*$



$\nabla_{\text{sat}}$  Let  $\phi * A$  and  $\phi * \neg A$  be  $\mathcal{N}\mathcal{K}_*$ -inconsistent. We show that  $\phi$  is  $\mathcal{N}\mathcal{K}_*$ -inconsistent.

# Lemma: Saturated $\mathcal{A}cc_*$



$\nabla_{\text{sat}}$  Let  $\Phi * A$  and  $\Phi * \neg A$  be  $\mathcal{N}K_*$ -inconsistent. We show that  $\Phi$  is  $\mathcal{N}K_*$ -inconsistent. Using  $\mathcal{N}K(\neg I)$ , we know  $\Phi \Vdash \neg A$  and  $\Phi \Vdash \neg\neg A$ .

# Lemma: Saturated $\mathcal{A}cc_*$

$\nabla_{\text{sat}}$  Let  $\Phi * \mathbf{A}$  and  $\Phi * \neg \mathbf{A}$  be  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. We show that  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. Using  $\mathfrak{N}\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg \mathbf{A}$  and  $\Phi \Vdash \neg \neg \mathbf{A}$ . By  $\mathfrak{N}\mathcal{K}(\neg E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .

# Lemma: Saturated $\mathcal{Acc}_*$

$\nabla_{\text{sat}}$  Let  $\Phi * \mathbf{A}$  and  $\Phi * \neg \mathbf{A}$  be  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. We show that  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. Using  $\mathfrak{N}\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg \mathbf{A}$  and  $\Phi \Vdash \neg \neg \mathbf{A}$ . By  $\mathfrak{N}\mathcal{K}(\neg E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .

Thus we have shown that  $\Gamma_{\Sigma}^{\beta}$  is saturated and in  $\mathcal{Acc}_{\beta}$ .

# Lemma: Saturated $\mathcal{Acc}_*$

$\nabla_{\text{sat}}$  Let  $\Phi * \mathbf{A}$  and  $\Phi * \neg \mathbf{A}$  be  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. We show that  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. Using  $\mathfrak{N}\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg \mathbf{A}$  and  $\Phi \Vdash \neg \neg \mathbf{A}$ . By  $\mathfrak{N}\mathcal{K}(\neg E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .

Thus we have shown that  $\Gamma_\Sigma^\beta$  is saturated and in  $\mathcal{Acc}_\beta$ .

Now let us check the conditions for the additional properties  $\eta$ ,  $\xi$ ,  $\mathfrak{f}$ , and  $\mathfrak{b}$ .

# Lemma: Saturated $\mathcal{Acc}_*$

$\nabla_{\text{sat}}$  Let  $\Phi * \mathbf{A}$  and  $\Phi * \neg \mathbf{A}$  be  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. We show that  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. Using  $\mathfrak{N}\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg \mathbf{A}$  and  $\Phi \Vdash \neg \neg \mathbf{A}$ . By  $\mathfrak{N}\mathcal{K}(\neg E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .

Thus we have shown that  $\Gamma_\Sigma^\beta$  is saturated and in  $\mathcal{Acc}_\beta$ .

Now let us check the conditions for the additional properties  $\eta$ ,  $\xi$ ,  $\mathfrak{f}$ , and  $\mathfrak{b}$ .



# Lemma: Saturated $\mathcal{Acc}_*$

$\nabla_{\text{sat}}$  Let  $\Phi * \mathbf{A}$  and  $\Phi * \neg \mathbf{A}$  be  $\mathcal{N}\mathcal{K}_*$ -inconsistent. We show that  $\Phi$  is  $\mathcal{N}\mathcal{K}_*$ -inconsistent. Using  $\mathcal{N}\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg \mathbf{A}$  and  $\Phi \Vdash \neg \neg \mathbf{A}$ . By  $\mathcal{N}\mathcal{K}(\neg E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .

Thus we have shown that  $\Gamma_\Sigma^\beta$  is saturated and in  $\mathcal{Acc}_\beta$ .

Now let us check the conditions for the additional properties  $\eta$ ,  $\xi$ ,  $\mathfrak{f}$ , and  $\mathfrak{b}$ .

$\nabla_\eta$  If  $*$  includes  $\eta$ , then the proof proceeds as in  $\nabla_\beta$  above, but with the rule  $\mathcal{N}\mathcal{K}(\eta)$ .

# Lemma: Saturated $\mathcal{Acc}_*$

$\nabla_{\text{sat}}$  Let  $\Phi * \mathbf{A}$  and  $\Phi * \neg \mathbf{A}$  be  $\mathcal{N}\mathcal{K}_*$ -inconsistent. We show that  $\Phi$  is  $\mathcal{N}\mathcal{K}_*$ -inconsistent. Using  $\mathcal{N}\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg \mathbf{A}$  and  $\Phi \Vdash \neg \neg \mathbf{A}$ . By  $\mathcal{N}\mathcal{K}(\neg E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .

Thus we have shown that  $\Gamma_\Sigma^\beta$  is saturated and in  $\mathcal{Acc}_\beta$ .

Now let us check the conditions for the additional properties  $\eta$ ,  $\xi$ ,  $\mathfrak{f}$ , and  $\mathfrak{b}$ .

$\nabla_\eta$  If  $*$  includes  $\eta$ , then the proof proceeds as in  $\nabla_\beta$  above, but with the rule  $\mathcal{N}\mathcal{K}(\eta)$ .

# Lemma: Saturated $\mathcal{Acc}_*$

$\nabla_{\text{sat}}$  Let  $\Phi * \mathbf{A}$  and  $\Phi * \neg \mathbf{A}$  be  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. We show that  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. Using  $\mathfrak{N}\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg \mathbf{A}$  and  $\Phi \Vdash \neg \neg \mathbf{A}$ . By  $\mathfrak{N}\mathcal{K}(\neg E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .

Thus we have shown that  $\Gamma_\Sigma^\beta$  is saturated and in  $\mathcal{Acc}_\beta$ .

Now let us check the conditions for the additional properties  $\eta$ ,  $\xi$ ,  $\mathfrak{f}$ , and  $\mathfrak{b}$ .

$\nabla_\eta$  If  $*$  includes  $\eta$ , then the proof proceeds as in  $\nabla_\beta$  above, but with the rule  $\mathfrak{N}\mathcal{K}(\eta)$ .

$\nabla_\xi$  Suppose  $*$  includes  $\xi$ ,  $\neg(\lambda X. \mathbf{M} \dot{=}^{\alpha \rightarrow \beta} \lambda X. \mathbf{N}) \in \Phi$ , and  $\Phi * \neg([\mathbf{w}/X] \mathbf{M} \dot{=}^\beta [\mathbf{w}/X] \mathbf{N})$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent for some parameter  $\mathbf{w}_\alpha$  which does not occur in any sentence of  $\Phi$ .

# Lemma: Saturated $\mathcal{Acc}_*$

$\nabla_{\text{sat}}$  Let  $\Phi * \mathbf{A}$  and  $\Phi * \neg \mathbf{A}$  be  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. We show that  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. Using  $\mathfrak{N}\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg \mathbf{A}$  and  $\Phi \Vdash \neg \neg \mathbf{A}$ . By  $\mathfrak{N}\mathcal{K}(\neg E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .

Thus we have shown that  $\Gamma_{\Sigma}^{\beta}$  is saturated and in  $\mathcal{Acc}_{\beta}$ .

Now let us check the conditions for the additional properties  $\eta$ ,  $\xi$ ,  $\mathfrak{f}$ , and  $\mathfrak{b}$ .

$\nabla_{\eta}$  If  $*$  includes  $\eta$ , then the proof proceeds as in  $\nabla_{\beta}$  above, but with the rule  $\mathfrak{N}\mathcal{K}(\eta)$ .

$\nabla_{\xi}$  Suppose  $*$  includes  $\xi$ ,  $\neg(\lambda X. \mathbf{M} \dot{=}^{\alpha \rightarrow \beta} \lambda X. \mathbf{N}) \in \Phi$ , and  $\Phi * \neg([\mathbf{w}/\mathbf{X}]\mathbf{M} \dot{=}^{\beta} [\mathbf{w}/\mathbf{X}]\mathbf{N})$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent for some parameter  $w_{\alpha}$  which does not occur in any sentence of  $\Phi$ . By  $\mathfrak{N}\mathcal{K}(\text{Contr})$ , we have  $\Phi \Vdash ([\mathbf{w}/\mathbf{X}]\mathbf{M} \dot{=}^{\beta} [\mathbf{w}/\mathbf{X}]\mathbf{N})$ .

# Lemma: Saturated $\mathcal{Acc}_*$

$\nabla_{\text{sat}}$  Let  $\Phi * \mathbf{A}$  and  $\Phi * \neg \mathbf{A}$  be  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. We show that  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. Using  $\mathfrak{N}\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg \mathbf{A}$  and  $\Phi \Vdash \neg \neg \mathbf{A}$ . By  $\mathfrak{N}\mathcal{K}(\neg E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .

Thus we have shown that  $\Gamma_\Sigma^\beta$  is saturated and in  $\mathcal{Acc}_\beta$ .

Now let us check the conditions for the additional properties  $\eta$ ,  $\xi$ ,  $\mathfrak{f}$ , and  $\mathfrak{b}$ .

$\nabla_\eta$  If  $*$  includes  $\eta$ , then the proof proceeds as in  $\nabla_\beta$  above, but with the rule  $\mathfrak{N}\mathcal{K}(\eta)$ .

$\nabla_\xi$  Suppose  $*$  includes  $\xi$ ,  $\neg(\lambda X. \mathbf{M} \dot{=}^{\alpha \rightarrow \beta} \lambda X. \mathbf{N}) \in \Phi$ , and  $\Phi * \neg([\mathbf{w}/\mathbf{X}]\mathbf{M} \dot{=}^\beta [\mathbf{w}/\mathbf{X}]\mathbf{N})$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent for some parameter  $w_\alpha$  which does not occur in any sentence of  $\Phi$ . By  $\mathfrak{N}\mathcal{K}(\text{Contr})$ , we have  $\Phi \Vdash ([\mathbf{w}/\mathbf{X}]\mathbf{M} \dot{=}^\beta [\mathbf{w}/\mathbf{X}]\mathbf{N})$ . By  $\mathfrak{N}\mathcal{K}(\beta)$ , we have  $\Phi \Vdash ((\lambda X. \mathbf{M} \dot{=}^\beta \mathbf{N})\mathbf{w})$ .

# Lemma: Saturated $\mathcal{Acc}_*$

$\nabla_{\text{sat}}$  Let  $\Phi * \mathbf{A}$  and  $\Phi * \neg \mathbf{A}$  be  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. We show that  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. Using  $\mathfrak{N}\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg \mathbf{A}$  and  $\Phi \Vdash \neg \neg \mathbf{A}$ . By  $\mathfrak{N}\mathcal{K}(\neg E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .

Thus we have shown that  $\Gamma_{\Sigma}^{\beta}$  is saturated and in  $\mathcal{Acc}_{\beta}$ .

Now let us check the conditions for the additional properties  $\eta$ ,  $\xi$ ,  $\mathfrak{f}$ , and  $\mathfrak{b}$ .

$\nabla_{\eta}$  If  $*$  includes  $\eta$ , then the proof proceeds as in  $\nabla_{\beta}$  above, but with the rule  $\mathfrak{N}\mathcal{K}(\eta)$ .

$\nabla_{\xi}$  Suppose  $*$  includes  $\xi$ ,  $\neg(\lambda X.M \dot{=}^{\alpha \rightarrow \beta} \lambda X.N) \in \Phi$ , and  $\Phi * \neg([w/X]M \dot{=}^{\beta} [w/X]N)$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent for some parameter  $w_{\alpha}$  which does not occur in any sentence of  $\Phi$ . By  $\mathfrak{N}\mathcal{K}(Contr)$ , we have  $\Phi \Vdash ([w/X]M \dot{=}^{\beta} [w/X]N)$ . By  $\mathfrak{N}\mathcal{K}(\beta)$ , we have  $\Phi \Vdash ((\lambda X.M \dot{=}^{\beta} N)w)$ . By  $\mathfrak{N}\mathcal{K}(III)$ ,  $\Phi \Vdash (\forall X.M \dot{=}^{\beta} N)$ .

# Lemma: Saturated $\mathcal{Acc}_*$

$\nabla_{\text{sat}}$  Let  $\Phi * \mathbf{A}$  and  $\Phi * \neg \mathbf{A}$  be  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. We show that  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. Using  $\mathfrak{N}\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg \mathbf{A}$  and  $\Phi \Vdash \neg \neg \mathbf{A}$ . By  $\mathfrak{N}\mathcal{K}(\neg E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .

Thus we have shown that  $\Gamma_\Sigma^\beta$  is saturated and in  $\mathcal{Acc}_\beta$ .

Now let us check the conditions for the additional properties  $\eta$ ,  $\xi$ ,  $\mathfrak{f}$ , and  $\mathfrak{b}$ .

$\nabla_\eta$  If  $*$  includes  $\eta$ , then the proof proceeds as in  $\nabla_\beta$  above, but with the rule  $\mathfrak{N}\mathcal{K}(\eta)$ .

$\nabla_\xi$  Suppose  $*$  includes  $\xi$ ,  $\neg(\lambda X.M \dot{=}^{\alpha \rightarrow \beta} \lambda X.N) \in \Phi$ , and  $\Phi * \neg([w/X]M \dot{=}^\beta [w/X]N)$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent for some parameter  $w_\alpha$  which does not occur in any sentence of  $\Phi$ . By  $\mathfrak{N}\mathcal{K}(\text{Contr})$ , we have  $\Phi \Vdash ([w/X]M \dot{=}^\beta [w/X]N)$ . By  $\mathfrak{N}\mathcal{K}(\beta)$ , we have  $\Phi \Vdash ((\lambda X.M \dot{=}^\beta N)w)$ . By  $\mathfrak{N}\mathcal{K}(III)$ ,  $\Phi \Vdash (\forall X.M \dot{=}^\beta N)$ . By  $\mathfrak{N}\mathcal{K}(\xi)$ ,  $\Phi \Vdash (\lambda X.M \dot{=}^{\alpha \rightarrow \beta} \lambda X.N)$ .

# Lemma: Saturated $\mathcal{Acc}_*$

$\nabla_{\text{sat}}$  Let  $\Phi * \mathbf{A}$  and  $\Phi * \neg \mathbf{A}$  be  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. We show that  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent. Using  $\mathfrak{N}\mathcal{K}(\neg I)$ , we know  $\Phi \Vdash \neg \mathbf{A}$  and  $\Phi \Vdash \neg \neg \mathbf{A}$ . By  $\mathfrak{N}\mathcal{K}(\neg E)$ , we have  $\Phi \Vdash \mathbf{F}_o$ .

Thus we have shown that  $\Gamma_\Sigma^\beta$  is saturated and in  $\mathcal{Acc}_\beta$ .

Now let us check the conditions for the additional properties  $\eta$ ,  $\xi$ ,  $\mathfrak{f}$ , and  $\mathfrak{b}$ .

$\nabla_\eta$  If  $*$  includes  $\eta$ , then the proof proceeds as in  $\nabla_\beta$  above, but with the rule  $\mathfrak{N}\mathcal{K}(\eta)$ .

$\nabla_\xi$  Suppose  $*$  includes  $\xi$ ,  $\neg(\lambda X.M \dot{=}^{\alpha \rightarrow \beta} \lambda X.N) \in \Phi$ , and  $\Phi * \neg([w/X]M \dot{=}^\beta [w/X]N)$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent for some parameter  $w_\alpha$  which does not occur in any sentence of  $\Phi$ . By  $\mathfrak{N}\mathcal{K}(\text{Contr})$ , we have  $\Phi \Vdash ([w/X]M \dot{=}^\beta [w/X]N)$ . By  $\mathfrak{N}\mathcal{K}(\beta)$ , we have  $\Phi \Vdash ((\lambda X.M \dot{=}^\beta N)w)$ . By  $\mathfrak{N}\mathcal{K}(III)$ ,  $\Phi \Vdash (\forall X.M \dot{=}^\beta N)$ . By  $\mathfrak{N}\mathcal{K}(\xi)$ ,  $\Phi \Vdash (\lambda X.M \dot{=}^{\alpha \rightarrow \beta} \lambda X.N)$ . By  $\mathfrak{N}\mathcal{K}(\neg E)$ ,  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.



# Lemma: Saturated $\mathcal{A}cc_*$



$\nabla_f$  This case is analogous to the previous one, generalizing  $\lambda X.M \doteq \lambda X.N$  to arbitrary  $G \doteq H$  and using the extensionality rule  $\mathfrak{N}\mathcal{R}(f)$  instead of  $\mathfrak{N}\mathcal{R}(\xi)$ .

# Lemma: Saturated $\mathcal{A}cc_*$



$\nabla_f$  This case is analogous to the previous one, generalizing  $\lambda X.M \doteq \lambda X.N$  to arbitrary  $G \doteq H$  and using the extensionality rule  $\mathfrak{N}\mathcal{R}(f)$  instead of  $\mathfrak{N}\mathcal{R}(\xi)$ .

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_f$  This case is analogous to the previous one, generalizing  $\lambda X.M \doteq \lambda X.N$  to arbitrary  $G \doteq H$  and using the extensionality rule  $\mathfrak{N}\mathfrak{E}(f)$  instead of  $\mathfrak{N}\mathfrak{E}(\xi)$ .
- $\nabla_b$  Suppose  $*$  includes  $b$ .

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_f$  This case is analogous to the previous one, generalizing  $\lambda X.M \doteq \lambda X.N$  to arbitrary  $G \doteq H$  and using the extensionality rule  $\mathfrak{R}(f)$  instead of  $\mathfrak{R}(\xi)$ .
- $\nabla_b$  Suppose  $*$  includes  $b$ . Assume that  $\neg(A \doteq^o B) \in \Phi$  but both  $\Phi * \neg A * B \notin \Gamma_\Sigma^*$  and  $\Phi * A * \neg B \notin \Gamma_\Sigma^*$ .

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_f$  This case is analogous to the previous one, generalizing  $\lambda X.M \doteq \lambda X.N$  to arbitrary  $G \doteq H$  and using the extensionality rule  $\mathfrak{N}\mathcal{K}(f)$  instead of  $\mathfrak{N}\mathcal{K}(\xi)$ .
- $\nabla_b$  Suppose  $*$  includes  $b$ . Assume that  $\neg(A \doteq^\circ B) \in \Phi$  but both  $\Phi * \neg A * B \notin \Gamma_\Sigma^*$  and  $\Phi * A * \neg B \notin \Gamma_\Sigma^*$ . So both are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent and we have  $\Phi * A \Vdash B$  and  $\Phi * B \Vdash A$  by  $\mathfrak{N}\mathcal{K}(Contr)$ .

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_f$  This case is analogous to the previous one, generalizing  $\lambda X.M \doteq \lambda X.N$  to arbitrary  $G \doteq H$  and using the extensionality rule  $\mathfrak{N}\mathcal{K}(f)$  instead of  $\mathfrak{N}\mathcal{K}(\xi)$ .
- $\nabla_b$  Suppose  $*$  includes  $b$ . Assume that  $\neg(A \doteq^\circ B) \in \Phi$  but both  $\Phi * \neg A * B \notin \Gamma_\Sigma^*$  and  $\Phi * A * \neg B \notin \Gamma_\Sigma^*$ . So both are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent and we have  $\Phi * A \Vdash B$  and  $\Phi * B \Vdash A$  by  $\mathfrak{N}\mathcal{K}(Contr)$ . By  $\mathfrak{N}\mathcal{K}(b)$ , we have  $\Phi \Vdash (A \doteq^\circ B)$ .

# Lemma: Saturated $\mathcal{A}cc_*$

- $\nabla_f$  This case is analogous to the previous one, generalizing  $\lambda X.M \doteq \lambda X.N$  to arbitrary  $G \doteq H$  and using the extensionality rule  $\mathfrak{N}\mathcal{K}(f)$  instead of  $\mathfrak{N}\mathcal{K}(\xi)$ .
- $\nabla_b$  Suppose  $*$  includes  $b$ . Assume that  $\neg(A \doteq^\circ B) \in \Phi$  but both  $\Phi * \neg A * B \notin \Gamma_\Sigma^*$  and  $\Phi * A * \neg B \notin \Gamma_\Sigma^*$ . So both are  $\mathfrak{N}\mathcal{K}_*$ -inconsistent and we have  $\Phi * A \Vdash B$  and  $\Phi * B \Vdash A$  by  $\mathfrak{N}\mathcal{K}(Contr)$ . By  $\mathfrak{N}\mathcal{K}(b)$ , we have  $\Phi \Vdash (A \doteq^\circ B)$ . Since  $\neg(A \doteq^\circ B) \in \Phi$ ,  $\Phi$  is  $\mathfrak{N}\mathcal{K}_*$ -inconsistent.

# Thm.: Henkin's Theorem for $\mathfrak{N}_*$

Let  $*$   $\in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$ . Every sufficiently  $\Sigma$ -pure  $\mathfrak{N}_*$ -consistent set of sentences has an  $\mathfrak{M}_*(\Sigma)$ -model.

Proof:



# Thm.: Henkin's Theorem for $\mathcal{N}\mathcal{K}_*$

Let  $*$   $\in \{\beta, \beta\eta, \beta\xi, \beta\mathfrak{f}, \beta\mathfrak{b}, \beta\eta\mathfrak{b}, \beta\xi\mathfrak{b}, \beta\mathfrak{f}\mathfrak{b}\}$ . Every sufficiently  $\Sigma$ -pure  $\mathcal{N}\mathcal{K}_*$ -consistent set of sentences has an  $\mathfrak{M}_*(\Sigma)$ -model.

Proof: Let  $\Phi$  be a sufficiently  $\Sigma$ -pure  $\mathcal{N}\mathcal{K}_*$ -consistent set of sentences.

# Thm.: Henkin's Theorem for $\mathcal{N}\mathcal{K}_*$

Let  $*$   $\in \{\beta, \beta\eta, \beta\xi, \beta\mathfrak{f}, \beta\mathfrak{b}, \beta\eta\mathfrak{b}, \beta\xi\mathfrak{b}, \beta\mathfrak{f}\mathfrak{b}\}$ . Every sufficiently  $\Sigma$ -pure  $\mathcal{N}\mathcal{K}_*$ -consistent set of sentences has an  $\mathfrak{M}_*(\Sigma)$ -model.

Proof: Let  $\Phi$  be a sufficiently  $\Sigma$ -pure  $\mathcal{N}\mathcal{K}_*$ -consistent set of sentences. By the previous lemma we know that the class of sets of  $\mathcal{N}\mathcal{K}_*$ -consistent sentences constitute a saturated  $\mathcal{A}cc_*$ ,

# Thm.: Henkin's Theorem for $\mathcal{N}\mathcal{K}_*$

Let  $* \in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$ . Every sufficiently  $\Sigma$ -pure  $\mathcal{N}\mathcal{K}_*$ -consistent set of sentences has an  $\mathfrak{M}_*(\Sigma)$ -model.

Proof: Let  $\Phi$  be a sufficiently  $\Sigma$ -pure  $\mathcal{N}\mathcal{K}_*$ -consistent set of sentences. By the previous lemma we know that the class of sets of  $\mathcal{N}\mathcal{K}_*$ -consistent sentences constitute a saturated  $\mathcal{A}cc_*$ , thus the Model Existence Theorem guarantees an  $\mathfrak{M}_*(\Sigma)$  model for  $\Phi$ .

# Thm.: Completeness Theorem for $\mathfrak{N}\mathfrak{K}_*$



Let  $\Phi$  be a sufficiently  $\Sigma$ -pure set of sentences,  $A$  be a sentence, and  $*$   $\in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$ . If  $A$  is valid in all models  $\mathcal{M} \in \mathfrak{M}_*(\Sigma)$  that satisfy  $\Phi$ , then  $\Phi \Vdash_{\mathfrak{N}\mathfrak{K}_*} A$ .

Proof:

# Thm.: Completeness Theorem for $\mathfrak{N}\mathfrak{K}_*$



Let  $\Phi$  be a sufficiently  $\Sigma$ -pure set of sentences,  $A$  be a sentence, and  $*$   $\in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$ . If  $A$  is valid in all models  $\mathcal{M} \in \mathfrak{M}_*(\Sigma)$  that satisfy  $\Phi$ , then  $\Phi \Vdash_{\mathfrak{N}\mathfrak{K}_*} A$ .

Proof: Let  $A$  be given such that  $A$  is valid in all  $\mathfrak{M}_*(\Sigma)$  models that satisfy  $\Phi$ .

# Thm.: Completeness Theorem for $\mathfrak{N}\mathfrak{K}_*$



Let  $\Phi$  be a sufficiently  $\Sigma$ -pure set of sentences,  $A$  be a sentence, and  $*$   $\in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$ . If  $A$  is valid in all models  $\mathcal{M} \in \mathfrak{M}_*(\Sigma)$  that satisfy  $\Phi$ , then  $\Phi \Vdash_{\mathfrak{N}\mathfrak{K}_*} A$ .

Proof: Let  $A$  be given such that  $A$  is valid in all  $\mathfrak{M}_*(\Sigma)$  models that satisfy  $\Phi$ . So,  $\Phi * \neg A$  is unsatisfiable in  $\mathfrak{M}_*(\Sigma)$ . Since only finitely many constants occur in  $\neg A$ ,  $\Phi * \neg A$  is sufficiently  $\Sigma$ -pure.

# Thm.: Completeness Theorem for $\mathfrak{N}\mathfrak{K}_*$



Let  $\Phi$  be a sufficiently  $\Sigma$ -pure set of sentences,  $A$  be a sentence, and  $*$   $\in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$ . If  $A$  is valid in all models  $\mathcal{M} \in \mathfrak{M}_*(\Sigma)$  that satisfy  $\Phi$ , then  $\Phi \Vdash_{\mathfrak{N}\mathfrak{K}_*} A$ .

Proof: Let  $A$  be given such that  $A$  is valid in all  $\mathfrak{M}_*(\Sigma)$  models that satisfy  $\Phi$ . So,  $\Phi * \neg A$  is unsatisfiable in  $\mathfrak{M}_*(\Sigma)$ . Since only finitely many constants occur in  $\neg A$ ,  $\Phi * \neg A$  is sufficiently  $\Sigma$ -pure. So,  $\Phi * \neg A$  must be  $\mathfrak{N}\mathfrak{K}_*$ -inconsistent by Henkin's theorem above.

# Thm.: Completeness Theorem for $\mathfrak{N}\mathfrak{K}_*$



Let  $\Phi$  be a sufficiently  $\Sigma$ -pure set of sentences,  $A$  be a sentence, and  $*$   $\in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$ . If  $A$  is valid in all models  $\mathcal{M} \in \mathfrak{M}_*(\Sigma)$  that satisfy  $\Phi$ , then  $\Phi \Vdash_{\mathfrak{N}\mathfrak{K}_*} A$ .

Proof: Let  $A$  be given such that  $A$  is valid in all  $\mathfrak{M}_*(\Sigma)$  models that satisfy  $\Phi$ . So,  $\Phi * \neg A$  is unsatisfiable in  $\mathfrak{M}_*(\Sigma)$ . Since only finitely many constants occur in  $\neg A$ ,  $\Phi * \neg A$  is sufficiently  $\Sigma$ -pure. So,  $\Phi * \neg A$  must be  $\mathfrak{N}\mathfrak{K}_*$ -inconsistent by Henkin's theorem above. Thus,  $\Phi \Vdash_{\mathfrak{N}\mathfrak{K}_*} A$  by  $\mathfrak{N}\mathfrak{K}(Contr)$ .



We can use the completeness theorems obtained so far to prove a compactness theorem for our semantics:

Let  $\Phi$  be a sufficiently  $\Sigma$ -pure set of sentences and  $* \in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$ .  $\Phi$  has an  $\mathfrak{M}_*(\Sigma)$ -model iff every finite subset of  $\Phi$  has an  $\mathfrak{M}_*(\Sigma)$ -model.

Proof:

We can use the completeness theorems obtained so far to prove a compactness theorem for our semantics:

Let  $\Phi$  be a sufficiently  $\Sigma$ -pure set of sentences and  $* \in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$ .  $\Phi$  has an  $\mathfrak{M}_*(\Sigma)$ -model iff every finite subset of  $\Phi$  has an  $\mathfrak{M}_*(\Sigma)$ -model.

Proof: If  $\Phi$  has no  $\mathfrak{M}_*(\Sigma)$ -model, then by the previous Henkin Theorem  $\Phi$  is  $\mathfrak{N}_*$ -inconsistent.

We can use the completeness theorems obtained so far to prove a compactness theorem for our semantics:

Let  $\Phi$  be a sufficiently  $\Sigma$ -pure set of sentences and  $* \in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$ .  $\Phi$  has an  $\mathfrak{M}_*(\Sigma)$ -model iff every finite subset of  $\Phi$  has an  $\mathfrak{M}_*(\Sigma)$ -model.

Proof: If  $\Phi$  has no  $\mathfrak{M}_*(\Sigma)$ -model, then by the previous Henkin Theorem  $\Phi$  is  $\mathfrak{N}\mathfrak{K}_*$ -inconsistent. Since every  $\mathfrak{N}\mathfrak{K}_*$ -proof is finite, this means some finite subset  $\Psi$  of  $\Phi$  is  $\mathfrak{N}\mathfrak{K}_*$ -inconsistent.

We can use the completeness theorems obtained so far to prove a compactness theorem for our semantics:

Let  $\Phi$  be a sufficiently  $\Sigma$ -pure set of sentences and  $* \in \{\beta, \beta\eta, \beta\xi, \beta f, \beta b, \beta\eta b, \beta\xi b, \beta f b\}$ .  $\Phi$  has an  $\mathfrak{M}_*(\Sigma)$ -model iff every finite subset of  $\Phi$  has an  $\mathfrak{M}_*(\Sigma)$ -model.

Proof: If  $\Phi$  has no  $\mathfrak{M}_*(\Sigma)$ -model, then by the previous Henkin Theorem  $\Phi$  is  $\mathfrak{N}\mathfrak{K}_*$ -inconsistent. Since every  $\mathfrak{N}\mathfrak{K}_*$ -proof is finite, this means some finite subset  $\Psi$  of  $\Phi$  is  $\mathfrak{N}\mathfrak{K}_*$ -inconsistent. Hence,  $\Psi$  has no  $\mathfrak{M}_*(\Sigma)$ -model.